

# Analysis of Cesarean Section Rate using Robson 10 Group Classification System in a Tertiary Hospital: An Observational Study

Dr. Abhilasha Yadav<sup>1</sup>, Dr. Rachna Agrawal<sup>2</sup>, Dr. K Romila Chawang<sup>3</sup> and Dr. Ruchika Garg<sup>4</sup>

<sup>1</sup> Sarojini Naidu Medical College

Received: 5 June 2021 Accepted: 2 July 2021 Published: 15 July 2021

---

## Abstract

Background: The Cesarean Section rate has increased from 12

---

**Index terms**— cesarean section rate, robson's criteria, groups.

## 1 I.

background ver the last few decades, there has been a progressive increase in the rate of cesarean section deliveries in our country. WHO has recommended that this rate should be between 10%and 15% but the driver behind this trend is not completely understood especially in developing countries. 1,2 Immediate and long-term complications of CS included increased risk of maternal mortality and morbidity, increased need for blood transfusion, longer hospitalization, postpartum infections, retained placenta, stillbirths, postpartum hemorrhage.

Over the last few decades, there has been a progressive increase in the rate of cesarean section deliveries in our country. WHO has recommended that this rate should be between 10%and 15% but the driver behind this trend is not completely understood especially in developing countries. (1,2). The rising cesarean section trend is a major public health concern due to potential maternal and perinatal risks associated with this. Immediate and long-term complications of CS included increased risk of maternal mortality and morbidity, increased need for blood transfusion, longer hospitalization, postpartum infections, retained placenta, stillbirths, postpartum hemorrhage.

To understand this and to implement effective measures to reduce cesarean section rates, a tool is required to monitor and compare cesarean section rates in the same setting over time and among different settings. Traditionally, at the facility level, we monitor cesarean section rates using the overall percentage of deliveries by cesarean section. But because of some intrinsic differences in hospital factors and infrastructure, a difference in characteristics of the population, and differences in clinical management protocols, this "overall cesarean section rate" becomes difficult to interpret and compare. Ideally, there should be a classification system to monitor and compare cesarean section rates. Such a system should be simple, clinically relevant, accountable, replicable, and verifiable (3,4,5). In 2015, WHO introduced Robson's criteria in the year 2015 as a standardized method to determine the rate and indications of cesarean section and issued an implementation manual(6). ? Obstetric history (parity and previous cesarean section) ? The onset of labor (spontaneous, induced, or cesarean section before the onset of labor) ? Fetal presentation or lie(cephalic, breech, or transverse) ? Number of neonates ? Gestational age (preterm or term). Nulliparous women with a single cephalic pregnancy, ?37 weeks gestation in spontaneous labor 2 Nulliparous women with a single cephalic pregnancy, ?37 weeks gestation who had labor induced or were delivered by cesarean section before labor 2a Cesarean section performed after induction of labor 2b Cesarean section performed before onset of labor 3

## 2 O

Multiparous women without a previous cesarean section, with a single cephalic pregnancy, ?37 weeks gestation in spontaneous labor 4 Multiparous women without a previous cesarean section, with a single cephalic pregnancy,

## 9 LOOK AT THE CS RATE FOR GROUP3

---

44 ?37 weeks gestation who had labor induced or were delivered by cesarean before labor 4a Cesarean sections  
45 performed after induction of labor 4b Cesarean sections performed before onset of labor 5

46 All multiparous women with one or more previous cesarean sections, with a single cephalic pregnancy, ?37  
47 weeks gestation 5.1 With one previous cesarean section 5.2 With two or more previous cesarean sections 6

48 All nulliparous women with a single breech pregnancy 7 All multiparous women with a single breech pregnancy  
49 including women with previous cesarean section 8 All women with multiple pregnancies including women with  
50 previous cesarean section 9

51 All women with a single pregnancy with a transverse or oblique lie, including women with previous cesarean  
52 section 10

53 All women with a single cephalic pregnancy < 37 weeks gestation, including women with previous cesarean  
54 section II.

## 55 3 Materials and Methods

56 This study is a cross-sectional study of 5744 women delivered in the department of obstetrics of S.N. Medical  
57 College, Agra performed from 1 st April 2020 to 31 st March 2021. S.N.Medical College is a tertiary referral  
58 hospital where around 6000 deliveries take place annually.

59 All women who underwent cesarean section in the hospital during the specified period were included in the  
60 study.

61 Cases with incomplete information, doubtful gestational age, and laparotomy for uterine rupture were excluded  
62 from the study. The identity of women who underwent cesarean section was obtained from the delivery register,  
63 admission and discharge register, and operation register. The admission and discharge register and delivery  
64 register contained information about all women who delivered in the hospital regardless of the mode of delivery  
65 (vaginal, Caesarean section) while the operation register contained only information about women who underwent  
66 Caesarean section. Using the medical registration number of each woman, we accessed all Caesarean section  
67 files performed during the study.10 groups and their characteristics are shown in table 1. Gestational age was  
68 categorized as a term ( $\geq 37$  weeks) or preterm ( $< 37$  weeks). Patient's demographic data, patient's parity,  
69 obstetric history, the onset of labor, fetal presentation or lie, number of neonates, gestational age were collected.  
70 The indications for cesarean section were grouped using a flowchart (figure 1).

## 71 4 Result

72 To make the most of the information provided by the Robson Classification in local settings and to allow  
73 comparisons between settings, the data is best reported in a standardized way (the "Robson Classification Report  
74 Table ??) 7 12.9% Can be higher in tertiary hospitals as women with high-risk factors are being referred to our  
75 hospital. These women require induction of labor, so it is accompanied by a high rate of cesarean section in this  
76 group.

## 77 5 Look at the Ratio of the size of Group1 versus Group2

78 It is usually 2:1 or higher It is always higher than the ratio of Group1/Group2 in the same institution. This is a  
79 very reliable finding in confirming data quality and culture of the organization.

## 80 6 5.2

81 The Ratio of group3:group4 is larger than the ratio of group 1:group2 which strongly signifies the reliability of  
82 data 9. Look at the Ratio of the size of Group 6 and Group7.

83 The ratio of nulliparous breech/multiparous breech is usually 2:1 because breeches are more frequent in  
84 nulliparous women than in multiparous women. Ratio 0.94 It can be because of an unusual nullipara/multipara  
85 ratio.

86 C. Steps to assess the rate of cesarean section Table ??: Steps to assess cesarean section rates using the Robson  
87 Report Table

## 88 7 Steps

89 Robson guideline Result Interpretation

90 1. Look at the CS rate for Group1

91 Rates under 10% are achievable 12.23% It is slightly higher in our hospital.

## 92 8 Look at the CS rate for Group2

93 Consistently around 20-35%

94 33.83% It is within the appropriate range.

## 95 9 Look at the CS rate for Group3

96 Normally, no higher than 3.0%.

---

## 10 3.57%

Slightly higher than 3%

4. Look at the CS rate for Group 4 It rarely should be higher than 15%

22.39% CS rates in Group 4 reflect the size and rates in 4a and 4b. If the size of Group 4 is large (there are more cases of prelabour cesarean sections), the overall CS rate in Group 4 is also to be high.

## 11 Look at the CS rate

for Group 5

Rates of 50-60% are considered appropriate.

## 12 69.07%

Rates are higher, this is due to a large number of women with 2 or more previous CS.

## 13 Look at the CS rate for Group 8

It is usually around 60%.

## 14 38.02%

The lower rate can be due to more number of multiparous twins or due to nulliparous/multiparous with or without a previous scar.

## 15 Look at the CS rate in Group 10

In most populations, it is usually around 30%

## 16 Discussion

Cesarean sections are becoming more common over the world, with rates rising from 12% in 2000 to 21% in 2015. In India, the caesarean section rate has been steadily growing from 8.5 percent in 2005-2006 to 17.2 percent in 2015-2016, following global trends. Many authors have expressed their satisfaction with this classification and have suggested that it be utilised more widely. Madhav Prasad wrote in a 2015 article that now is an ideal moment for India to use the 10-group Robson's classification to rate caesarean sections. The bulk of caesarean sections were found in groups 2 and 5, according to Deepika Jamwal et al. Group 5 was responsible for 40.3 percent of all caesarean sections, while group 2 was responsible for 29.2%. At a study conducted in a private tertiary care centre in northwest India, Priyanka D. Jogia et al. discovered that group 5 (women with a history of CS) contributed the most (37 percent) to overall surgical deliveries, with group 2 being the second highest contributor (21 percent). Pratima et al conducted a study of 81,784 deliveries (62,336 vaginal and 19,448 Caesarean deliveries) over 3 years. The year-wise CS rate was 22.4%, 23.5% and 25.5%, respectively. The largest contributor was by group 5 followed by group 2 and group 1. Based on 3-year data, it was predicted in the study that the CS rate will increase by 0.905% annually in the coming 3 years. (15). During the study period, our institution's overall caesarean section rate was 26.22 percent, which looks to be higher than the national norm. Robson group 5 was the most significant single contribution to our institution's CS rate. Caesarean section rate in group 1 was higher than recommended i.e. 12.23% (rate <10% are achievable) and in group 3 CS rate was 3.57% against the recommended rate of <3%. In group 5 the caesarean section rate between 50-60% is recommended but in our institution CS rate in group 5 was 69% as a large number of women with previous 2 or more caesarean sections were being admitted. The absolute contribution of group 5 is very high (40%), this indicates high CS rates. The size of group 5 is frequently connected to the overall caesarean section rate. The size of group 5 in our study was 15%, which explains our institution's overall increased caesarean section rate. It's also worth noting that Robson group 10 has grown in size (preterm deliveries). Because women with high-risk conditions are referred to our hospital, Robson group 5 can be greater in tertiary hospitals. Induction of labour is necessary for these women, hence there is a high rate of caesarean section in this group. In group 1 and group 2 in previous years.

## 17 V.

## 18 Conclusion

Reducing the primary caesarean section rate in group 1 and group 2 will reduce the overall caesarean section rate in the institution.

## 19 Bibliography




Figure 1: Figure 1 :

1

Figure 2: Table 1 :

2

Group Number	Number of CS in group	Number of women in the group	Group Size1 (%)	Group CS rate2 (%)	Absolute contribution to overall CS rate3 (%)	The relative contribution of a group to overall CS rate4 (%)
1	214	1750	30.43	12.23	3.73	14.21
2	204	603	10.49	33.83	3.55	13.55
3	48	1346	23.43	3.57	0.84	3.19
4	58	259	4.51	22.39	1.01	3.85
5	603	873	15.12	69.07	10.50	40.04
6	60	77	1.34	77.92	1.04	3.98
7	36	82	1.43	43.90	0.63	2.39
8	27	71	1.24	38.02	0.47	1.79
9	39	42	0.73	92.86	0.68	2.59
10	217	741	12.90	29.28	3.78	14.41
Total* number CS 1506	Total number women delivered 5744	Total number delivered 5744	100%	Overall CS rate 26.22%	Overall CS rate 26.22%	100%

[Note: Unclassifiable: Number of cases and % [Number unclassifiable cases/(Total Number women delivered classified+unclassified)X100] \* These totals and percentages come from the data in the table.1. Group size(%)=no of women in the group/total no of women delivered in the hospital x 100 2. Group CS rate(%)=no of CS in the group/total no of women in the group x 100 3. Absolute contribution (%)=no of CS in the group/total no of women delivered in the hospital x 100 4. Relative contribution(%)=no of CS in the group/total no of CS in the hospital x 100 A. Steps to assess the quality of data]

Figure 3: Table 2 :

3

Steps	Robson guideline	Result	Interpretation
1. The total number of cesarean sections and women delivered in our hospital	These numbers should be the same as the total number of cesarean sections performed and of women delivered in the hospital.	57	We excluded the cases with incomplete data
2. Look at the size of Group 9 i.e. all women with single ton transverse or oblique lie.	It should be less than 1%.	0.73%	It is less than 1%
3. Look at the CS rate of Group 9	It should be 100% by convention.	92.86 %	extremely high pregnancies with intrauterine death of fetus were delivered vaginally
B. Steps to assess the type of population			

Figure 4: Table 3 :

4

Steps	Robson guideline	Result	Interpretation
1. Look at the of Group 1+	This usually represents	40.9%	It is within the acceptable range
2i.e. All nulliparous women ?37 weeks gestationsing letoncephalic	35-42% of the obstetric population hospitals.	of	most
2. Look at the size of Groups 3+4 i.e. All multiparous women ?37 weeks gestation cephalic, without previous CS	This usually represents about 30% of women. which is accompanied by a high overall CS torate.	26.2% The reason for low size of Groups 3 and 4 could be that the size of Group 5 is very high	
3. Look at the size of Group 5. (Multiparous women with previous cesarean section ?37 weeks gestation with singleton pregnancy).	It is related to the overall CS rate. Group 5 usually contributes to about half of the total CS rate. In settings with low overall CS rates it is usually under 10%.	15.12% Overall CS rate is usually related to the size of group 5 and the size of this group is larger (>15%) if the institute has high CS rate in the pasty ears mainly in Groups 1 and 2.	
4. Look at the size of Groups 6+7 i.e. Breeches in nulliparous women + breeches in multiparous women	It should be 3-4%	2.77%	It is within the acceptable range
5. Look at the size of Groups 8	It should be 1.5 -2%	1.43%	It is nearly within the acceptable range
6. Look at the size of Groups 9	It should be less than 5%		

Multiples



- 
- 145 [ WHO Statement on Caesarean Section Rates. Geneva: World Health Organization ()] , WHO/RHR/15.02.  
146 *WHO Statement on Caesarean Section Rates. Geneva: World Health Organization* 2015.
- 147 [Mittal (2019)] , Pratima Mittal . 10.1007/s13224-019-01275-7. *J ObstetGynaecol India* 2020 Apr. 2019 Oct 11.  
148 70 (2) p. .
- 149 [Jamwal (2021)] , D Jamwal . *Int J Reprod ContraceptObstet Gynecol* 2021 Jun. 10 (6) p. .
- 150 [Fesseha et al. ()] ‘A national review of cesarean delivery in Ethiopia’. N Fesseha , A Getachew , M Hiluf . *Int J*  
151 *GynaecolObstet* 2011. 115 p. .
- 152 [Betra´n et al. ()] ‘A systematic review of the Robson classification for cesarean section: what works, doesn’t  
153 work and how to improve it’. A P Betra´n , N Vindevoghel , J P Souza . 10.1371/journal.pone.0097769. *PLoS*  
154 *One* 2014. 9 (6) p. e97769. (PMC free article. Google Scholar)
- 155 [Priyanka and Jogia] *Analysis of cesarean sections according to modified Robson’s ten group*  
156 *classification system at a tertiary care center in Western India*, D Priyanka , Jogia .  
157 [https://scholar.google.com/scholar\\_lookup?journal=PLoS+One&title=Classifications+for+cesarean+section:+a+systematic-](https://scholar.google.com/scholar_lookup?journal=PLoS+One&title=Classifications+for+cesarean+section:+a+systematic-)  
158 <https://dx.doi.org/10.18203/2320-1770.ijrcog20190264>
- 159 [Torloni et al. ()] ‘Classifications for cesarean section: a systematic review’. M R Torloni , A P Betran , J P Souza  
160 . doi: 10.1371/ journal.pone.0014566. *PLoS One* 2011. 6 (1) p. e14566. (PMC free article. Google Scholar)
- 161 [CSA] [Ethiopia] nd ICF. Ethiopia Demographic and Health Survey ()b2 *CSA [Ethiopia] and ICF. Ethiopia*  
162 *Demographic and Health Survey*, 2016. 2016. Addis Ababa, Ethiopia, and Rockville, Maryland, USA. Central  
163 Statistical Agency (Ethiopia Demographic and Health Survey: CSA and ICF)
- 164 [Boeremat ()] ‘Global epidemiology of use of and disparities in cesarean sections’. Boeremat . *Lancet* 2018. 392 p. .
- 165 [Prasad (2015)] ‘It is a good time to adopt 10 group Robsons classification to rate cesarean sections in India’.  
166 Madhav Prasad . <https://dx.doi.org/10.1007%2> *J ObstetGynaecol India* 2015 Jul. 65 (4) p. .
- 167 [NFHS-4(National Family Health Survey-4)-2015-2016 -India fact sheet ()] *NFHS-4(National Family Health*  
168 *Survey-4)-2015-2016 -India fact sheet*, <http://rchiip.org/nfhs/pdf/NFHS4/TGFactSheet.pdf>  
169 2017. (International Institute for Population Sciences)
- 170 [Betrán et al. ()] ‘The increasing trend in cesarean section rates: global, regional and national estimates: 1990-  
171 2014’. A P Betrán , J Ye , A B Moller . *PLoS One* 2016. 11 p. e0148343.
- 172 [World Health Organization Human Reproduction Programme Reprod Health Matters (2015)] ‘World Health  
173 Organization Human Reproduction Programme’. *Reprod Health Matters* 10 April 2015. 2015. 23 p. . (WHO  
174 statement on cesarean section rates)