Online ISSN: 2249-4618
Print ISSN: 0975-5888

Global Journals

of Medical Research: G

Veterinary Science and Veterinary Medicine

Highlights

Cross-Sectional Survey

Bovine Fasciolosis at Elkadaro

Prevalence of Bovine Mastitis

Implications in Selected Commercial

Discovering Thoughts, Inventing Future

VOLUME 15 ISSUE 2 VERSION 1.0

© 2001-2015 by Global Journal of Medical Research , USA

Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine

Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine

Volume 15 Issue 2 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Medical Research . 2015.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric SciencesPurdue University Director
National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Issue

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue

1. A Cross-Sectional Survey of Bovine Fasciolosis at Elkadaro Abattoir, Khartoum State, Sudan. *1-7*
2. Effects of Feeding Processed Kidney Bean Meal (*Phaseolus Vulgaris*) by Replacing Soybean Meal on Egg Fertility and Qualities of Chicks of White Leghorn Hens. *9-16*
3. Prevalence of Bovine Mastitis in Lactating Cows and its Public Health Implications in Selected Commercial Dairy Farms of Addis Ababa. *17-24*

v. Fellows and Auxiliary Memberships
vi. Process of Submission of Research Paper
vii. Preferred Author Guidelines
viii. Index

Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine
Volume 15 Issue 2 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

A Cross-Sectional Survey of Bovine Fasciolosis at Elkadaro Abattoir, Khartoum State, Sudan

By Mohammed B. Badreldeen & Abdelhamid A. Elfadil
Sudan University of Science and Technology, Sudan

Abstract- A cross-sectional study was conducted from May to July 2011. The multivariate analysis exposed significant associations by postmortem examination: age >4 years (P= 0.000, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed (P= 0.000, OR= 77.6, 95% CI= 9.3, 81.6), light weight (P= 0.000, OR= 3.0, 95% CI= 1.9, 5.3), Ethiopian cattle (P= 0.000, OR= 76.1. 95% CI= 8.4, 83.7) and small size (P= 0.000, OR= 3.1, 95%CI= 1.9, 5.3). while, the coprology demonstrated significant associations among: age >4 years (P= 0.000, OR= 28.8, 95% CI= 3.9, 34.2), foreign breed (P= 0.000, OR= 94.4, 95% CI= 13.2, 102.6), light weight (P= 0.000, OR= 53.4, 95% CI= 7.1, 61.3), Ethiopian cattle (P= 0.000, OR= 60.2, 95% CI= 8.3, 70.1) and small size (P= 0.000, OR= 54.9, 95% CI= 6.3, 63.8). A higher prevalence was recorded by postmortem examination (X^2= 1.669, P= 0.000). This study determined the prevalence of bovine fasciolosis.

Keywords: fasciolosis, cattle, prevalence, risk factors, abattoir, khartoum, sudan.

GJMR-G Classification : NLMC Code: QW 70

ACROSSSECTIONALSURVEYOFBOVINEFASCIOLOSISATELKADAROABATTOIRKHARTOUMSTATESUDAN

Strictly as per the compliance and regulations of:

© 2015. Mohammed B. Badreldeen & Abdelhamid A. Elfadil. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

A Cross-Sectional Survey of Bovine Fasciolosis at Elkadaro Abattoir, Khartoum State, Sudan

Mohammed B. Badreldeen [α] & Abdelhamid A. Elfadil [σ]

Abstract- A cross-sectional study was conducted from May to July 2011. The multivariate analysis exposed significant associations by postmortem examination: age >4 years (P= 0.000, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed (P= 0.000, OR= 77.6, 95% CI= 9.3, 81.6), light weight (P= 0.000, OR= 3.0, 95% CI= 1.9, 5.3), Ethiopian cattle (P= 0.000, OR= 76.1. 95% CI= 8.4, 83.7) and small size (P= 0.000, OR= 3.1, 95%CI= 1.9, 5.3). while, the coprology demonstrated significant associations among: age >4 years (P= 0.000, OR= 28.8, 95% CI= 3.9, 34.2), foreign breed (P= 0.000, OR= 94.4, 95% CI= 13.2, 102.6), light weight (P= 0.000, OR= 53.4, 95% CI= 7.1, 61.3), Ethiopian cattle (P= 0.000, OR= 60.2, 95% CI= 8.3, 70.1) and small size (P= 0.000, OR= 54.9, 95% CI= 6.3, 63.8). A higher prevalence was recorded by postmortem examination (X^2= 1.669, P= 0.000). This study determined the prevalence of bovine fasciolosis.

Keywords: fasciolosis, cattle, prevalence, risk factors, abattoir, khartoum, sudan.

I. Introduction

Fasciolosis is among the most neglected important tropical diseases. Although it has significant economic impact on livestock industry, particularly cattle and sheep and occasionally can infects human beings (CDC, 2013). Among the estimated 91.1 million humans at risk for infection worldwide, as many as 17 million may be infected (Tolan, 2011). However, the disease is also consider as one of the major parasitic diseases contributing to loss in productivity estimated at over 200 US$ million per annum worldwide (Sturat, 1998). The infection is due to the food- and water-borne route. The two species most commonly implicated, as the etiological agents of fasciolosis are *F. hepatica* and *F. gigantica* (CDC, 2013).

Sudan possesses one of the highest livestock populations in Africa but productivity is low as a result of diseases, malnutrition and other management problems. Fasciolosis is one of these diseases which is responsible for considerable economic losses in livestock production (Boray, 1985). The disease is proved to be endemic in certain districts that characterized by intensive sheep or cattle production, in addition to the existence of favorable habitats for the snails host, like: White Nile, Eljazeera and Sennar regions (Ali, 1983). Another negative economic impacts on indigenous livestock result from inefficient conversion of feed, retarded growth, death, condemnation of infected livers, cost of preventive and treatment programs, reduced production, predisposition to other diseases, restricted use of infested lands and protein deficiencies among livestock dependant people (Saad, 2004).

Diagnosis is made serologically most often, although fecal examination for the eggs is fruitful if obtained when the adult worm is laying eggs (Tolan et al., 2011).

Therefore, the present study was designed to estimate the prevalence of bovine fasciolosis among cattle slaughtered at Elkadaro Abattoir, to identify potential risk factors associated with the occurrence of fasciolosis and to evaluate the accuracy of fecal examination.

II. Materials and Methods

a) Study Abattoir

Elkadaro Abattoir was designed for a processing capacity of 30 cattle per hour, with shifting work system every 10 hours (two shifts per day), in five days weekly. The abattoir was classified as code no.1, certified via O.I.E categories. The plant is situated within an open free disease area (recognized as free zone referring to O.I.E scientific terms). The main task of the abattoir is to process fresh and frozen meat, mainly for export orders. The abattoir is managed directly by the Federal Ministry of Animal Resources and Fisheries.

b) Duration of Study

A field investigation was launched in Sunday 21/ May and completed in Saturday 23/ July 2011.

c) Study Animals

According to the latest estimation of the livestock population in Khartoum State 2010, about 33800 of cattle are raised. The target animals were provided from different sites in and out of Sudan. The local cattle were fetched from: Nyala, Kordofan, Kosti, Kassala, Eljazeera, Khartoum and Upper Nile (South Sudan). While, the foreign ones were from Ethiopia (eastern neighboring country).

d) Sample Size Determination and Sampling Methods

The sample size was calculated using the formula: $4PQ/L^2$, given by (Martin et al., 1988) where:

Author α σ: College of Veterinary Medicine, Sudan University of Science and Technology, P.O. Box: 204, Khartoum north 13314, Sudan.
e-mails: mohammed_bushra2000@yahoo.com, aamelfadil@yahoo.com

© 2015 Global Journals Inc. (US)

P≡ prevalence
Q≡ 1-P
L^2≡ allowable error, for systematic random sampling with 7.4% reported prevalence (Eldoush, 1995) and 3% allowable error. Accordingly, the sample size was determined to be 307.

The sampling procedure was carried out in such a way that from daily 120 slaughtered cattle, 10 were randomly selected. There are five slaughter days a week and accordingly, 50 cattle were examined weekly. Hence, 307 cattle were examined within two months of the study period. A fresh feces was collected instantly after slaughtering of the selected animals. Then livers were subjected to detailed postmortem examination.

e) Study Methodology

Coprology: Fecal samples for parasitological examination were collected directly from the rectum of each animal immediately after slaughter using disposable plastic gloves and placed in new plastic bags. Prior to slaughter, each selected animal was given an identification number. Then each fecal sample was clearly labeled with the cattle identification number. Samples were kept at the room temperature and examined fresh. In laboratory, coproscopic examinations were performed to detect *Fasciola* eggs using standard sedimentation technique as previously described (Coles, 1986).

Liver Inspection: Liver of each cattle was strictly examined for the presence of liver flukes separately to correlate the coprology and postmortem examination of each animal. Examination of livers for *Fasciola* was carried out immediately after removal of liver from abdominal cavity. The inspection was made according to the procedures certified by FAO (2003).

f) Sensitivity and Specificity of the Fecal Examination Method

One of the objectives of this study was to evaluate the accuracy of the direct coprological examination method, which is routinely employed at field to examine the presence of *Fasciola* species eggs in feces. The sensitivity and specificity of the method was computed by taking liver inspection at postmortem as gold standard for the diagnosis of fasciolosis. Kappa statistic was used to determine the degree of agreement between the two methods of liver fluke diagnosis. The kappa value was interpreted as: slight agreement (K <0.2); fair agreement (K= 0.2→0.4); moderate agreement (K= 0.4→0.6); substantial agreement (K= 0.6→0.8); and almost perfect agreement (K >0.8) (Thrusfield, 2005).

g) Data Management and Analysis

Both fecal examination and liver inspection results were recorded on specially designed forms and preliminary analysis was done in Microsoft Excel. The outcome variable was the cases of fasciolosis detected during routine postmortem inspection (positive or negative) and fecal examinations for *Fasciola spp* eggs (positive or negative). Descriptive statistics were carried out to summarize the prevalence and proportion of infection in each category of investigated potential risk factors. Univariate and multivariate logistic regression analysis were conducted to see the significance and strength of association between potential risk factors and the occurrence of the infection. 95% confidence interval and p-value (P=<0.05) were used to notice the significance of association. Also, Odds Ratios (Exp B) was employed to assess the strength and direction of this association using SPSS statistical software (SPSS 16.0).

III. Results

a) Abattoir (Postmortem) Prevalence

Of the total 307 slaughtered cattle that subjected to detailed postmortem examination at Elkadaro Abattoir, 31.6% (97/307) were found positive for fasciolosis (Table I). The highest prevalence was recorded in age greater than 4 years (44.5%), foreign breed (64.1%), light weight (47.4%), Ethiopian source (65.5%) and small size (47.8%) (Table III).

b) Risk Factor Analysis for the postmortem results

The occurrence of fasciolosis significantly varied with age, breed, weight, source and animal size (P=<0.05). The likelihood of fasciolosis occurrence was significantly higher in age >4 years (P=0.000, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed (P=0.000, OR= 77.6, 95% CI= 9.3, 81.6), light weight (P=0.000, OR= 3.0, 95% CI= 1.9, 5.3) Ethiopian source (P=0.000, OR=76.1, 95% CI=8.4, 83.7), Kassala source (P=0.027, OR= 1.6, 95% CI= 1.1, 3.1), small size (P=0.000, OR=3.1, 95% CI= 1.9, 5.3) (Table III).

c) Prevalence by Coprology

Of the total 307 collected fecal samples 20.2% (62/307) were positive for coprological examination by sedimentation technique (Table II). The highest prevalence was recorded in age >4 years (28.9%), foreign breed (42.1%), light weight (43.8%), Ethiopian source (43.0%) and small size (44.1%) (Table IV).

d) Risk Factors Analysis for the Coprological Results

The results of coprological examination revealed significant association (P=<0.05) between the occurrence of fasciolosis and the risk factors: age, breed, source, weight and animal size. The likelihood of fasciolosis occurrence was significantly higher in age >4 years (P= 0.000, OR = 28.8, 95% CI = 3.9, 34.2), foreign breed (P= 0.000, OR = 94.4, 95% CI= 13.2, 102.6), Ethiopian source (P= 0.000, OR = 60.2, 95% CI = 8.3, 70.1) light weight (P= 0.000, OR = 53.4, 95% CI= 7.1, 61.3) and small size (P= 0.000, OR = 54.9, 95% CI= 6.3, 63.8) (Table IV).

© 2015 Global Journals Inc. (US)

e) Difference in Prevalence between the Two Diagnostic Methods

Based on the proportions comparison test there was significant difference (X^2=1.669, P=0.000) between fasciolosis prevalence estimated by coprology and postmortem examinations. Hence, in this study, higher prevalence of infection was observed by postmortem examination (31.6%) than by coprology (20.2%) (Table V).

f) The Sensitivity and Specificity of the Fecal Examination Technique Considering the Presence of Fasciola spp in the liver as a Gold Standard Test

As indicated in (Table VI), no animal that was positive with fecal examination and negative during postmortem examination. This revealed that postmortem examination was the golden test for diagnosis of fasciolosis when compared with coprology. The table set out the number of positive and negative tests in animals with and without flukes in their livers (Smith, 1995). The sensitivity and the specificity of fecal examination were found to be 63.9% and 100%, respectively. The calculated Kappa value (Kappa= 0.69) indicated substantial agreement between the two techniques.

IV. Discussion

Fasciolosis is a wide spread ruminant health problem and causes significant economic losses to the livestock industry in some areas in Sudan. The abattoir prevalence of fasciolosis obtained from the present study (31.6%) is very high compared to 7.4% (Eldoush, 1995) and nearly similar with 30% (Elmannan, 2001) and slightly lower than 34.4% (Abu-rigaila, 1983). These differences within the country could be attributed mainly to variations such as altitude, rainfall and temperature, although differences in livestock management system and the ability of the meat inspectors to detect the infection may playapart (Abu-rigaila, 1983). From African countries, a higher prevalence of 63.8% from Tanzania (Keyyu et al., 2006) and 53.9% from Zambia (Phiri et al., 2006) were reported. The observed prevalence may reflect suitable ecological and climatic conditions for the snail intermediate host in the areas from which the study animals came from.

Regarding the risk factors analysis, the results of this study indicate that the occurrence of fasciolosis in cattle varied with sex, age groups, breeds, weights, sources, size of animals and other diseases concurrent with fasciolosis.

The association between the occurrence of bovine fasciolosis and sex of the animals by both coprological and postmortem examinations revealed that the prevalence of *Fasciola* infection was found to be higher in males in agreement with a previous report (Shiferaw et al., 2009). The significant effects of sex on the prevalence of bovine fasciolosis might be attributed to the management system in which males are kept outdoor while females are kept indoor at the beginning of lactation (Balock et al., 1985).

Higher *Fasciola* infection rate was recorded among older cattle (>4 years) with both postmortem and coprological examinations in agreement with a previous report (Andrade et al., 2002). The higher prevalence in older animals by both examinations could be associated with the degree of exposure to the parasite which is normally greater in old animals than young animals. In contrast to our finding, higher prevalence in young cattle than older ones has been reported (Mulugeta et al., 2011).

The current study determined a higher prevalence of *Fasciola* infection among Ethiopian zebu cattle by both postmortem and coprological examinations in agreement with previous reported (Kassaye et al., 2012) and (Chakiso et al., 2014) This might be attributed to the difference in resistance to parasitic infection between different breeds (Tasawar et al., 2007).

In this study a higher prevalence of fasciolosis was observed among Ethiopian source animals by postmortem and coprological examinations in consistence with another finding (Yilma et al., 2000). The higher prevalence of the disease among Ethiopian source animals could be associated with the existence of suitable ecological conditions for the intermediate snail host in the areas where animals graze (Abebe et al., 2010).

We determined a higher prevalence of *Fasciola* infection rate among light weight animals with both postmortem and coprological examinations in agreement with previous reports (Kassaye et al., 2012) and (Nega et al., 2012). This signifies that the light weight animals are more susceptible to the infection.

The results of our study also showed a higher infection rate among small size animals by postmortem and coprological examinations in agreement with a previous report (Bekele et al., 2012). This attributed to low resistance against the disease.

In this study the association between the occurrence of fasciolosis with concurrent infections with both coprological and postmortem examinations revealed that there is no significant difference between the concurrent infections with fasciolosis by both examinations.

The prevalence of fasciolosis reported by using coproscopy was lower than that obtained by the abattoir results indicating that the latter is more sensitive in detecting the disease. The detection of *Fasciola* eggs can be unreliable as the eggs are expelled intermittently, depending on the evacuation of the gall bladder (Briskey et al., 1998). Similar study suggested that about 36% infected animals may pass undetected with single fecal examination technique. This might be attributed partly to the fact that *Fasciola* eggs only appear in feces 8-15

© 2015 Global Journals Inc. (US)

weeks post infection, so most of pathological lesions had already occurred (Mulugeta et al., 2011).

The present sensitivity value (63.9%) is comparable to other reports: 65.9% from Ethiopia (Regassa et al., 2012) and 69% from Switzerland (Rapsch et al., 2006). The latter stated that traditional coproscopy can be very efficient if there is repeated sampling, resulting in sensitivity of approximately 92%. Therefore, worm counts at liver necropsy can only be considered as a gold standard if the livers are sliced and soaked. Even then very light or prepatent infections could still be missed, affecting the calculated sensitivity and specificity of the evaluated tests.

The current study revealed that the infection with fasciolosis varies according to different regions in Sudan and demonstrated that the disease had a high prevalence among Ethiopian cattle. It also elucidated that coprological examination for the parasite eggs has significant limitations in detecting the presence or absence of fasciolosis in animals. On the other hand, it has helped to illustrate the usefulness of meat inspection in monitoring disease situation and demonstrating possible long term trends. This study also showed that bovine fasciolosis is significantly associated with age, breed, weight, source and size of animal.

V. Acknowledgements

Special appreciation to Elkadaro Abattoir officials and all the workers for their willingness and collaboration which was indispensable inputs for the accomplishment of this study. All contributions and supports are gratefully acknowledged.

References Références Referencias

1. Abebe, R., F. Abunna., M. Berhane ., S. Mekuria., B. Megersa., and A. Regassa. 2010. Fasciolosis: Prevalence, financial losses due to liver condemnation and evaluation of a simple sedimentation diagnostic technique in cattle slaughtered at Hawassa Municipal abattoir, Southern Ethiopia. Ethiopian Veterinary Journal **14**:39-5.
2. Abu-rigaila, H. A. 1983. Studies on naturally occurring and experimental ovine Fasciolosis in Sudan. M.V.Sc Thesis. University of Khartoum, 7 p.
3. Ali, M. A. 1983. Epidemiological studies on bovine fasciolosis in the White Nile Province. M.V.Sc Thesis. University of Khartoum, 5 p.
4. Andrade, S. R., A. Paz-Silva., L. J. Suarez., R. Panadero., J. Pedreira., C. Lopez., P. Diez-Banos., and P. Morrondo. 2002. Influence of age and breed on natural bovine fasciolosis in an endemic area (Galicia, NW Spain). Veterinary Research Communications **26**:361-370.
5. Balock, F. C., and R. J. Arthur. 1985. A survey of fasciolosis beef cattle killed at abattoir in southern Queensland. Australian Veterinary Journal **62**:324-326.
6. Bekele, M., H. Tesfay., and Y. Getachew. 2012. Bovine Fasciolosis: Prevalence and its economic loss due to liver condemnation at Adwa Municipal Abattoir, North Ethiopia. Ethiopian Journal of Applied Science and Technology **3**:319-324.
7. Boray, J. C. 1985. Flukes of Domestic Animals. In: Gaafar, S.M., Howard, W.S. and R.E. Marsh, (Eds.) Parasites, Pests and Predators. Elsevier 178-218.
8. Briskey, D. W. 1998. Diagnosis of liver fluke infections in cattle, Veterinary Bulletin 68-72.
9. Centre of Disease Control and Prevention. Va. USA. 2013. Available @www.dpd, cdc, gov/ dpdx/htm/ fasciolosis, htm.
10. Chakiso, B., S. Menkir., and M. Desta. 2014. On farm study of bovine fasciolosis in lemo districts and its economic loss due to liver condemnation at Hossana Municipal Abattoir, Southern Ethiopia. International Journal of Current Microbiology and Applied Sciences **3**:1122-1132.
11. Coles, E. 1986. Veterinary Clinical Pathology. Professor of Clinical pathology and Immunology. 4th ed. Kansas State University, 374-376 p.
12. Eldoush, M. E. 1995. Some studies on liver and gastro-intestinal parasites of cattle at Atbara town, Northern State, Sudan. M.V. Sc Thesis. University of Khartoum, 6 p.
13. Elmannan, A. M. 2001. Studies on biological control of lymnea natalensis snails using a molluscicidal indigenous plant pulicaria crispa (forsk) oliv compositae. Ph. D Thesis. University of Khartoum, 9 p.
14. FAO. 2003. Diagnostic Manual on Meat Inspection for developing countries.
15. Kassaye, A., N. Yehualashet., D. Yifat., and S. Desie. 2012. Fasciolosis in slaughtered cattle in Addis Ababa abattoir, Ethiopia. Global Veterinaria **8**:115-118.
16. Keyyu, J. D., A. A. Kassuku., L. P. Msalilwa., J. Monrad., and N. C. Kyvsgaard. 2006. Cross-sectional prevalence of helminth infections in cattle on traditional, small-scale and large-scale dairy farms in Iringa District, Tanzania. Veterinary Research Communications **30**:45-55.
17. Martin, W. S., L. A. Meek., and P. Willeberg. 1988. Veterinary Epidemiology. 1st ed. Ontario: Ontario Veterinary College, 33 p.
18. Mulugeta, S., F. Begna., and E. Tesgaye. 2011. Prevalence of Bovine Fasciolosis and its Economic Significance in and Around Assela, Ethiopia. Global Journal of Medical Research **11** (3):1-7.
19. Nega, M., B. Bogale., M. Chanie., A. Melaku., and T. Fentahun. 2012. Comparison of Coprological and Postmortem Examinations Techniques for the

© 2015 Global Journals Inc. (US)

Determination of Prevalence and Economic Significance of Bovine Fasciolosis. Journal Advanced Veterinary Research **2**:18-23.

20. Phiri, A. M., I. K. Phiri., and J. Monrad. 2006. Prevalence of amphistomiasis and its association with *Fasciola gigantica* infections in Zambian cattle from communal grazing areas. Journal of Helminthology **80**:65-73.
21. Rapsch, C., G. Schweizer., F. Grimm., L. Kohler., C. Bauer., P. Deplazes., U. Braun., and P. R. Torgerson. 2006. Estimating the true prevalence of *Fasciola hepatica* in Cattle slaughtered in Switzerland in the absence of an absolute diagnostic test. International Journal of Parasitology **36**:1153-1161.
22. Regassa, A., T. Woldemariam., S. Demisie., N. Moje., D. Ayana., and F. Abunna. 2012. Bovine Fasciolosis: Coprological, Abattoir Survey and Financial loss Due to Liver Condemnation in Bishooftu Municipal Abattoir, Central Ethiopia. European Journal of Biological Sciences **4**:83-90.
23. Saad, A. A. 2004. A survey of diseases prevalent in dairy cows in the White Nile and Gezira State, Sudan. M.V.Sc Thesis. University of Khartoum, 4 p.
24. Shiferaw, S., A. Kumar., and K. Amssalu. 2009. Organs condemnation and economic loss at Mekelle Municipal Abattoir, Ethiopia. Haryana Veterinarian **48**:17-22.
25. Smith, R. 1995. Veterinary clinical pathology. A problem-oriented Approach, 2nd ed. Florida: CRC Press-Boca Raton, 52 p.
26. Sturat, J. A. 1998. The life cycle of F. hepatica. In Dalton J.P. (Ed), Fasciolosis. CBA International Publication Cambridge, p. 1.
27. Tasawar, Z., U. Minir., C. Hayat., and M. H. Lashari. 2007. The prevalence of *Fasciola hepatica* in Goats around Multan. Pakistan Veterinary Journal **27**:5-7.
28. Thrusfield, M. 2005. Veterinary Epidemiology. 3rd ed. UK: Black Well Science, 183p.
29. Tolan, R.W. Jr. 2011. Fasciolosis Due to *Fasciola hepatica* and *Fasciola gigantica* Infection: An Update on this 'Neglected' Neglected Tropical Disease. Laboratory Medicine **42** (2):107-116.
30. Yilma, J., A. Mesfin. 2000. Dry season bovine fasciolosis in North Western part of Ethiopia. Revue de Médecine Vétérinaire **6**:493-500.

Table 1 : Prevalence of fasciolosis in 307 cattle examined by postmortem examination at Elkadaro abattoir, Khartoum, Sudan

Disease	Frequency	Relative frequency (%)	Cumulative frequency (%)
Negative	210	68.4% (210/307)	68.4%
Positive	97	31.6% (97/307)	100%
Total	307		

Table 2 : Prevalence of fasciolosis in 307 cattle examined by coproscopic examination at Elkadaro abattoir, Khartoum, Sudan

Disease	Frequency	Relative frequency (%)	Cumulative frequency (%)
Negative	245	79.8% (245/307)	79.8%
Positive	62	20.2% (62/307)	100%
Total	307		

© 2015 Global Journals Inc. (US)

Table 3 : Multivariate analysis for the association between fasciolosis (examined by postmortem) and potential risk factors using logistic regression

Risk Factors	No. tested	No. positive (%)	Exp (B)	95% C.I. for Exp (B)	p-value
Age (years)					
<2	25	0	- - -	- - -	1.000
2-4	71	3 (4.2%)	1	- - -	.999
>4	211	94 (44.5%)	18.5	3.1, 21.7	0.000
Breed					
Local	128	3 (2.3%)	1	- - -	1.000
Cross	34	1 (2.9%)	1.3	0.4, 3.2	0.641
Foreign	145	93 (64.1%)	77.6	9.3, 81.6	0.000
Weight					
Light	137	65 (47.4%)	3.0	1.9, 5.3	0.000
Medium	139	32 (23.0%)	1	- - -	1.000
Heavy	31	0	---	- - -	1.000
Source					
Nyala	79	2 (2.5%)	1	- - -	1.000
Ethiopia	142	93 (65.5%)	76.1	8.4, 83.7	0.000
Khartoum	35	0	- - -	- - -	1.000
Kordofan	13	0	- - -	- - -	0.999
Eljazeera	1	1 (100%)	- - -	- - -	1.000
White Nile	9	0	- - -	- - -	0.999
Kassala	25	1 (4 %)	1.6	1.1, 3.1	0.027
South Sudan	3	0	- - -	- - -	1.000
Animal Size					
Small	136	65 (47.8%)	3.1	1.9, 5.3	0.000
Medium	141	32 (22.7%)	1	- - -	1.000
Large	30	0	- - -	- - -	1.000

Table 4 : Multivariate analysis for the association between fasciolosis (examined by coprology) and potential risk factors using the logistic regression

Risk Factors	No. tested	No. positive (%)	Exp (B)	95% C.I. for Exp (B)	p-value
Age (years)					
<2	25	0	- - -	- - -	1.000
2-4	71	1 (1.4 %)	1	- - -	1.000
>4	209	61 (28.9%)	28.8	3.9,34.2	0.000
Breed					
Local	128	1 (0.7%)	1	- - -	1.000
Cross	34	0	- - -	- - -	1.000
Foreign	145	61 (42.1%)	94.4	13.2,102.6	0.000
Weight					
Light	137	60 (43.8%)	53.4	7.1,61.3	0.000
Medium	139	2 (1.4%)	1	- - -	1.000
Heavy	31	0	- - -	- - -	1.000
Source					
Nyala	79	1 (1.2%)	1	- - -	1.000
Ethiopia	142	61 (43.0%)	60.2	8.3,70.1	0.000
Khartoum	35	0	- - -	- - -	1.000
Kordofan	13	0	- - -	- - -	1.000
Eljazeera	1	0	- - -	- - -	1.000
White Nile	9	0	- - -	- - -	1.000
Kassala	25	0	- - -	- - -	1.000
South Sudan	3	0	- - -	- - -	1.000
Animal Size					
Small	136	60 (44.1%)	54.9	6.3,63.8	0.000
Medium	141	2 (1.4%)	1	- - -	1.000
Large	30	0		- - -	1.000

© 2015 Global Journals Inc. (US)

Table 5 : The difference in fasciolosis prevalence estimated based on coprology and postmortem examination at Elkadaro abattoir during May -July 2011

Test Type	No. Positive	Prevalence	Diff. (95% CI)	X^2	p-value
Coprology	62	20.2%	7.3 (3.1, 9.4)	1.669	0.000
Postmortem	97	31.6%			

Diff = difference

Table 6 : The presence or absence of *Fasciola spp* eggs in the feces of cattle with and without *Fasciola* in the liver for measuring the sensitivity, specificity and agreement level of the two tests

Faecal examination	Presence of *Fasciola spp* in liver		Total
	Fluke (+ve)	Fluke (-ve)	
Egg present (+ve)	62	0	62
Egg absent (-ve)	35	210	245
Total	97	210	307

© 2015 Global Journals Inc. (US)

This page is intentionally left blank

© 2015 Global Journals Inc. (US)

Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine
Volume 15 Issue 2 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Effects of Feeding Processed Kidney Bean Meal *(Phaseolus Vulgaris)* by Replacing Soybean Meal on Egg Fertility and Qualities of Chicks of White Leghorn Hens

By Taju Hussein, Mengistu Urge, Getachew Animut & Sisay Fikru
Jigjiga University, Ethiopia

Abstract- The study was to evaluate feeding processed kidney bean meal (PKBM) instead of soybean meal (SBM) on fertility, hatchability, embryonic mortality and chick quality of white leghorn (WL) hens. Replacement of SBM with PKBM in the diet did not affect the fertility, hatchability and embryonic mortality. Chick length (15.63, 15.00, 15.33, 15.03 and 14.33 (SEM $=\pm 0.02$)) and chick weight (34.13, 34.20, 33.13, 33.06 and 32.47(SEM$=\pm 1.39$)) for T1, T2, T3, T4 and T5 respectively, were significantly ($P<0.05$) lower for T5 than the rest treatments. Visual assessment of chick quality is lower for treatments containing higher proportion of PKBM than treatments containing lower proportion of PKBM. Therefore, as it affects the quality of chicks at 100% replacement, it is only up to 75% replacement of SBM by PKBM (dosed at 195 g/kg concentrate diet) is possible without having significant negative effect on chick quality.

Keywords: embryonic mortality, fertility, hatchability, kidney bean, soybean meal.

GJMR-G Classification : NLMC Code: QW 70, WA 390

Strictly as per the compliance and regulations of:

© 2015. Taju Hussein, Mengistu Urge, Getachew Animut & Sisay Fikru. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Effects of Feeding Processed Kidney Bean Meal (*Phaseolus Vulgaris*) by Replacing Soybean Meal on Egg Fertility and Qualities of Chicks of White Leghorn Hens

Taju Hussein [α], Mengistu Urge [σ], Getachew Animut [ρ] & Sisay Fikru [ω]

Abstract- The study was to evaluate feeding processed kidney bean meal (PKBM) instead of soybean meal (SBM) on fertility, hatchability, embryonic mortality and chick quality of white leghorn (WL) hens. Replacement of SBM with PKBM in the diet did not affect the fertility, hatchability and embryonic mortality. Chick length (15.63, 15.00, 15.33, 15.03 and 14.33 (SEM =±0.02)) and chick weight (34.13, 34.20, 33.13, 33.06 and 32.47(SEM=±1.39)) for T1, T2, T3, T4 and T5 respectively, were significantly (P<0.05) lower for T5 than the rest treatments. Visual assessment of chick quality is lower for treatments containing higher proportion of PKBM than treatments containing lower proportion of PKBM. Therefore, as it affects the quality of chicks at 100% replacement, it is only up to 75% replacement of SBM by PKBM (dosed at 195 g/kg concentrate diet) is possible without having significant negative effect on chick quality.

Keywords: embryonic mortality, fertility, hatchability, kidney bean, soybean meal.

I. Introduction

Chick quality is affected by pre-incubation storage conditions, time in the Hatcher, and size of egg. Tona et al., (2004) Found that increased incubation storage produced poor quality chick. Larger eggs tend to have significantly poor quality chicks as compared to other egg size Tona *et al.,* (2004).

The fertility of an egg is affected by several factors originating from the hen such as her ability to mate successfully, to store sperm, to ovulate an egg cell, and to produce a suitable environment for the formation and development of the embryo. The fertility also depends on the cock's ability to mate successfully, quantity and quality of semen deposited Brillard (2007). When fertility is low, it can affect other categories because of the lack of uniformity of embryo temperature inside the egg set (not as much heat provided by the developing embryos).

Author α: Depertment of Animal and Range Science, Collage of Agriculture, Wolaita Soddo University, Ethiopia; Po. Box 138, Wolaita Soddo, Ethiopia. e-mail: tajuh47@gmail.com

Author σ ρ: Departments of Animal and Range Science, College of Agriculture and Environmental science, Haromaya University, Ethiopia, po. Box, 138 Dire Dawa, Ethiopia.

Author ω: Depertment of Animal and Range Science, Collage Dry Land Agriculture, Jig-jiga University, Ethiopia; Po. Box 1020 Jig-jiga, Ethiopia. e-mail: sisayfikru69@gmail.com

Hatchability is a process that has several critical points that can be monitored and controlled to produce consistently healthy and mature hatchlings. These includes assessing hatching egg, fertility, egg storage and care, evaluation of hatch residue, poultry processing, sanitation, and poultry health and viability Hullet (2007).

The cost of poultry feed is very high and it accounts for 60-70% of the layer production cost Wilson and Beyer (2000). In recent years, the price of conventional or basic feed ingredients has tremendously increased. This has made poultry and live-stock production very expensive. In Ethiopia where soybean and its meal are in short supply and very expensive, the use of soybean meal as protein source of poultry ration is limited. Thus, an alternative protein source should be assessed and used. Sisay et al., (2015b) suggested that 100% replacement of SBM with PKBM is possible at 10% (100 g/kg) of soybean in layer ration. Therefore, the present research was initiated to evaluate the effect of feeding processed kidney bean meal (*Phaseolus vulgaris*) by replacing soybean meal at 19.5% (195g/kg) on egg fertility and qualities of chicks of white leghorn hens.

II. Materials and Methods

a) *Management of Experimental Animals*

Experimental house which were partitioned into 15 pens with wire-mesh and covered with grass litter material of 10 cm depth were used for experiments. Before the commencement of the actual experiment, the experimental pens, (2.5*2m), watering equipment, feeding troughs and laying nests were disinfected, sprayed against external parasites and thoroughly cleaned. Disinfectant was placed at the gate of the experimental house for workers and other visitors to step on before they enter into the house, which is in addition to the one placed at the main gate of poultry farm, for prevention of disease introduction.

The birds were vaccinated against Newcastle disease according to the vaccination program of the farm. The birds were offered with experimental diets for 7 days as period of adaptation before actual data

© 2015 Global Journals Inc. (US)

collection takes places. The feed was measured and given to the birds in groups twice a day at 8 am and 2 pm on *ad libitum* basis by dividing the daily offer into two equal portions. Feed refusals were collected every morning at 7:30 am before providing new feed, external contaminants were removed by visual inspection, weighed and recorded for each pen separately. Feed were offered into two metal tubular feeders per pen that was hanged approximately at a height of the backs of the birds. Water was provided in a plastic fountains placed on a flat stone at the center of the pen. The watering trough was cleaned every morning before feeding. Clean and fresh water was available to the birds' *ad libtum*. The experimental duration lasted for 12 weeks.

b) Treatments and Experimental Design

The ration of the experiment consists of PKB, which replaced SBM at five levels, namely 0, 25, 50, 75 and 100% PKB for T1, T2, T3, T4 and T5, respectively (Table 1). Initially weighed two hundred ten (210) birds consisting of 180 White leghorn layers and 30 cocks of mature white leghorn breeds of similar age and weight, which were obtained from the University's poultry farm, were randomly assigned to the five treatment rations. Thus, the design employed was a completely randomized design (CRD) with five treatments and three replications (pen) per treatment (Table 1).

c) Chemical Composition of Experimental Feeds

Representative samples of feed ingredients was taken and analyzed before formulating the actual dietary treatment rations. The results of the analysis were used to formulate the treatment rations (Table 2). The feed ingredient offer samples were analyzed for dray matter (DM), ether extracts (EE), crude fiber (CF) and ash following the procedure of Weende (proximate analysis method) of the AOAC (1990). Kjeildhal procedure was employed to determine the nitrogen (N) content of the feeds and crude protein (CP) was determined by multiplying the N value with 6.25. The total metabolizable energy (ME) contents were calculated indirectly by using the formula presented by Wiseman (1987). ME (Kcal/kg DM) = 3951 + 54.4 EE – 88.7 CF – 40.8 Ash.

d) Data Collection and Measurements

i. *Fertility and hatchability*

Hundred ninety-five, that is 39 and 13 eggs from each treatment and replication, respectively, of medium sized, non-defected and normal shaped eggs were collected in three consecutive days. The eggs were kept at room temperature until incubated within 4 days after the first day of collection. The eggs were lightly coded by marker before they were placed into the incubator. The incubation temperature and relative humidity during the 18 days of incubation was auto fixed at 37-38°C and 65-70%, while that of hatchery unit was adjusted to 38.5-39°C and 90% relative humidity.

The eggs kept in the tray with small end down and turned automatically by slanting the tray at 45°. The incubator is equipped with the turner that facilitates the turning operation at an interval of two hours. Fertility was determined by candling the incubated eggs on the 7th day of incubation. Candling was done at dark room with egg Candler powered by electricity. Eggs found to be infertile, which are characterized by clear appearance, egg with blood adhering to one sides of the eggs were drown from the incubator. Finally eggs found fertile, i.e. eggs having small dark spot, numerous blood vessels arising from those dark spot of yolk at day of candling, clearly visible thick and dark and well fill structure was further kept in the incubator for hatching North (1984). Eggs with living embryo were transferred to the hatching section of the incubator at the end of the 18thday. Hatched chicks were counted and chick quality was determined. Fertility and hatchability were determined by using the following formulae, respectively.

$$\%\text{Fertility} = \frac{\text{number of fertile eggs}}{\text{total egg set}} x100$$

$$\%\ \text{Hatchability on fertile eggs base} = \frac{\text{number of chick hatched}}{\text{fertile eggs}} x100$$

$$\%\ \text{Hatchability on total eggs base} = \frac{\text{number of chick hathced}}{\text{total eggs set}} x100$$

ii. *Embryonic mortality*

Embryonic mortality was determined by candling eggs at 7th, 14th and 18th days of incubation and at hatching. Eggs that had a structure encircled with blood ring, absence of blood vessels, adhering to the shell membrane and absence of clear demarcation between embryo and air cell was considered as dead embryo and removed from the incubator North (1984). The dead embryo further categorized in to early, mid, late dead and pip embryo, based on the classification criteria set by Butcher (2009). The embryonic mortality was computed by using the formulae indicated by Rashed (2004).

$$\%\ \text{of early mortality} = \frac{total\ numbr\ of\ early\ dead\ embryo}{total\ number\ of\ fertile\ eggs} x100$$

$$\%\ \text{of mid mortality} = \frac{total\ number\ of\ mid\ dead\ embryo}{total\ number\ of\ fertile\ eggs} x100$$

$$\%\ \text{of late mortality} = \frac{total\ number\ of\ late\ dead\ embryo}{total\ number\ of\ fertile\ eggs} x100$$

$$\%\ \text{of pip mortality} = \frac{total\ number\ of\ pip\ dead\ embryo}{total\ number\ of\ fertile\ eggs} x100$$

© 2015 Global Journals Inc. (US)

iii. *Chick quality*

Chicks' quality were measured using two different methods, which includes visual scoring and measuring of day old chick weight and length. Visual scoring of chicks was done by visual examination based on the quality standards described by North (1984). Accordingly, a chick that was not malformed, physically active, stands up well, and looks live has been taken as good quality chicks. The researcher and two technicians were under taken the visual quality assessment. Based on common decisions, the chicks were classified in to poor and normal chicken. After the chicks were classified in to two groups, namely quality (chicks with dried body, stand well, and active) and non-quality chicks (chicks with wet body, not firm, non-straight, and not having perpendicular leg to the ground), five chicken from normal chicken were taken randomly from each replicate and their weight and length were measured using sensitive balance and ruler for further quality assessments respectively. The chick's length was taken by stretching the chicks on the table and taking the length from the tip of the beak to the middle toe by a ruler. Percentage of quality chicks were calculated as

Quality chicks =

$$\frac{\text{total number of quality chicks}}{\text{total number of hatched chicks}} x100$$

e) *Statistical Analysis and Models*

The data collected during the study period were subjected to statistical analysis using SAS computer software version 9.1.3. SAS (2008). during data analysis, chick weight and length were analyzed following one way analysis of variance procedure. When the analysis of variance indicated the existence of significant difference between treatment means, list significant difference (LSD) method was used to locate the treatment means that were significantly different from the other Gomez and Gomez (1984).

The model used for statistical analysis was:

$Yij = \mu + Ti + eij$

Where: Yij=Individual observation,

Ti=Treatment effect, μ=Overall mean, eij= Error term

General logistic regression analysis was employed for analysis of data recorded on fertility (fertile/infertile), hatchability (hatched/un-hatched), embryonic mortality (alive/dead), and visual scoring (normal/poor). The general logistic regression model used is given below:

Model: $1n\left\{\frac{\pi}{1-\pi}\right\} \beta 0+\beta 1^{*}$ (x)

Test H0: No treatment effect (*i.e.*, β1=0) vs.

HA: Significant treatment effect (β1≠ 0).

Where, π=probability,

β=slope, x=treatment

III. Results and Discussion

a) *Nutrient Composition of Ingredients and the Treatment*

The results of the chemical analysis of ingredients used and nutritional composition of the ration–for each treatment are given in Table 3 and 4 respectively. The DM contents of PKB obtained in the present study was slightly lower than reported by Marzo et al., (2002) (93.2%) and Audu and Aremu (2011) (96.8) but very close to Sisay et al., (2014) and Sisay et al. (2015a) (88) while CP contents is slightly higher than reported by Marzo et al., (2002), Audu and Aremu (2011), Emiola (2011), Sisay et al., (2014), Sisay et al., (2015b) and Emiola and Olghobo (2006) which were 20.9%, 23.6%, 24.7%, 25.8% and 26.8%, respectively. The EE content of kidney bean is the same with that reported by Emiola (2011) but Marzo et al., (2002) reported lower than the current results. The CF content of kidney bean used in the present study was comparable to that reported by Arija et al., (2006), Emiola (2011), Sai-ut et al., (2009), Audu and Aremu (2011), Sisay et al., (2014) and Sisay et al., (2015b) who reported 5.1, 5.0, 6.0, 4.7% and 4.5% respectively. The result of chemical composition of kidney bean used in the present experiment showed comparable ME contents to that noted by Ofongo and Aloghobo (2007) and Arija et al., (2006) who reported 3342.2 and 3365 kcal/kg, respectively.

In general, as different report showed the processing (treatments) could cause drop or increase in nutrients compared to the raw seed. The treatment method (boiling) employed was suggested increased total carbohydrates and decrease the CP, EE, CF and ash contents Marzo et al., (2002), Audu and Aremu (2011). Similarly, Akaerue and Onwuka, (2010) and Mubarak (2005) reported reduced CP, EE, CF and ash for Mung bean (*Vigna radiata*). The low content of CF of PKB and the further reduction as a result of boiling might favored the feed intake of the layers by decreasing the problem of feed digestibility.

SBM and NSC are rich in CP content that make these ration to be ideal source of protein supplement for poultry. The NSC used to formulate experimental ration composed 26% CP and 2006 kcal ME. The CP values are comparable with that of Shewangzaw et al., (2011) and Meseret et al., (2011) but low in ME. Previously under taken studies indicated that the CP content of SBM to be in the range of 41 to 50% Waldroup (2002), Ekeren et al., (2006). However, similar to the value obtained in this study, a 38% CP content of SBM were reported in Ethiopia Meseret et al., (2011).

When SBM and PKB are compared, SBM contain by far higher CP, EE and ME. On the other hand, PKB has lower CF and ash than SBM. As a result, the CP of the formulated ration ranged between 16% (diet 5) and 18% (diet 1). Crude fiber increased, EE

© 2015 Global Journals Inc. (US)

decreased because of total replacement of SBM by PKB. The energy content of SBM is higher than that of PKB. For this reason, the energy content decreased with increasing level of processed kidney bean. Nevertheless, energy and protein content of all rations ranged within the recommended level for layers. Lesson and summer (2001) recommended 16-18% of CP and 2500-3300 kcal/kg ME, respectively for white leghorn layers. Furthermore, the energy of compound ration is the same with that used by Zebiba (2012) and Senayt et al., (2011) in their experiment for the same bird in the same farm.

b) Fertility and Hatchability of Eggs

Mean values of fertility and hatchability for the treatments are presented in Table 5. The logistic regression results for fertility and hatchability of eggs showed no significant difference among treatments. However, there was a numerical decrease in fertility and hatchability percentage for T5 as compared toT1 which is similar with Sisay et al., (2015b) finding. The numerical decrease in fertility and hatchability from T1 to T5 could be the result of increased, level of kidney bean in the ration that caused reduction of level of protein, calcium and phosphorus found in the rations. The diets of breeder poultry should be adequate in both quantity and quality to meet the recommended levels of feed standard Brillard (2007). The level of dietary protein significantly affected egg fertility and hatchability Gareil et al., (2006). Poor hatching happen when nutritionally deficient feed is used for layers Hocking et al., (2002). The authors have shown that low calcium levels tended to decrease percent of fertility and hatchability. On the contrary, El-Ghamry et al., (2010) reported improved fertility (83% and 77%) and hatchability (78% and 71%) for the group fed 2.4% and 2.6% calcium, respectively.

c) Embryonic Mortality

The mean values of embryonic mortality at different stages of development are presented in Table 5. The logistic regression result showed no significant differences among treatments for early, mid, late and pip embryonic mortality. Nevertheless, the mortality tends to numerically increase with increased level of PKB compared to control diets that do not contain PKB. This might occurred because of the decreased crude protein level with increased level of PKB. Embryonic mortality of eggs of breeder hens' fed low protein is reported to be higher than hens fed high protein diets. Low-protein rearing rations were associated with higher rates of food intake, higher mortalities and lower rates of egg production than the conventional protein ration Hocking et al., (2002).

d) Chick Quality

i. *Visual Scoring (Observation)*

The visual scoring of chicks is presented in Table 6. Wald chi-square statistics indicated that visual scoring of chicks was insignificant (0.64) at (α = 0.05) among the treatments, but Sisay et al., (2015b) indicated that visual scoring of chicken was not significant (*pr>chisq* 0.641 α=0.05). The visual assessment showed that the quality of chicken is better in the order of T2>T1>T3>T4, while the chick quality of T5 is inferior or poor on visual scoring. The chicken in T5 were not well standing, they are dehydrated and seems inactive at their day old age. This assessment was supported by length and weight of chicks that indicated significant differences among treatments. Utilization of visual score parameters such as naval quality, firmness of leg, size of beak, eye and vital chick are recommended ways of determining highest quality chicks Petek et al., (2010).

ii. *Chick Length*

The length of chicks hatched from eggs of hens fed diet containing 0, 25, 50 and 75% of PKB are not significantly different which is similar with Sisay et al., (2015b) finding that there was no significant difference ($p>0.05$) among treatments in chick weight and chick length. However, the chicken hatched for eggs harvested from layers fed 100% PKB (T5) had significantly lower length as compared to that from the layers fed SBM at different levels. This occurred either because of the egg size that accommodate larger embryo of chicken compared to the small eggs that has less environment to hold large chicken or the nutrition of the layers that promote better growth of the chicks. Mukhtar et al., (2013) Found similar result and stated that hatching length is an easy and repeatable quality evaluation parameter for newly hatched chicks. This important trait has a positive correlation with the size of the egg and the chick's weight. It is an important economic trait to predict chick development because it is positively related to yolk-free body mass at hatch and potential of chicks for optimum future performance.

Furthermore, Petek et al., (2010) pointed out that each extra cm of hatchling length at day of hatch meant an increase of 18 g BW at seven days of age. Chicks with longer hatchling length have better-feed efficiency and survival rates as compared to smaller chicks. Petek et al., (2009) Classified length intervals in to short, middle and long for a day old chicks. Accordingly, layer chick with a length of < 17.8, 17.8-18.2 and > 18.2cm, respectively are grouped as short, medium and long chicks. As to this classification, chick lengths in the present experiment for all treatments fall within a short category. On other hand, Petek et al., (2008) reported that the body weight and chick length uniformity in long group in all poultry to be better than the shorter group

iii. *Chick Weight*

The chick weight of layers were higher (P<0.05) in ration that do not contain SBM as compared to the ration containing 100%PKB (Table 6). The variation in

© 2015 Global Journals Inc. (US)

chick weight may be due to the weight of eggs, which is slightly decreasing across treatment as well as the amino acid content of the rations, which is higher in control diet while decreasing across treatment because of decreased SBM. The present finding is in agreement with many previous of findings. For instance the chicks injected with amino acids invariably had higher plasma protein and lower plasma glucose on the day of hatch, because methionine and threonine are critical for the growth of chicken embryo Subrat et al., (2012). Egg weight has a direct impact on the weight of chick and there is a positive correlation between egg and chick weight Petek et al., (2010). Chicken hatched from large eggs are heavier than those hatched from comparatively smaller eggs Al-Murrani (1978). A heavy chick indicates a good development. However, this is sometimes not true and evaluating chick quality by measuring only body weight can be misleading Molenaar (2009).

IV. Acknowledgement

The authors' heartfelt appreciation goes to the Ethiopian Ministry of Education (MOE) for fully sponsoring this study and Haramaya University for provision of research facilities.

References Références Referencias

1. Akaerue Blessing, I. and Onwuka Gregory, I. (2010) Effect of Processing on the Proximate Composition of the Dehulled and Undehulled Mungbean [Vigna radiata (L.) Wilczek] Flours. Pakistan Journal of Nutrition 9 (10): 1006-1016.
2. Al- Murrani, W.K. (1978) Maternal effects on Embryonic and Post-embryonic growth in poultry. British Poultry Science 19: 277-281.
3. AOAC (Association of Official Analytical Chemist) (1990) Official Method of Analysis (13th ed) 15th ed. AOAC Arlington, Verginia, USA. pp. 12-98.
4. Arija, Centeno, CA., Viveros, Brenes, A., Marzo, F. et al. (2006) Nutritional Evaluation of Raw and Extruded Kidney Bean (Phaseolus vulgaris L. var. Pinto) in Chicken Rations. Poultry Science 85: 635–644.
5. Audu, S.S. and Aremu, M.O. (2011) Effect of processing on chemical composition of red kidney bean (Phaseolus Vulgaris L.) Flour. Pakistan journal of Nutrition 10(11): 1069-1075.
6. Brillard, J.P. (2007) Control of fertility in turkeys: the impact of environment, nutrition and artificial reduction by maternal diets high in unsaturated fatty insemination technology. Poultry Industry Technical articles.
7. Butcher, G.D. and Nilipour, A.H. (2009) Chicken Embryo Mal positions and Deformities. Institute of Food and Agricultural Sciences, University of Florida. http://edis.ifas.ufl.edu
8. Eekeren, A.N. Maas, H.W. Saatkamp and Verschuur, M. (2006) Small Scale Chicken Production. Digigrafi, Wageningen press, Netherlands 99p.
9. El-Ghamry, El-Allawy, Hewida, Yassein, M.S.A. and El Mallah, G.M. (2010) Evaluation of Rationary Calcium Requirements in Fayoumi Laying Hens. Iranian Journal of Applied Animal Science 1(2): 81-86.
10. Emiola. I. A. (2011) Processed african yam bean (sphenostylis stenocarpa) inbroiler feeding: performance characteristicsand nutrient utilization. Journal of Environmental Issues and Agriculture in Developing Countries 3(3): 123-131.
11. Emiola, I. A. and Ologhobo, A. D. (2006) Nutritional assessment of raw and different processed Legume seed in Broiler ration. Journal of animal and veterinary advance 5(2):96-101
12. Gabreil, M. B. Babatunde, and Fetuga, L. (2006) Effects of different rationary oils superimposed on different rationary protein levels on the laying performance, egg weights, fertility and hatchability in the tropical environment. Journal of the Science of Food and Agriculture 27: 54- 62.
13. Gomez, K. A. and Gomez, A. A. (1984) Statistical Procedures for Agricultural Research. 2nd ed. John Willey and Sons, New York p720
14. Hocking, P. M. R. Bernard, and Robertson, G. W. (2002) Effects of low dietary protein and different allocations of food during rearing and restricted feeding after peak rate of lay on egg production, fertility and hatchability in female broiler breeders. British Poultry Scienc, 43:1, 94-103.
15. Hulet, R. M. (2007) Managing incubation, Where are we and why? Poultry Science 86: 1017-1019.
16. Leeson, S. and Summers, J. D. (2001) The Nutrition of Chicken. 2nd ed. University Books, Guelph, Canada 355p.
17. Marzo, F. Alonso, R. Urdaneta, E. Arricibita, F. J. and Ibáñez F (2002) Nutritional quality of extruded kidney bean (Phaseolus vulgaris L. var. Pinto) and its effects on growth and skeletal muscle nitrogen fractions in rats. Journal of Animal science 80:875-879.
18. Meseret Girma, Mengistu Urge, and Getachu Animtu, (2011) Ground Prosopis Juliflora poods as feed ingredient in poultry Ration: Effects growth and carcass characteristics of Broiler. International journals of poultry Sciences 10 (12):970-976.
19. Molenaar, R. (2009) Evaluations of chick quality; which method do you choose? HatchTech B.V., the Netherlands. www.hatchtech.nl
20. Mubarak, A. E. (2005) Nutritional composition and anti nutritional factors of mung bean seeds (Phaseolus aureus) as affected by some home traditional processes. Food Chemistry 89(4): 489-495.

© 2015 Global Journals Inc. (US)

21. Mukhtar, N. Khan, S. H. and Anjum, M. S. (2013) Hatchling length is a potential chick quality parameter in meat type chickens. World's Poultry Science Journal 69:889-895.
22. North, Mack, O. (1984) Commercial Chicken Production Manual. 3rd ed. AVI Publishing Company, Inc. West Port, Connecticut, Georgia 451p.
23. Ofongo, S. T. and Ologhobo, A. D. (2007) Processed Kidney Bean (Phaseolus vulgaris) in Broiler Feeding: Performance Characteristics. Conference on International Agricultural Research for Development pp-3.
24. Petek, M. Orman, A. Dikmen, S. and Alpay, F. (2010) Physical chick parameters and effects on growth performance in broiler. Archiv Tierzucht 53 (1):108-115.
25. Petek, M. Orman, A. Ddkmen, S. and Alpay, F. (2009) Relations between day –old chick length and body weight in broiler,quail and layer. Turkey Journal of Animal Sciences 27(2): 25-28.
26. Petek, M. Orman, A. Dikmen, S. and Alpay, F. (2008) Relations between dayold chick length and body weight in broiler, quail and layer. Journal of Veteriner Fakültesi Dergisi, Uludağ Üniversitesi 27 (1-2): 25-28
27. Rashed, M. (2004) Effects of feeding systems on the egg production of Fayoumi hens of model breeding unit under PLDP programin Bangladesh. MScThesis, http://www.smallstock.info/research/reports/Dan00pdf
28. Sai-Ut, S. Ketnawa, S. Chaiwut, P. and Rawdkuen, S. (2009) Biochemical and functional properties of proteins from red kidney, navy and adzuki beans. Asian Journal of Food and Agro-Industry 2(04), 493-504.
29. SAS (Statistical analysis system), (2004) SAS user guide. Version 9.0 institute Inc, NC.
30. Senayt Abraha, (2011) the effect of feeding different levels of soybean meal on egg production, quality, fertility and hatchability of white leghorn layers. M.sc Thesis, Haramaya University, Haramaya.
31. Shewangizaw Wolde, Tegene Negesse, and Aberra Melesse, (2011) Effect of dietary protein concentration on feed intake, body mass gain and carcass traits of Rhode Island Red chicken. Journal of Science and Development 1(1): 53-64.
32. Sisay Fikru, M, Mengistu Urge, L. and Getachew Animu, (2014) Effects of Replacing Soybean Meal with Processed Kidney Bean Meal (Phaseolus vulgaris) on Egg Production of White Leghorn Hens. World Applied Sciences Journal 32 (9): 1918-1926.
33. Sisay Fikru, M. Mengistu Urge, L. and Getachew Animu, (2015a) Effects of Replacing Soybean Meal with Processed Kidney Bean Meal (Phaseolus vulgaris) on qualities of Eggs of White Leghorn Hens. International Journal of Agricultural Science Research Vol. 4(3), pp. 049-056.
34. Sisay Fikru, M. Mengistu Urge, L. and Getachew Animu, (2015b) Effects of Replacing Soybean Meal with Processed Kidney Bean Meal (Phaseolus vulgaris) on egg fertility and chick quality of White Leghorn Hens. J. Adv. Vet. Anim. Res. 2(2): 146-152.
35. Subrat, K. Bhanja, Asit, Baran, Mandal, Sushil, K. Agarwal and Samir Majumdar, (2012) Modulation of Post hatch-growth and immune competence through in ovo injection of limiting amino acids in broiler chickens. Indian Journal of Animal Sciences. Abstract
36. Tona, K. Bruggeman, V. Onagbesan, O. Bamelis, F. Gbeassor, K. (2004) Effects of age of broiler breeder and egg storage on egg quality, hatchability, chick chick weight and quality chick post-hatch growth to forty-two days. Journal of Applied Poultry Research 13:10-18
37. Waldroup, W. P. (2002) Soybean meal in poultry nutrition. Poultry science, 0.210 faretteville AR 72701. Arakansas. www.soymeal.org
38. Wilson, K. J. Beyer, R. S. (2000) Poultry Nutrition Information for the Small Flock. Kansas State University Agricultural Experiment Station and Cooperative Extension Service. http://www.oznet.ksu.edu. (Accessed on January 31, 2015).
39. Wiseman, J. (1987) Feeding of Non-Ruminant Livestock. Butterworth and C.Ltd. 370 p.
40. Zebib Abisa (2012) Effects of substituting sorghum for maize on egg production, quality, fertility and hatchability of white leghorn layers. M.sc Thesis, Haramaya University, Haramaya.

Table 1: Experiment lay out

Treatments	Replications	Layers per replication	Cocks per replication
T1(100%SBM:0%PKB)	3	12	2
T2 (75% SBM:25%PKB)	3	12	2
T3 (50% SBM:50%PKB)	3	12	2
T4 (25% SBM:75%PKB)	3	12	2
T5 (0% SBM:100% PKB)	3	12	2

SBM=Soybean meal; PKB=Processed kidney bean; 100% PKB represents replacement of maximum SBM (260g/kg) as recommended by earlier study Senayt et al. (2011)[6]

© 2015 Global Journals Inc. (US)

Table 2 : Proportion of feed ingredient used in formulating experimental rations

Feed (%)	Treatments				
	T1	T2	T3	T4	T5
CG	56	53	52	48	37
WS	7	7	7	7	16
SBM	26	19.5	13	6.5	0
PKB	0	6.5	13	19.5	26
NSC	4	7	8	12	14
LS	5.5	5.5	5.5	5.5	5.5
Salt	0.5	0.5	0.5	0.5	0.5
VPM	1	1	1	1	1
Total	100	100	100	100	100

CG= corn grain; WS= wheat short; SBM= soybean meal; PKB= processed kidney bean; NSC= noug seed cake; LS= limestone; VPM= vitamin pre mix; T1 100%SBM: 0%PKB; T2= 75%SBM: 25%PKB; T3= 50%SBM: 50%PKB; T4=25%SBM: 75%PKB; T5= 0%SBM: 100%PKB

Table 3 : Ingredient used in the study and its nutrients compositions

Feed type	Nutrient composition (% for DM and % DM for others)					
	DM	CP	EE	CF	Ash	ME kcal/kg
CG	89.5	8.7	4.3	8.0	6.21	3230.5
WS	90.3	12	3.3	6.2	6.8	3303.1
SBM	90.2	38	7.0	9	7.8	3215
PKB	87.5	28	0.9	6	7.0	3182.2
NSC	91.5	26	6.0	21.0	10	2006.0

CG=Corn grain; WS=Wheat short; SBM=Soybean meal; PKB=processed Kidney bean; NSC=Noug seed cake; DM=dry matter; CP=Crude protein; EE=ether extract; CF=crude fiber; ME=methabolizable energy

Table 4 : Nutritional composition of treatment diets containing different levels of processed kidney bean as a replacement for soybean meal

Treatments	Nutrient composition (% for DM and % DM for others)							
	DM	CP	EE	CF	Ash	Ca	P	ME kcal/kg
T1	91.85	18	5.64	6.26	9.96	3.4	0.39	3296.20
T2	91.56	17.8	5.63	6.36	9.97	3.26	0.38	3286.40
T3	91.17	17.6	5.58	6.52	9.98	3.28	0.38	3269.00
T4	90.21	16.3	5.40	6.56	9.98	3.01	0.36	3255.70
T5	89.86	16.0	4.90	6.86	10.20	2.79	0.32	3192.90

DM= dry matter; CP= crude protein; EE = ether extract; CF= crude fiber; SBM= soybean meal; PKB processed kidney bean; T1 100%SBM: 0%PKB; T2= 75%SBM: 25%PKB; T3= 50%SBM: 50%PKB; T4=25%SBM: 75%PKB; T5= 0%SBM: 100%PKB

Table 5 : Fertility and hatchability of eggs and embryonic mortality in layers fed different levels of processed kidney bean as a replacementz to soybean meal

Parameter	Treatments					SEM	SL
	T1	T2	T3	T4	T5		
Fertility (%)	94.87	94.87	94.87	94.87	92.3	2.29	NS
HTES (%)	71.79	71.79	71.79	64.10	61.54	4.59	NS
HFES (%)	75.43	75.64	75.43	69.44	66.67	3.50	NS
EEM (%)	5.56	5.56	5.56	8.33	8.33	2.14	NS
MEM (%)	5.56	8.12	8.12	11.11	8.33	1.76	NS
LEM (%)	5.56	5.56	5.56	8.30	13.89	2.48	NS
PIEM (%)	10.89	8.12	10.89	11.11	11.11	3.07	NS

SBM= soybean meal; PKB= processed kidney bean;T1=100% SBM:0% PKB; T2=75%SBM:25%PKB; T3=50%SBM:50%PKB; T4=25%SBM:75%PKB; T5=0%SBM:100PKB; HTES= hatchability on total egg set; HFES= hatchability on fertile egg set; EEM =early embryonic mortal; MEM= mid embryonic mortality; LEM = late embryonic mortality; PIEM =pip embryonic mortality

© 2015 Global Journals Inc. (US)

Table 6 : Chick quality parameters of white leghorn layers fed different levels of processed kidney bean as a replacement to soybean meal

Parameters	Treatment					SEM	SL
	T1	T2	T3	T4	T5		
Visual (%)	84.93	85.09	82.36	81.02	70.83	3.89	NS
Length (cm)	15.63[a]	15.00[a]	15.33[a]	15.03[a]	14.33[b]	0.20	*
Weight (g)	34.13[a]	34.20[a]	33.13[a]	33.06[a]	32.47[b]	0.28	*

[a-b] means in the row without common superscript are significant; NS = Non- significant, SL= significance level; SEM= standard error of mean ; cm = cent meter, g = gram; % = percent; PKB = processed kidney bean meal; SBM= soybean meal; T1=100% SBM:0% PKB; T2=75%SBM: 25%PKB; T3=50%SBM: 50%PKB; T4=25%SBM: 75%PKB; T5=0%SBM: 100PKB;

© 2015 Global Journals Inc. (US)

Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine
Volume 15 Issue 2 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Prevalence of Bovine Mastitis in Lactating Cows and its Public Health Implications in Selected Commercial Dairy Farms of Addis Ababa

By Alebachew Tilahun & Alemu Aylate
Wolaita Sodo UniversityWolaita Sodo University, Ethiopia

Abstract- A cross sectional study was conducted in Addis Ababa from October 2011 to May 2012 to determine prevalence of bovine mastitis and discuss its public health implications. A total of 444 systematically selected lactating cows of different cattle breed from thirty seven (37) dairy farms were investigated. The herds were visited and the farmers interviewed about the management, housing, feed and feeding, and milking conditions. California Mastitis Test (CMT) was applied on milk samples collected from 1776 individual quarters. The overall prevalence of bovine mastitis was 68.0% (302/444) of which sub-clinical mastitis accounted for 46.8% (208/444) while 21.2% (94/444) were found to be clinical forms. There was significant difference ($P<0.05$) in the prevalence of mastitis among the different breeds, age groups, parity and lactation stage. Relatively higher number of farmers interviewed (20.8%) replied that they do not withhold milk from cows treated for mastitis and continue to avail it to the public without interruption. As it is economically damaging, the need to establish diagnostic facility to enable early detection for screening large number of samples was emphasized. Further work on identification of the causative agents and conducting public awareness creation about major zoonotic diseases were also recommended.

Keywords: *california mastitis test, interview, prevalence, mastitis, zoonotic.*

GJMR-G Classification : *NLMC Code: WA 360*

PREVALENCEOFBOVINEMASTITISINLACTATINGCOWSANDITSPUBLICHEALTHIMPLICATIONSINSELECTEDCOMMERCIALDAIRYFARMSOFADDISABABA

Strictly as per the compliance and regulations of:

© 2015. Alebachew Tilahun & Alemu Aylate. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Prevalence of Bovine Mastitis in Lactating Cows and its Public Health Implications in Selected Commercial Dairy Farms of Addis Ababa

Alebachew Tilahun [α] & Alemu Aylate [σ]

Abstract- A cross sectional study was conducted in Addis Ababa from October 2011 to May 2012 to determine prevalence of bovine mastitis and discuss its public health implications. A total of 444 systematically selected lactating cows of different cattle breed from thirty seven (37) dairy farms were investigated. The herds were visited and the farmers interviewed about the management, housing, feed and feeding, and milking conditions. California Mastitis Test (CMT) was applied on milk samples collected from 1776 individual quarters. The overall prevalence of bovine mastitis was 68.0% (302/444) of which sub-clinical mastitis accounted for 46.8% (208/444) while 21.2% (94/444) were found to be clinical forms. There was significant difference ($P<0.05$) in the prevalence of mastitis among the different breeds, age groups, parity and lactation stage. Relatively higher number of farmers interviewed (20.8%) replied that they do not withhold milk from cows treated for mastitis and continue to avail it to the public without interruption. As it is economically damaging, the need to establish diagnostic facility to enable early detection for screening large number of samples was emphasized. Further work on identification of the causative agents and conducting public awareness creation about major zoonotic diseases were also recommended.

Keywords: california mastitis test, interview, prevalence, mastitis, zoonotic.

I. Introduction

Despite many years of research, mastitis subclinical remains the most economically damaging and zoonotic potential disease for dairy industry and consumers worldwide irrespective of species of animal (Ojo *et al*., 2009). Economic losses caused by mastitis include value of discarded milk (Radostits *et al*., 2007). Bacterial contamination of milk from affected cows may render unsuitable for human consumption by causing food poisoning or interference with manufacturing process or in rare cases provides mechanism of spread of disease to humans. Zoonotic diseases potentially transmitted by raw cow milk include brucellosis, caseous lymphadenitis, leptospirosis, listeriosis, melioidosis, Q-Fever, Staphylococcal food poisoning, toxoplasmosis and tuberculosis (Mungube *et al*., 2005; Radostits *et al*., 2007).

The prevalence of subclinical mastitis in dairy herds is often surprising to producers, moreover, sub-clinically infected udder quarters can develop clinical mastitis and the rate of new infections can be high (Zdunczyk *et al*., 2003). Previous studies conducted in different countries indicate the distribution and economic importance of the disease. Contreras *et al*. (1997) from Spain; Moshi *et al*. (1998) from Tanzania; Ameh and Tari (2000) from Nigeria; Ndegwa *et al*. (2000) from Kenya and Kozacinski *et al*. (2002) from Croatia reported different prevalence rates of mastitis in dairy cattle. The disease has been reported by several authors in different parts of the Ethiopian country (Mungube *et al*., 2005; Lakew *et al*., 2009; Gebreyohannes *et al*., 2010; Megersa *et al*., 2010). Several of these studies have been shown the occurrence a range of mastitis causing bacteria indicating *Staphylococcus*, *Escherichia coli* and *Streptococcus* as dominant and pathogenic species. Some authors (Mungube *et al*., 2005) reported a substantial economic loss in Ethiopian highland crossbred dairy cows due to subclinical mastitis.

Subclinical mastitis can be recognized indirectly by several diagnostic method including the California mastitis test (CMT), the Modified White Side test, Somatic cell count, pH, and catalase tests. These tests are preferred to be screening tests for subclinical mastitis as they can be used easily, yielding rapid as well as satisfied results (Joshi and Gokhale, 2006).

In some parts of Ethiopia, the disease is insufficiently investigated and information relating to its magnitude, distribution and risk factors is scant. Such information is important to envisage when designing appropriate strategies that would help to reduce its prevalence and effects (Mekebib *et al*., 2009; Megersa *et al*., 2010).

This study aimed (i) to evaluate the prevalence of subclinical mastitis in apparently healthy dairy cows in Holeta district, (ii) to determine the most frequency of intramammary infection, causative agents, and (iii) to evaluate associated risk factors affecting on subclinical mastitis.

II. Materials and Methods

a) Study area

The study was conducted in Addis Ababa city administration, the capital of the Federal Democratic

Author α σ: School of Veterinary Medicine, Wolaita Sodo University, Wolaita Sodo, Ethiopia. e-mail: wsusvm2014@gmail.com

© 2015 Global Journals Inc. (US)

Republic of Ethiopia. The city covers an area of 530.14 km^2 and is sub divided into ten sub-cites namely, Arada, Bole, Addis Ketema, Nefas Silk Lafto, Kolfe Keranio, Akaki Kality, Yeka, Lideta, Kirkos and Gulele sub-cites. Addis Ababa lies at an altitude of 2000-3000 meters above sea level and is a grass land biome located between 9.03 North latitude and 38.74 East longitudes. The city has alternating dry and rainy seasons with the long rainy season that extends from June to September and short rainy season that lasts from March to May. The mean annual minimum and maximum temperatures range between 14°C and 21°C respectively with an overall overage of 17°C. The mean relative humidity is 61.3% (CSA, 2003).

b) Study Animals and Sample Size Determination

The study was conducted on 444 lactating cows (local, Holstein-Friesian, Jersey and cross breeds) from 37 dairy farms in Addis Ababa. The farms were purposively selected based on the availability of lactating cows within the farm and the owners' willingness. Systematic random sampling method was applied for the selection of individual animals (lactating cows) in the farms. The sample size was determined by the formula given by Thrusfield (2007) considering an expected prevalence of 71% (Mekibib *et al*., 2009), 95% confidence level and 5% desired precision. Adding a few more samples to improve on the accuracy, a total of 444 lactating cows were considered for the study.

c) Study Design

A Cross sectional study was conducted. Three dairy farms were purposively selected for their ease accessibility. Simple random sampling technique was followed to select the study animal and the desire sample size was calculated according to the formula given by Thrusfield (2007).

The study was carried out from November 2011 to April 2012 by collection of events associated with mastitis in lactating cows from 37 small holder's dairy farms in Addis Ababa.

d) Study Methodology

i. *Clinical inspection of udder*

The udder was first examined visually and then by palpation to detect possible fibrosis, inflammatory swellings, visible injury, tick infestation, atrophy of the tissue and swellings of supra mammary lymph nodes. The teat condition (color changes, swelling at or near the teat base, swelling or firmness at or near the teat end, openness of the teat orifice, teat skin condition, signs of vascular damage like petechial hemorrhage, etc.) was evaluated during clinical examination (More, 1989). Upon palpation, one can feel hot, painful swelling on udder and ventral abdomen and was manifested by loss of appetite, depression, recumbence and blood mixed milk in acute mastitis. In chronic mastitis, continuous or intermittent discharge of pus, clots, flakes or watery secretion will be seen from the udder (Chauhan and Agarwal, 2006).

e) California mastitis test (CMT)

The California Mastitis Test (CMT) was performed according to the manufacturer's instruction. In brief, a small sample of milk (approximately ½ teaspoon) was collected from each quarter into a plastic paddle that has 4 shallow cups marked A, B, C and D. An equal amount of CMT reagent was added to the milk and the paddle rotated to mix the contents. After approximately 10 seconds, the score was read while continuing to rotate the paddle. Results were recorded as T (trace), 1, 2 or 3 based on the level of precipitation (coagulation) (Mellenberger and Carol, 2000).

i. *Risk factor assessment*

Information on animal and farm-based risk factors was collected in two separate pre-designed questionnaires, by observation, and by interviewing of the different farm attendants and owners. A check-list was used to record such information as the cows' age, breed, parity, lactation stage, and body condition, problems of leaking milk and previous history of mastitis. Farm-based risk factors considered were teat drying, teat cleaning, floor types, teat dipping, milkers, bedding and treatment history.

ii. *Assessment of public health risks*

This was done by asking respondents weather they adapt the behavior of boiling milk before consumption, stripping of the foremilk at the start of milking, and by asking them the time duration of time they withheld milk before distribution to the public if the animals were treated for mastitis.

f) Statistical analysis

The data was compiled and analyzed with SPSS statistical package version 17. Prevalence estimation of commonly isolated pathogens in Holeta town dairy farms was determined using standard formulae (i.e., the number of positive animals/samples divided by the total number of animals/samples examined). Descriptive statistics such as percentages and frequency distributions was used to describe/present the nature and the characteristics of the data.

III. Results

a) Prevalence of Mastitis at Individual Cow and Quarter Level

Three hundred forty three Holstein-Friesian (HF), 20 Jersey, 15 local and 32 cross (HF X Local) breeds were included in the study. Of the total 444 lactating cows, 302 (68%) were found to be affected with clinical or sub clinical mastitis based on clinical examination of the udder and CMT results. From these, 94 (21.2%) was clinical and 208 (46.8%) was sub-clinical mastitis (Table 1). Out of 1776 quarters examined, clinical, non-functional and sub-clinical abnormalities

© 2015 Global Journals Inc. (US)

were found in 288 (16.2%), 55 (3.1%), 456 (25.7%) quarters respectively.

Table 1 : Prevalence of mastitis

Types of mastitis	Total number examined	Positive (%)	χ^2	P- Value
Clinical	444	94 (21.20)		
Subclinical	444	208 (46.80)	52.078	0.000
Total	444	302 (68.00)		

b) Prevalence of Bovine Mastitis across Different Categories of Cows

Breed, age, parity and lactation stages have significant influence ($P<0.05$) on the prevalence of bovine mastitis. There was a significant difference in prevalence between animals of different age categories ($P<0.05$). The highest prevalence (86.5%) was found in lactating cows of ages 7-10 years, followed by cows of ages 11-13 years (81.8%), and the lowest prevalence (59.1%) was recorded in cows of ages 3-6 years. Higher prevalence (90.8%) was recorded in cows which gave birth to 4-7 calves and the lower prevalence (61.6%) was recorded in cows that gave birth to 1-3 calves. The difference was statistically significant ($P<0.05$) (Table 3).

The effect of lactation stage on the current prevalence of mastitis was studied and analyzed and the result revealed that lactation stage had significant effect ($P<0.05$) on the prevalence of mastitis. Higher prevalence (89.3%) of mastitis was observed and recorded in cows of late lactation stage (9-14 month) followed by cows in mid (83.65%) lactation (5-8 month) and early lactation stage (3 week-4 month) that had a prevalence of 50.7%. The effect of breed on the prevalence of mastitis was also studied and analyzed and the result revealed that breed had significant effect ($P<0.05$).

Among the different breeds studied, the highest mastitis prevalence was observed in Holstein-Friesian breeds (71.8%) followed by Jersey (70.0%), local (66.7%), and cross (48.5%) breeds (Table 3).

Table 3 : The prevalence of bovine mastitis in association with potential predisposing factors

Variables	No. examined	Positive (%)	$\chi2$	P-Value
Breed				
Holstein-Friesian (HF)	343	246 (71.7)		
Jersey	20	14 (70.0)	13.786	.003
Cross (local x HF)	66	32 (48.5)		
Local	15	10 (66.7)		
Total	444	302 (68)		
Age				
3-6 years	296	175 (59.1)		
7-10 years	126	109 (86.5)	32.497	.000
11-13 years	22	18 (81.8)		
Total	444	302 (68)		
Parity				
1-3	346	213 (61.6)	30.048	.000
4-7	98	89 (90.8)		
Total	444	302 (68)		
Lactation stage				
3 week-4 month	223	113 (50.7)		
5-8 month	146	122 (83.65)	62.722	.000
9-14 month	75	67 (89.3)		
Total	**444**	**302 (68)**		

c) Questionnaires Survey, Observation and Interviewing

Questionnaires were distributed to 24 farms among the 37 farms included in the study. One questionnaire per farm owner/attendant was distributed. The entire farms included in the study followed manual milking (hand milking) system and most (80%) of the milkers were males. No specific sequence is followed during milking in many (87.5%) of the farms. Rather, it depends on the placement of the animal in the shed. Fifty four percent of the farm owners were educated to

© 2015 Global Journals Inc. (US)

high school level while 12.5% were educated up to university level. The remaining (33.5%) attended elementary schools. Overall, educated people had better know how about the zoonotic implications of consuming raw milk, predisposing factors for mastitis and drug residue effect post treatment of mastitic animals. A few (12.5%) farmers emphasized the need to milk healthy cows first and the diseased cows later to prevent transmission of disease. Most (66.7%) of the milkers used disinfectant before milking only while 8 (33.3%) milkers said that they use disinfectant both before and after milking. Tap water is the primary source of water to clean teats and hands in many (91.7%) of the farms while few (8.3%) milkers use river water for teats and hands cleansing.

Eighteen (75%) farms strip the foremilk first while few undertake direct milking to the material used for milking. Among the 24 farms, 8 (33.3%) used individual towels, 10 (41.7%) communal towel and 6 (25%) did not use towel for drying of teats before or after milking. Among the 24 farms, 14 (58.3%) milkers disinfect their hands before proceeding to milk the next cow while 10 (41.4%) milkers disinfect their hands only at the beginning of milking. Most (75%) of the farmers boil milk before consumption while few (25%) milkers consume raw milk. In almost all of the farms, `animals were previously treated for mastitis while few animals (heifers that gave the first and second calf) were not treated for mastitis cases. Few (20.8%) farms distribute the milk for public consumption starting from the same day the animals were treated while most (79.2%) withhold the milk depending on the withdrawal period of the drug as prescribed by veterinarians. The management (housing, bedding, feeding, etc.) and the degree of sanitation were also observed. Among the 24 farms, there were leakage of urine, feces and milk during milking in 7 (29.2%) while in the remaining (14 farms), the bedding, housing and other degree of sanitary measures like milking procedures, use of disinfectant etc. were good.

IV. Discussion

A total of 444 dairy cows, from which 343 HF, 20 Jersey, 66 cross (HF x local), and 15 local breeds from Addis Ababa were investigated in a cross sectional study conducted between November 2011 and April 2012. The current prevalence of mastitis was 68.0%. The finding in this study is greater than that of Girma (2010), who reported 44.1% and Nibret *et al.* (2011), who reported 32.6% in different parts of Ethiopia. The high prevalence of sub-clinical mastitis may be attributed to improper milking hygiene, lack of post milking teat dipping and contact labors used, absence of order in milking cows of different ages and milking of mastitic animals before the healthy ones all of which might have increased the prevalence (Radostits *et al.,* 2007). The quarter level prevalence was 41.9% (744/1776). This finding was greater than that of Benta *et al.* (2011), who reported 31.4% (349/1112). This difference in the observed prevalence of mastitis among studies may be attributed to various factors like management, environmental, animal risk factors and causative agents (Radostits *et al*., 2007).

This study revealed a higher prevalence of sub-clinical mastitis (46.8%) than clinical mastitis (21.2%). A similar result was found by Mekibib *et al*. (2009) who reported 22.4% and 48.6% for clinical and sub-clinical mastitis respectively, in dairy farms of Holeta town, central Ethiopia. In case of sub-clinical mastitis, the cow-level prevalence (46.8%) obtained in this study was greater than the finding by Girma (2010) and Nibret *et al.* (2011) who reported 33.8% and 31.67%, respectively, in different parts of Ethiopia. However, it was lower than that reported by a study carried out in Sudan (88.1%) (Abdelrahim *et al.*, 1989). This may be attributed to the difficulty of detecting sub-clinical mastitis by the owners compared to the easily detectable clinical cases which prompt owners seek treatment for their animals (Radostits *et al*., 2007).

Increasing age, lactation stage, parity and poor management increased the risk of mastitis. This is line with previous reports on mastitis in Ethiopia (Kerro Dego and Tareke, 2003) and industrialized countries (Schukken *et al.*, 1989). Stage of lactation was a risk factor for mastitis (Mungube *et al*., 2004). In late lactation the risk of mastitis increased. Two reports on Ethiopian conditions found higher prevalence of mastitis during early lactation than late lactation (Hussien, 1999). The reason may be due to excluding of lactating cows below 3 weeks to avoid false positive result since SCC increases during early lactation (Tesfu *et al*., 1999).

Manual milking methods in the entire farms that included in this study was the major predisposing factors to increase the prevalence of mastitis. Most of employed milkers have little educational background and have limited knowledge about the mechanism/s of disease transmission. Often, they do not disinfect their hands and teats during and between milking of different cows, use of communal towel for drying of teats and also, they have no special preference between tape and river water. This study also noted a high prevalence of mastitis in farms that use river water for sanitation. Sequence of milking cows also seemed to have a role on the prevalence of mastitis. For example, in farm A which employ a specific sequence (first milk healthy heifers, healthy cows and last diseased cows), the prevalence was lower as compared to the other farms in which they apply random milking procedures in the placement of cows in the shed.

In this study, 33.3% of the farm attendants reported to consume raw milk. This practice can be said as risky as raw milk can contain a variety of disease-causing pathogens, as demonstrated by numerous

© 2015 Global Journals Inc. (US)

scientific studies. These studies, along with numerous milk borne out breaks, clearly demonstrated the risk associated with drinking raw milk. For instance, in the US alone, there were 85 reported outbreaks of human infections over the years 1998-2008 due to the consumption of contaminated milk, 1614 illness, 187 hospitalizations and two deaths (Thorne, 2011).

There is also concern that small amounts of certain antimicrobial agents (residue) may significantly shift the resistance patterns in the microbial population in human intestinal tract, allergy from residue of penicillin etc. (Jones, 1999). The present study also found reluctance in 79.2% of the farm owner's to withhold milk from mastitic cows after treatment. Pasteurization effectively kills raw milk pathogens without any significant impact on milk nutritional quality. Stripping of the foremilk is also necessary as it contains many microbes that affect human health negatively.

V. Conclusion and Recommendations

In a spite of a large research efforts aimed to gain prevalence and to develop a new control tools for mastitis, the subclinical occurrence of the mastitis remains a substantial problem for dairy producers. The result of the present study indicated a relatively high prevalence of subclinical mastitis in dairy cattle of the study area. The relatively high prevalence reported in this study was clearly indicated lack of strategic control measures against the disease as well as poor surveillance measures. Lack of maintenance of strict hygiene and good sanitary environment may be contributory factors in the cause of subclinical mastitis. It is therefore important that farmers should ensure strict personal hygiene and that of animals and general sanitary condition of the farms should be improved and maintained. Furthermore, all dairy producers know that early detection of intramammary infection is important for selecting and implementing proper therapy. Unfortunately, most infections are not detected until they become clinical, and by then extensive and costly damage can result. Routine milk cultures should be an ongoing part of any mastitis control program. The sampling strategies for any ongoing program require the input of the herd veterinarian as well as herd management.

Conflict of Interest

The authors have no declared any conflict of interest

VI. Acknowledgements

Authors would like to thank Aklilu Lemma Institute of Patho-biology for their material and technical supports.

References Références Referencias

1. Abdelrahim, A. I., Shommein, A. M., Suliman, H. B. and Shaddad, S. A. I. (1989). Prevalence of mastitis in imported Friesian cows in Sudan. *Revue Elev. Med. Vet. Pays Trop*. **42**: 512-514.
2. Ameh JA, Tari LS (2000). Observation on the prevalence of caprine mastitis in relation to predisposing factors in Maiduguri. Small Ruminant Res., 35: 1-5.
3. Azmi, D., Hawari, F., Dabbas, A. I. (2008). Prevalence and distribution of mastitis pathogens and their resistance against antimicrobial agents in dairy cows Jordan. *American Journal of Animal and veterinary sciences*, **3** (1): 36-39.
4. Barnouin J, Bord S, Bazin S, Chassagne M (2005). Dairy management practices associated with incidence rate of clinical mastitis in low somatic cell score herds in France. Journal of Dairy Sci., 88: 3700–3709.
5. Bekele T, Molla B (2001). Mastitis in lactating camels (Camels dromedarus) in Afar Region, Northeast Ethiopia. Berl. Munch Tieriaztl Wochenschr., 114: 169- 172.
6. Benta, D. B. and Habtamu, T. M. (2011). Study on Prevalence of Mastitis and its Associated Risk factors in Lactating Dairy cows in Batu and its Environs, Ethiopia. *Global veterinarian* . **7**(6): 632-637.
7. Biffa D, Debela E, Beyene F (2005). Prevalence and Risk factors of mastitis in lactating dry cows in southern Ethiopia. Inter. J. of Applied Research in Veterinary Medicine., 3(3): 189-198.
8. Bitew M, Tafere A, Tolosa T (2010). Study on bovine mastitis in dairy farms of Bahir Dar town and its environ. J. Anim. Vet. Adv., 9: 2912-2917.
9. Central statistical Authority (CSA). (2003). In Ethiopia Livestock Estimate, vol. 1, Bulletin, and No. 52. Addis Ababa, Ethiopia.
10. Chauhan, R. S. and Agarwal, D. K. (2006). Textbook of Veterinary Clinical and Laboratory Diagnosis. 2ned. Jaypee Brothers Medical Publishers (P), New Delhi. Pp. 103.
11. Contreras A, Corrales JC, Sanchez A, Sierra D (1997). Persistence of Caprine intramammary pathogens throughout lactation. J. of Dairy Sci., 80: 2815-2819.
12. Coulon JB, Gasqui P, Barnoun J, Ollier A, Pardel P, Dominique P (2002). Effect of mastitis and related germ yield and composition during naturally occurring udder infections in dairy cows. Anim. Res., 51: 383–393.
13. CSA (2010). Ethiopian Agricultural Sample Enumeration (EASE), Addis Ababa, Ethiopia.
14. FAO (2003). Livestock Sector, Brief Livestock Information, Sector Analysis and Policy Branch:Rome, Italy, Pp.1-15.
15. Fekadu, K. (1995). Survey on the prevalence of bovine mastitis and the predominant causative agents in Chaffa valley. Proceedings of the 9th

© 2015 Global Journals Inc. (US)

Conference of Ethiopian Veterinary Association: Addis Ababa, Ethiopia; 101–111.

16. Gebreyohannes YT, Regassa FG, Kelay B (2010). Milk yield and associated economic losses in quarters with subclinical mastitis due to *Staphylococcus aureus* in Ethiopian crossbred dairy cows. Trop. Anim. Health Prod., 42: 925-931.
17. Girma DD (2010). Study on prevalence of mastitis on cross breed Dairy cow around Holeta areas. Global veterinaria., 5(6): 318-323.
18. Girma, D. D. (2010). Study on Prevalence of Bovine Mastitis on cross Breed Dairy cows Aroun Holeta Areas, West Shewa Zone of Oromia, Ethiopia. *Global Veterinaria* **5** (6): 318-323.
19. Girma, T. (2001). Prevalence of mastitis at Alemaya University dairy farm. *Journal Ethiopian Veterinary.*
20. Harmon RJ (1994). Physiology of mastitis and factors affecting somatic cell counts. J. Dairy Sci., 77: 2103.
21. Hogan JS, Gonzalez RN, Harmon RJ, Nickerson SC, Oliver SP, Pankey JW, Smith KL (1999). Laboratory and field handbook on bovine mastitis. Madison: National Mastitis Council, 12: 1-33.
22. Hogan SJ, Gonzalez RN, Harmon JR, Nickerson SC, Oliver SP, Pankey JW, Smith LK (1999). Laboratory Handbook on Bovine Mastitis. Published by National Mastitis Council, Inc., W D Hoard, Fort Atkinson, USA.
23. Hussein, N. (1999). Cross-sectional and longitudinal study of bovine mastitis in urban peri-urban dairy systems in the Addis Ababa region. Ethiopia, (MSc thesis, Free University of Berlin, Germany and Addis Ababa University, Ethiopia, Joint programme).
24. Iqbal M, Khan MA, Daraz B, Saddique U (2004). Bacteriology of mastitic milk and in vitro antibiogram of the isolates. Pak. Vet. J., 24: 161-164
25. Jones, G. M. Professor and Extension Dairy Scientist, Milk Quality and Milking Management, Department of Dairy Science, Virginia Tech. (1999). On-farm Tests for Drug Residues in Milk. The cattle site. Ltd.
26. Joshi S, Gokhale S (2006). Status of mastitis as an emerging disease in improved and periurban dairy farms in India. Ann. New York Acad. Sci., 1081: 74–83.
27. Kahn, C. M. (2005). The Merck Veterinary Manual. 9th edition. Merck and Co., Inc. Whithhouse Station, N. J., U.S.A. Pp. 1120-1129.
28. Karimuribo ED, Fitzpatrick JL, Bell CE, Swai ES, Kambarage DM, Ogden NH, Bryant MJ, French NP (2006). Clinical and subclinical mastitis in smallholder dairy farms in Tanzania. Risk intervention and knowledge transfer. Prev. Vet. Med., 74: 84-98.
29. Kerro Dego, O. and Tareke, F. (2003). Bovine mastitis in selected areas of southern Ethiopia. *Tropical Animal Health and Production.* **35**, 197-205.
30. Kozacinski LM, Iladziosmanovi T, Majic IK, Jole, Cvrtila Z (2002). Relationships between the results of mastitis tests, somatic cell counts and the detection of mastitis agents in milk. Paraxis Vet., 57: 255-260.
31. Lakew M, Tolosa T, Tigre W (2009). Prevalence and major bacterial causes of bovine mastitis in Asella, South Eastern Ethiopia. Trop. Anim. Hlth. Production., 41: 1525-1530.
32. Megersa B, Chala T, Abunna F, Regassa A, Berhanu M, Etana D (2010). Occurrence of mastitis and associated risk factors in lactating goats under pastoral management in Borana, Southern Ethiopia. Trop. Anim. Hlth. Production., 42: 1249-1255.
33. Mekebib B, Furgasa M, Abunna F, Megersa B, Furgasa A (2009). Bovine mastitis prevalence, risk factors and major pathogens in dairy farms of Holeta Town, central Ethiopia. Veterinary World., 13(9): 397-403.
34. Mekibib, B., Furgasa, M., Abunna, F., Megersa, B. and Regassa, A. (2009). Bovine Mastitis: Prevalence, Risk Factors and Major Pathogens in Dairy Farms of Holeta Town, central Ethiopia. *Veterinary World vol*. **3** (9): 397-403.
35. Mellenberger, R. Dept. of Animal Sciences, Michigan State University and Carol, J. Roth, Dept. of Dairy Science, University of Wisconsin-Madison April, 2000.
36. Mohammed, A., and Etana, D. (1996). Bacterial causes of mastitis in south Ethiopia, annual research report, Hawasa college of agriculture.
37. More, P. C. (1989). Tick borne Diseases of Livestock in Africa. In: Manual of Tropical Veterinary Parasitology. English version. C. A. B. International: Wallingford, UK: 331-
38. Moshi NG, Kitaro GC, Minga UM (1998). Prevalence of mastitis in dairy goats in some selected farms in Morgoro and Arusha, Tanzania. Tanzania J. Agric. Sci., 1: 173-180.
39. Mungube ED, Tenghagen BA, Regassa F, Kyule MN, Shiferaw Y, Kassa T, Baumann MPO (2005). Reduced milk production in udder quarters with subclinical mastitis and associated economic losses in crossbred dairy cows in Ethiopia. Trop. Anim. Hlth. Produ., 37(5): 503-512.
40. Mungube, E. O., Tenhagen, B. A., Kyule, M. N., Greiner, M., Baumann, M. P. O., Kassa, T. and Regassa, F. (2004). Risk Factors for Dairy Cow Mastitis in the Central Highlands of Ethiopia. *Tropical Animal Health and Production*. **36**: 463-472.
41. National Mastitis Council (2004). Microbiological procedures for the diagnosis of udder infection. 3rd ed. Arlington, VA: National Mastitis Council Inc.
42. Ndegwa EN, Mulei CM, Munyna SJ (2000). The prevalence of subclinical mastitis in dairy goats in Kenya. J. South Afr. Vet. Assoc., 71: 25-27.

© 2015 Global Journals Inc. (US)

43. Ojo OE, Oyekunle MA, Ogunleye AO, Otesile EB (2009). *Escherichi coli*, O157:H7 in Food animals in part of south-western Nigeria. Prevalence and invitro antimicrobial susceptibility. Trop. Vet., 26 (3): 23-30.
44. Okeke IN, Laxminarayan R, Bhutta ZA, Duse AG, Jenkins P, O'Brien TF, Pablos-Mendez A, Klugman KP (2005). Antimicrobial resistance in developing countries. Part I: recent trends and current status The Lancet Infectious Diseases, 5(8): 481-493.
45. Quinn PJ, Carter ME, Markey B, Carter GR (2004). Clinical Veterinary Microbiology. London Wild life Publisher. 95-101.
46. Quinn, P. J., Carter, M. E., Markey, B. K. and Carter, G. R. (1999). Mastitis. In Clinical Veterinary Microbiology, Mosby International Limited, London, and Pp. 327-344.
47. Quinn, P. J., Carter. M. E., Marky, B. K., Carter. G. R. (1994). Clinical Veterinary Microbiology. 1 ed. London: Wolfe publishing.
48. Radostits OM, Gay CC, Hinchcliff KW, Constable PD (2007). Veterinary Medicine. A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses. 10th edition, Saunders Elsevier, Spain. 1045-1046.
49. Radostits, O. M. (2007). A text book of the diseases of cattle, horses, sheep, pigs, and goats. Pp. 673-753.
50. Riekkerink RGMO, Barkema HW, Kelton DF, Scholl T (2008). Incidence rate of clinical mastitis on Canadian dairy farms. J. of Dairy Sci., 91, 1366-1377.
51. Sandholom, M. (1995). Detection of inflammatory changes in milk. In: Sandholom, M., Honkanen_Buzalski, T., Kaartinan, L., Pyorala. (eds), In: The bovine udder and mastitis Finland, Gummerus Kirjapaino Oy. Juvaskyla.Pp. 89-104.
52. Schukken, Y. H., Grommers, F. J., van de Geer, D. and Brand, A. (1989). Incidence of clinical mastitis on farms with low somatic cell counts in bulk milk. *Veterinary record*, **125**: 60-63.
53. Sears PW, Wilson DJ, Gonzalez RN, Hancock DD (1991). Microbiological results from milk samples obtained pre-milking and post-milking. J. of Dairy Sci., 74: 4183-4188.
54. Shakoor A (2005). Preparation and evaluation of *Staphylococcus aureus* vaccines for the control of mastitis in dairy buffaloes (Bubalus bubalis). PhD Dissertation, Department of Veterinary Clinical Medicine and Surgery, College of Agriculture, Faisalabad, Pakistan. 34-40.
55. Sharif A, Umer M, Muhammad G (2009). Mastitis control in dairy production. J. Agri. Soc. Sci., 5: 102-105.
56. Sori H, Zerihun A, Abdicho S (2005). Dairy Cattle Mastitis in and around Sebeta, Ethiopia. Inter. J. Applied Research in Veterinary Medicine., 3: 332-338.
57. Tefera G (2001). Prevalence of mastitis at Alemaya University Dairy Farm, J. Eth. Vet. Ass., 3: 17-21.
58. Tegegne, T., Alemayehu, A. and Ayele, G. M. (1999). Cross Border Livestock Trade and Food Security in the Southern and Southeastern Ethiopia Borderlands. *OSSREA (Organization for Social Science Research in Eastern and Southern Africa) Development Research Report Series*. No.1. Commercial Printing Enterprise: Addis Ababa, Ethiopia; 1–6.
59. Tesfu, K., Gemechu, W. and Azage, T. (1999). Survey of Mastitis in Dairy Herds in the Ethiopian Central Highlands. *SINET: Ethiop. J. Sci*., **22**(2): 291-301.
60. Thorne, R. (2011). Side effects of Dairy Raw milk. *Proud Partner of the Lance Armstrong Foundation*.**8** (5): 319-328.
61. Thrusfield M (2007). Veterinary epidemiology. 3rd edition, Blackwell Publishing.
62. Thrusfield, M. (1995). Veterinary Epidemiology. 2nd. Black well science Ltd. UK.
63. Vaarst, M. and Envoldsen, C. (1997). Patterns of clinical mastitis in Danish organic dairy herd. *J. Dairy Sci.* **64**: 23.
64. Walshe, M. T., Grindle, T., Nell, A. and Baschaman, M. (1995). Dairy Development in Sub-Saharan Africa: A study of Issues and option, World Bank Technical Paper, 135, Africa Technical Development Series, Washington D. C., The World Bank.
65. Williams, T. O., DeRosa, D. A., Badiane, O. (1995). Macroeconomic, international trade and sectoral policies in livestock development: an analysis with particular reference to low income countries. In: Wilson, R. T., Etui, S., and Mack, S. (eds). Livestock development strategies for low income countries. Proceeding of the Joint FAO/IL Roundtable on livestock development strategies for low income countries, ILRI, Addis Ababa, Ethiopia, 27 February 02 March, Pp. 45-68.
66. Workineh SM, Bayleyegne M, Mekonnen H, Potgieter LND (2002). Prevalence and etiology of mastitis in cows from two major Ethiopian dairies. J. Trop. Anim. Hlth. Prod., 34: 19–25.
67. Yohannes, A. (2003). Clinical and sub-clinical mastitis in primiparous dairy heifers in Jordan. MSc Thesis. Faculty of Agricultural technology, Al-Salt University College, Al-salt, Jordan.Yoseph, M., Azage, T., Alemu, Y. and Umunna, N. N. (1998). Reproductive management and reproductive performance of dairy herds in urban and peri-urban dairy production systems in Addis Ababa milk shed. In: proceeding of the 6th annual conference of ESAP Addis Ababa, Ethiopia, 312-315.
68. Zdunczyk S, Zerbe H, Hoedemaker M (2003). Importance of oestrogen and oestrogen active compounds for udder health in cattle. A review. Dtsch Tierarztl Wochenschr., 110: 461.

© 2015 Global Journals Inc. (US)

69. Zerihun T (1996). A Study on Bovine Subclinical Mastitis at Stela Dairy Farm. Thesis, Debrezeit: Faculty of Veterinary Medicine, Addis Ababa University: Ethiopia. 25–27.

© 2015 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2015

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSM" title to individuals.The'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.

- The "FARSM" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall,Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.

You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.

As FARSM, you will be given a renowned, secure and free professional email addres with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.

The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.

The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of you credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.

The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.

The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.

MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.

The “MARSM” is a dignified ornament which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.

MARSM accrediting is an honor. It authenticates your research activities. Afterbecoming MARSM,you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.

MARSM designated members are entitled to avail a 25% discount whilepublishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you willbe given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.

Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

Auxiliary Memberships

Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).

The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.

The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.

The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:

Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.

Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone. "

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.

To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .

Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

Index

B

Brillard · 18, 24, 26

C

Coprological · 3, 4, 5, 6, 7

E

Eljazeera · 1, 2, 11
Embryonic · 18, 21, 22, 24, 26, 30, 31

F

Fasciolosis · 1, 3, 5, 7, 8, 9, 11, 14, 16

G

Gebreyohannes · 35, 44

L

Leghorn · 18, 20, 22, 24, 26, 28, 30, 32
Leptospirosis · 34
Libitum · 20

M

Mulugeta · 6, 7, 8
Mungube · 34, 35, 41, 45

P

Phaseolus · 18, 19, 20, 22, 24, 26, 27, 28, 30, 32

V

Vulgaris · 18, 20, 22, 24, 26, 28, 30, 32

Z

Zdunczyk · 35, 46

save our planet

Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.MedicalResearchJournal.org
or email us at helpdesk@globaljournals.org

ISSN 9755896

© Global Journals