

Characterization of Maxillary Sinus in Patients with Facial Pain using Ultrasound

Caroline.E.Ayad¹ and Alsaffi Ahmed Abdalla²

¹ Sudan University of Science and Technology

Received: 11 December 2012 Accepted: 31 December 2012 Published: 15 January 2013

Abstract

This study dealt with characterization of maxillary sinus in patients with facial pain using ultrasound. Methods: It was carried out on 50 patients complaining of maxillary sinus problems who were referred to the National Center of Ear, Nose and Throat Khartoum-Sudan. The instruments used were; Aloka SSD 500, Honda HS2000 portable with curvilinear probe 3.5MHz, and General Electric LOGIQ5 Mobile with curvilinear and linear probe (3.5MHz, 10MHz). Results: patients were 27males (54

Index terms— ultrasound, pain, maxillary sinuses.

1 Introduction

sinusitis is one of the most common problems affecting the maxillary sinuses. Acute sinusitis may be caused by bacterial infection; Complications of acute bacterial sinusitis (ABRS) may include orbital, intracranial, or soft tissue involvement. Therefore, accurate diagnosis is of major importance and the radiographic imaging is considered as an accurate diagnostic tool. [1]

2 S

The American Academy of Pediatrics Clinical Practice Guideline on the management of sinusitis stated that imaging including radiographs, Computerized Tomography(CT) or magnetic resonance imaging(MRI) ;can serve only as confirmatory measures of sinus disease in patients whose clinical histories are supportive of the diagnosis [2]. The American Academy of Allergy, Asthma and Immunology stated that computed tomography (CT) is the preferred imaging technique for pre-operative evaluation of the paranasal sinuses and that ultrasonography has limited utility, but may be applicable in pregnant women and for determining the amount of retained secretions [3]. Coronal computed tomography (CT) scan of the paranasal sinuses was suggested to be considered in the diagnosis of acute sinusitis [4]. The diagnosis of both acute and chronic sinusitis should be made clinically, and not on the basis of imaging findings alone. CT remains the study of choice for the imaging evaluation of acute and chronic sinusitis, in addition magnetic resonance imaging of the sinuses, orbits, and brain should be performed whenever extensive or multiple complications of sinusitis are suspected. In chronic sinusitis, CT scanning is the "gold standard" for the diagnosis and the management, as well as when surgery is necessary. Nuclear medicine studies and ultrasound are rarely indicated in acute and chronic rhinosinusitis [5]. Diagnostic ultrasound is a noninvasive imaging modality that uses high-frequency sound waves. The great variability of test performance of diagnostic ultrasound in acute sinusitis cites systematic evidence [4]. No risks have been identified with ultrasound evaluation of the paranasal sinuses, but the accuracy of the ultrasound is dependent largely on the examiner's skills. [6] Therefore this study is to assess the role of ultrasound as another tool in diagnosis of maxillary sinuses in patients with facial pain in order to take part in the detection of maxillary sinuses disorders as well as to characterize normal echo texture of different maxillary sinuses diseases as compared to CT; and to correlate between the ultrasound findings with the patients clinical findings.

3 II.

4 Material and Methods

The study was conducted at the National Center of Ear, Nose, Throat (ENT) and Head and Neck surgery Khartoum-Sudan, in the period from August to November 2011.

5 b) Instrumentations

Ultrasound machines used were Aloka SSD 500 portable with curvilinear probe 3.5MHz, Honda HS2000 portable with curvilinear probe 3.5MHz and General Electric LOGIQ5 Mobile with curvilinear and linear probes (3.5MHz-10MHz).

6 c) Method

An approval from the review board and ethics committee of the College of Medical Radiological Science was taken. The patients were examined in sitting position; the U/S gel was applied to the area on the face where the sinuses are located, including the nose, cheekbones and maxillary sinuses. High frequency sound waves produced the image of the internal structures, and a highly ultrasound expertise Ear, Nose, Throat (ENT) specialist doctor had done the scan and interpreted the data. The patients were scanned by a frequency ranging from 3.5 to 10 MHz. The data were collected using variable patients complaints and they were diagnosed by U/S and the findings were correlated to CT findings.

To find out the accuracy, sensitivity and specificity of ultrasound(U/S) as compared to the Computed Tomography (CT) diagnosis; equations were applied including :A) For U/S accuracy (True positive cases(TP) + True negative cases(TN) / Total number of cases) X 100. B) For U/S Sensitivity(True positive cases(TP) / True positive(TP) + False negative(FN) X 100. C) for U/S specificity (True negative cases(TN) / True negative cases(TN) + False positive cases(FP) X 100.

7 III.

8 Results

The following results were for a sample consisted of 50 patients (23 females and 27 males) all were complaining of maxillary pain. The patients were classified according to age. Patients ages ranged from 20-30 years were 21(42%), 31-40 years 14(28%), 41-50 years 5(10%), 51-60 3(6%) years, 61-70 7(14%) (Figure1). The following tables showed the results according to the age, gender, complaints and the ultrasound findings.

9 Discussions

In recent decade, the Health Care Guideline (2011) stated that the diagnosis of acute sinusitis should be based on the presenting symptoms, history, and clinical examination. It noted that plain X-rays for sinus and other imaging tests are usually not necessary in making the diagnosis of acute sinusitis due to their poor sensitivity and specificity limits. Other published studies of ultrasound of the paranasal sinuses do not permit assessment of the sensitivity or specificity of the technique compared to the gold standard of CT scanning. [9] The Agency for Healthcare Research and Quality (AHRQ) evidence report on Acute Bacterial sinusitis does not address ultrasound of the sinuses [10] This study assessed the ultrasonography as another tool in the diagnosis of maxillary sinuses disorders in patients with pain in order to analyze whether ultrasonography with a reasonable degree of confidence and can replace radiography in the diagnosis of sinusitis. Fifty patients from the national center of ENT diseases and head and neck center who were previously diagnosed by CT, were investigated by ultrasound, the study showed different maxillary sinus ultrasound diagnosis related to age, gender and complaints.

The frequency distribution was according to gender (table 3), there were 27 males (54%) and 23 females (46%) out of total 50 patients (100%) with maxillary sinus pain. These results support the fact that the males were more affected than females.

The study showed the frequency and percentage regarding to age group (Figure1), the most affected age group was the group (20-30) years constituting 42%.

The frequency and percentage of the patient's complaints (table1) revealed that 50/50 (100%) complain of pain followed by stuffy nose 30/50 (60%) as compared to the other complaints.

The frequency distribution and percentage of ultrasound diagnosis (table 2), clarifies that the normal U/S appearance was found in 19/50 constituting 38%, followed by polyps, mucosal thickening and cysts.

Ultrasonography character and findings of the maxillary sinuses were 30% polyps with hypo echoic cavity, 12% fluid with hypo echoic cavity with posterior margin was seen, 2% cyst with hypo echoic area with well defined borders, 12% mucosal thickening, 38% normal sonographic appearance of hyper echoic anterior wall and cavity and the posterior margins were not seen, 6% polypoid mucosal thickening with hypo echoic cavity and echogenic bony walls.

From the study, the patients suffered from maxillary pain represented (100%), halitosis (56%), stuffy nose (60%), postnasal drip (58%), fever (54%), and malaise (34%).

97 When comparing the Ultrasound findings with the CT findings; the study showed that the ultrasound has
98 Sensitivity of 76.6% and specificity of 92.2%
99 Regarding the results and comparing to what was mentioned by (AHRQ); Ultrasound is proposed for
100 demonstrating mucosal wall thickening, focal soft tissue masses, and complex collections in the paranasal sinuses.
101 Ultrasound is a painless non-invasive diagnostic procedure and no risks have been identified with ultrasound
102 evaluation of the paranasal sinuses, but the accuracy of the ultrasound is dependent largely on the examiner's skills
103 and to achieve this purpose; this study was done by a highly experience ENT specialist with ultrasonography of
104 excellent performance. The study concluded that paranasal sinus ultrasound has been proposed as best diagnostic
imaging modality to confirm the diagnosis of clinical sinuses diseases.¹



Figure 1: Figure 1 :

¹© 2013 Global Journals Inc. (US)




Figure 2: Table 1 :

2

Ultrasound diagnosis	Frequency (%)
Fluid	6 (12%)
Polyp	15 (30%)
Cyst	1 (2%)
Mucosal Thickening	6 (12%)
Normal	19 (38%)
Polyploidy mucosal thickening	3 (6%)
Total	50 (100%)

Figure 3: Table 2 :

3

Gender	Fluid	Mucosa Thickening	Normal Polyps	Cyst	Polyploidy mucosal thickening	
Male	4	2	12	9	0	0
Female	2	4	7	6	1	3
Total	6	6	19	15	1	3

Figure 4: Table 3 :

4

Author	Ultrasound	Ultrasound	Modalities used	Number of Patients
	Sensitivity	Spesifity		
Current Study	76.6%	92.2%	CT and Ultrasound	50
Apostolos H. et al [7]	66.7%	94.9%	CT and Ultrasound	56
Fufezan et al [8]	94.9%	98.4	X-Rays and Ultrasound	67

[Note: © 2013 Global Journals Inc. (US) Characterization of Maxillary Sinus in Patients with Facial Pain using Ultrasound IV.]

Figure 5: Table 4 :

-
- [Ip et al. (2005)] *Agency for Healthcare Research and Quality. Update on acute bacterial rhinosinusitis. Health Technology Assessments*, S Ip , E Fu , Balk , Chew , J Devine , Lau . <http://www.ahrq.gov/downloads/pub/evidence/pdf/rhinoupdate/rhinoup.pdf>. Accessed on 2005 June. February 7, 2012. (No 05-E 020-2. Available at)
- [Bhattacharyya ()] 'Clinical and symptom criteria for the accurate diagnosis of chronic rhinosinusitis'. N Bhattacharyya . *Laryngoscope* 2006. 116 (7) p. . (Pt. Suppl 110)
- [Clinical practice guideline: Management of sinusitis American Academy of Pediatrics ()] 'Clinical practice guideline: Management of sinusitis'. *American Academy of Pediatrics* 2001. 108 (3) p. . (Pediatrics)
- [Varonen et al. ()] 'Comparison of ultrasound, radiography, and clinical examination in the diagnosis of acute maxillary sinusitis: a systematic review'. H Varonen , M Mäkelä , S Savolainen , E Läärä , J Hilden . *Journal of Clinical Epidemiology* 2000. 53 (9) p. .
- [Diagnosis and treatment of respiratory illness in children and adults. ICSI Health Care Guideline (2011)] *Diagnosis and treatment of respiratory illness in children and adults. ICSI Health Care Guideline*, http://www.icsi.org/respiratory_illness_in_children_and_adults_guideline_respiratory_illness_in_children_and_adults_guideline_13116.html. Accessed on 2011. February 7, 2012. (Institute for Clinical Systems Improvement (ICSI))
- [Reider et al. (2003)] 'Do imaging studies aid diagnosis of acute sinusitis?'. J M Reider , J Nashelsky , J Neher . *J Fam Pract* 2003 Jul. 52 (7) p. .
- [Triulzi and Zirpoli ()] 'Imaging techniques in the diagnosis and management of rhinosinusitis in children'. F Triulzi , S Zirpoli . *Pediatr Allergy Immunol* 2007. 18 p. . (Suppl)
- [Karantanias and Sandris (1997)] 'Maxillary sinus inflammatory disease: Ultrasound compared to computed tomography'. Apostolos H Karantanias , Vasilios Sandris . *Comput Med Imaging Graph* 1997 Jul-Aug; 21. (4) p. .
- [Spector et al. ()] 'Parameters for the diagnosis and management of sinusitis'. S L Spector , I L Bernstein , J T Li , W E Berger , M A Kaliner , D E Schuller . *J Allergy Clin Immunol* 1998. 102 (6) p. . (Pt 2)
- [Fufezan et al. (2010)] 'The role of ultrasonography in the evaluation of maxillary sinusitis in pediatrics'. O Fufezan , C Asavoia , Chereche? Panta , P Mihu? , G Bursa?iu , E . *Med Ultrason* 2010 Mar. 12 (1) p. .