Online ISSN: 2249-4618
Print ISSN: 0975-5888

Global Journal of Medical Research: B

Pharma, Drug Discovery, Toxicology and Medicine



Diabetic Patients

Toona Ciliata Leaves

Highlights

Anti-Ulcer Activity

Activity of Rhus Coriaria

Discovering Thoughts, Inventing Future

VOLUME 14 ISSUE 7 VERSION 1.0

© 2001-2014 by Global Journal of Medical Research , USA

Volume 14 Issue 56 was originally issued as B7

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine

Volume 14 Issue 56
Originally issued as B7

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine

Volume 14 Issue 7 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Medical Research . 2014.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric SciencesPurdue University Director
National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Issue

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue

1. Anti-Ulcer Activity of Rhus Coriaria in Indomethacin and Water Immersion Restraint Induced Gastric Ulcer in Experimental Rats. *1-7*
2. Antidepressant and Anti-Inflammatory Activities of Cationic Amphiphilic Complexes of Sulphadiazine. *9-16*
3. In Vivo Anti-Inflammatory and in Vitro Antioxidant Activities of Toona Ciliata Leaves Native to Bangladesh. *17-25*
4. Glycemic Control and Self-Care Practice among Ambulatory Diabetic Patients in Ambo General Hospital, West Showa, Ethiopia. *27-36*
5. Assessment of Knowledge, Attitude and Practices Regarding Life Style Modification among Type 2 diabetic Mellitus Patients Attending Adama Hospital Medical College, Oromia Region, Ethiopia. *37-48*

v. Fellows and Auxiliary Memberships
vi. Process of Submission of Research Paper
vii. Preferred Author Guidelines
viii. Index

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine
Volume 14 Issue 7 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Anti-Ulcer Activity of Rhus Coriaria in Indomethacin and Water Immersion Restraint Induced Gastric Ulcer in Experimental Rats

By Haqeeq Ahmad, Abdul Wadud, Nasreen Jahan & Izharul Hasan

Rajiv gandhi university, India

Abstract- The anti-ulcer activity of hydro alcoholic extract of Rhus coriaria Linn (HAERC) was investigated in Indomethacin and Water immersion-induced restraint gastric ulcer in Wistar rats. The assessment was carried out by using ulcer index, ulcer score and histopathological studies of the specimens. HAERC at doses of 145 and 248 mg/kg given orally produced significant inhibition of the gastric lesions induced by Indomethacin and Water Immersion restraint method and the results were comparable to the standard treatment regime. We observed that Rhus coriaria Linn extract exhibits significant anti ulcer activity and thus supports the Unani claims about the drug.

Keywords: rhus coriaria; indomethacin; water immersion induced ulcer model; ulcer index; post sumaq.

GJMR-B Classification : NLMC Code: WR 598, WI 310



Strictly as per the compliance and regulations of:

© 2014. Haqeeq Ahmad, Abdul Wadud, Nasreen Jahan & Izharul Hasan. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Anti-Ulcer Activity of Rhus Coriaria in Indomethacin and Water Immersion Restraint Induced Gastric Ulcer in Experimental Rats

Haqeeq Ahmad [α], Abdul Wadud [σ], Nasreen Jahan [ρ] & Izharul Hasan [ω]

Abstract- **The anti-ulcer activity of hydro alcoholic extract of Rhus coriaria Linn (HAERC) was investigated in Indomethacin and Water immersion-induced restraint gastric ulcer in Wistar rats. The assessment was carried out by using ulcer index, ulcer score and histopathological studies of the specimens. HAERC at doses of 145 and 248 mg/kg given orally produced significant inhibition of the gastric lesions induced by Indomethacin and Water Immersion restraint method and the results were comparable to the standard treatment regime. We observed that Rhus coriaria Linn extract exhibits significant anti ulcer activity and thus supports the Unani claims about the drug.**

Keywords: rhus coriaria; indomethacin; water immersion induced ulcer model; ulcer index; post sumaq.

I. Introduction

Peptic ulcer disease (PUD) is one of the most common gastro intestinal disorders, which causes a high rate of morbidity. An estimated 15,000 deaths occur each year because of PUD. The prevalence of duodenal ulcer is dominant in western population whereas gastric ulcer is more frequent in most Asian countries[1]. The lifetime prevalence of peptic ulcer disease is 5 to 10% with about equal prevalence in men and women. The incidence of ulcer increases with age because of excessive use of NSAIDs and the reduction in the tissue prostaglandins [2]. In India, PUD is common and the Indian pharmaceutical industries share 6.2 billion rupee and occupy 4.3% of the market share in consuming the antacids and antiulcer drugs [3]. Peptic ulcer which is usually an asymptomatic gastrointestinal disorder defined as a breach in the mucosa of the alimentary tract, which extends through the muscularis mucosa into the submucosa or deeper. Peptic ulcer disease commonly occurs when the linings of stomach or proximal duodenum are corroded by the acid-peptic juices which are secreted by the stomach cells [4,5]. Peptic ulcer is caused by *H. pylori* infection, long term and high doses of drugs such as non steroidal anti-inflammatory drugs (NSAIDs), diseases like Zollinger-Ellison syndrome, other factors such as smoking; emotional stress and excessive alcohol consumption also may contribute. In Unani Medicine, gastric ulcer is known as *Qarah-e-Medah*. Unani scholars mentioned its causes as, *Khilte Haad* (hot and irritant humour), Fuzlat (waste products), *intake* of hot and spicy foods, excessive intake of rotten food, alcohol and hard fibrous diet, desensitization of internal surface of stomach which causes excessive gastric secretions, chronic gastritis and indigestion, prolonged stress and strains and unabsorbed gastric secretions. The modern approach to control gastric ulceration is to inhibit gastric acid secretion, to promote gastro protection, to block apoptosis and to stimulate epithelial cell proliferation for effective healing [6]. Hence, Conventional medicine treats Peptic ulcer by proton pump inhibitors (PPI), H2-receptor antagonist, antacids and antibiotics for *H. pylori.* However, there are reports of adverse effects and relapse in the long run[7] that lead people to find the alternative medications. Furthermore, many of these drugs do not fulfill all the *beneficial necessities* [8]. The clinical evaluation of these drugs showed development of tolerance and incidence of relapse and side effects that make their efficacy debatable. This has been the rationale for the development of new, safe antiulcer drugs. Herbal drugs can provide lead for the development of such antiulcer drugs because these drugs are considered safer in view of their natural ingredients. In recent times, focus on plant research has increased all over the world and a large body of evidence has been collected to show immense potential of medicinal plants used in various traditional systems of medicine. More than 13,000 plants have been studied during the last few years [9].

Unani physicians in the treatment of gastritis, gastric ulcer and associated disorders due to its stomachic, astringent, desiccant, styptic, sedative and coolant activities [10] also use Post Sumaq (Fruit rind of Rhus coriaria Linn.) frequently. However, there is no scientific report regarding its efficacy in PUD. Therefore, the present study was carried out to examine the effect of Post Sumaq in gastric ulcer on animal model.

II. Material and Methods

Institutional Animal Ethics Committee (IAEC) of NIUM approved the present study. The test drug *Sumaq*

Author α σ ρ : Dept of Ilmul-Advia (Pharmacology), National Institute of Unani medicine, Bangalore-91, Karnataka India.
Author ω : Dept of preventive and social medicine, A & U Tibbia College and Hospital, New Delhi 05.

© 2014 Global Journals Inc. (US)

(Rhus coriaria Linn) was procured from local market of Bangalore, and was identified by Dr. H.B. Singh, Chief Scientist and Head of National Institute of Science Communication and Information Resources (NISCAIR) New Delhi, vide Reference No. NISCAIR/RHMD 2030/38.

III. Preparation and Dose of the Test Drug

The fruits of test drug were dried in shade, the Post (rind) was peeled off, and its therapeutic dose (5gm) as in Unani medicine was used to calculate the dose for experiment [11]. Thus, dose was found to be 580 mg /kg. Since, the test drug was studied at two different doses; therefore a second dose was also calculated by the method of Miller and Tainter (1944) [12] and found to be 990 mg/kg. As the hydro alcoholic extract was used for the study, the dose of the extract was calculated with reference to the dose of crude drug after obtaining the 25% yield percentage of extract. The hydro alcoholic extract of the drug was used in the dose of 145mg/kg and 248mg/kg. Standard drug, Omeprazole (Manufactured in India by Dr Reddys Laboratories Ltd. Village Manuja Thana) was used in the dose of 20mg/kg.

IV. Animals

The study was carried out in healthy Wistar rats of either sex, weighing 150-250 gm. The animals were procured from Biogen Laboratory Animal Facility (Reg. No. 971/bc/06/CPCSEA), a registered breeder in Bangalore. They ware acclimatized to the laboratory condition for 7 days before the experimental studies. The rats were housed in polypropylene cages under controlled conditions of light (12/12) and temperature (23±20C) under strict hygienic conditions. The animals were given Standard food pellets (Hindustan Lever Ltd.) and tap water ad libitum.

V. Induction of Gastric Ulcer

This test was carried out by the method described by Vogel[13] with minor modification in the treatment schedule. The animals were divided into 8 groups of 6 animals each. The animals in group I were administered with distilled water throughout study and served as Plain control and after 36 hours they were sacrificed while the animals in group II (after 24 hours of fasting) were treated with Indomethacin 20 mg/kg, once daily, orally for 5 days and served as negative control. The animals in group III, IV and V were treated with standard drug Omeprazole and hydro alcoholic extract of test drug in the dose 20 mg/kg, 145mg/kg and 248 mg/kg, respectively and served as pre-treated standard, Pre-treated test group A and Pre-treated test group B, respectively. These treatments were continued for five days; however, on 6th day after 24 hours of fasting ulcer was induced by the administration of Indomethacin in the dose of 20 mg/kg, for the next five days. Food was withdrawn for two hours after Indomethacin administration. On 5th day after 12 hours of fasting, the animals were treated with the last dose of Indomethacin and after five hours of administration of Indomethacin, the animals including negative control were sacrificed. While in Post treated standard and test groups the animals were first kept on fasting for 24 hours and ulcer induced by the administration of Indomethacin in the dose 20 mg/kg, daily for 5 days, thereafter the animals were treated with standard and test drug for next 5 days in the same dose and same manner as described above. On 6th day after 12 hours of fasting, the animals were sacrificed.

The water immersion restraint induced gastric ulcer was done by the method of Hayaso and Takeuchi[14]. The animals in this model were also divided in to 8 groups of 6 animals each. The animals in Group I and II were treated with distilled water and were serve as Plain control and Negative control, respectively. While the animals in Group III, IV and V were treated with standard drug, and hydro alcoholic extract of the test drug in the dose of 20 mg/kg, 145mg/kg and 248mg/kg and served as Pre-treated standard, Pre-treated test group A and Pre-treated test group B, respectively. All the animals were treated in this way once daily for 5 days. They had free access to food and water during the treatment period. However, on 4th day they were kept on fasting for 12 hours with water ad libitum. On 5th day, 12 hour fasted rats were treated routinely and after one hour of treatment animals in Group I were sacrificed while in rest of the groups, ulcer was induced by water immersion restraint method. Then animals were sacrificed. The animals in Group VI, VII and VIII were also subjected to gastric ulceration in the same manner as mentioned above. After one hour of ulcer induction animals were treated with standard and test drug and served as *Post*-treated standard group (VI) and Post-treated test group A and group B (VII and VIII) , respectively. All the animals were treated in this manner orally, once daily for five days. On 5th day, 12 hours fasted animals were treated routinely and after one hour of treatment, they were sacrificed. In all the above methods, the animals were sacrificed under The opent one anesthesia (40 mg/kg, IP). Stomach was removed from the body and opened along with the greater curvature, washed with fresh water and spread on cardboard with the mucous surface upwards. The mucosal surface was examined for ulceration with the help of magnifying lens (10 fold magnification) and scored by the method of Brzozowski et al [15,21,22]

VI. Assessment of Extent of Ulceration

The parameters viz. Ulcer score, ulcer index and reduction percentage in ulcer were taken to assess the

© 2014 Global Journals Inc. (US)

anti ulcerogenic effect. Histopathological studies were also carried out to determine the nature and amount of damage and the improvement after treatment.

VII. Statistical Analysis

The observations in various groups were expressed as Mean ± SEM. The ulcer score and index of various groups were compared with plain control group. The group comparison was analyzed by using ANOVA one way with Kruskall Wallis and Dunn's pair comparison test.

VIII. Results

Plain control (Group I), showed no pathological sign. In Group II (Negative control) where ulcer was induced by Indomethacin (20mg/kg) once daily for 5 days, the ulcer score was found to be 1.08±0.27. The ulcer scores in pre-treated standard and test groups where the animals were treated with Standard drug & test drug in low dose were found to be 1.16±0.30 &1.33±0.27 respectively when compared to negative control showed non- significant result. In pre-treated test Group B (Group V), the test drug was given orally in the dose of 248mg/kg ulcer score was found to be 0.66±0.27 with respect to negative control, showed non- significant decrease. In *Post* treated standard group (Group VI). Ulcer score was found to be 0.66±0.27 (insignificant) with respect to negative control. In Post treated test group A (VII) it was observed 1.08±0.27 (insignificant). In Post treated test group B (VIII) score was found to be 0.33±0.10 (insignificant). The ulcer index in Negative control Pre and Post treated standard, test group A and B were found to be 1.25, 1.63, 1.80, 0.44, 0.17, 1.67 and 0.22 respectively and percentage of ulcer reduction in Pre and *Post* treated standard, test group A and B were observed -7, -19, 39, 39, 0, and 47, respectively when calculated with Negative control. (Table no 1)

Table 1 : Effect of Hydro alcoholic extract of Post *Sumaq* on Indomethacin induced restraint gastric ulcer

Groups	Treatment	ADU(Mean± SEM)	%RU	Ulcer index	%Reduction
Group I Plain control	D W	0.08±0.08	17%	0.01	94
Group II Negative control	D W + IM 20mg/kg dissolved in CMC	1.08±0.27	100%	1.25	-----
Group III Pre-treated Stand	Omeprazole 20 mg/ kg + IM 20mg/kg Dissolved in CMC	1.16±0.30	100%	1.63	-7
Group IV Pre-treated test A	*Post Sumaq* 145mg/kg+ IM 20mg/kg dissolved in CMC	1.33±0.27	100%	1.80	-19
Group V Pre-treated test B	*Post Sumaq* 248mg/kg IM 20mg/kg dissolved in CMC	0.66±0.27	67%	0.44	39
Group VI *Post*-treated Stand	IM 20mg/kg. dissolved in CMC+ omeprazole 20 mg/kg	0.66±0.27	50%	0.17	39
GroupVII *Post*-treated test A	IM 20mg/kg. dissolved in CMC + *Sumaq*145 mg/kg	1.08±0.27	100%	1.67	0
Group VIII *Post*-treated B	IM 20mg/kg. dissolved in CMC + *Post Sumaq* 248 mg/kg	0.33±0.10	67%	0.22	47

(N=6 in each group. DW = Distilled water, IM = Indomethacin, CMC= Carboxy, methyl cellulose, %RU =Percentage of rats with ulceration, ADU = Average degree of ulceration)

Ulcer score in Negative control was found to be significantly increased ($p<0.01$) 1.16±0.21 when compared to plain control. The ulcer score in pre-treated standard, test group A and test group B, score was found to be 0.91±0.27, 0.66±0.16, and 0.83±0.21 respectively. No significant reduction was observed when compared to negative control. While in Post-treated Group VI, Group VII and Group VIII first ulcer was graded and ulcer score was found to be 0.41±0.08, 0.75±0.25 and 0.75±0.25 respectively, but no significant reduction was observed when compared to negative control. The ulcer index in Negative control, Pre and Post treated standard, test group A and B was observed 1.42, 0.76, 0.67, 0.83, 0.35, 0.76, and 0.83 respectively and percentage of ulcer reduction was found to be 45, 60, 50, 75, 55, and 55 respectively. (Table no 2).

Table 2 : Effects of hydro alcoholic extract of Post *Sumaq* on Water-immersion induced restraint gastric ulcer

Groups	Treatment	ADU(Mean± SEM)	%RU	Ulcer index	%Reduction
Group I Plain control	D.W	0.08±0.08	17%	0.01	93

© 2014 Global Journals Inc. (US)

Group II Negative control	D.W+ Ulcer induction	1.16±0.21*	100%	1.42	---
Group III Pre-treated Stand	Omeprazole 20 mg/ kg + Ulcer Induction	0.91±0.27	83%	0.76	45
Group IV Pre-treated test A	*Post Sumaq* 145 mg/kg+ Ulcer induction	0.66±0.16	100%	0.67	60
Group V Pre-treated test B	*Post Sumaq* 248mg/kg + Ulcer induction	0.83±0.21	100%	0.83	50
Group VI *Post*-treated Stand	Ulcer Induction + Omeprazole 20 mg/kg	0.41±0.08	83%	0.35	75
GroupVII *Post*-treated test A	Ulcer induction + *Post Sumaq* 145 mg/kg	0.75±0.25	83%	0.76	55
Group VIII *Post*-treated B	Ulcer induction. + *Post Sumaq* 248 mg/kg	0.75±0.25	83%	0.83	55


(N=6 in each group. Test used Kruskall Wallis test with Dunn, pair comparison test, N= 6 $p<0.05$ with respect to plain control, D.W= Distilled water)*

IX. Discussion


Gastric ulceration has long been viewed as the disease of stress, hence central nervous system may also play role in production of ulcer by causing hyperacidity. The techniques of restraint in albino rats provide a model for the study of stress induced gastric ulceration. Water immersion induced restraint gastric ulcer model is suitable to see anti stress effect of drugs. Therefore, the test drug was also evaluated by using this model. In Water Immersion Induced restraint gastric ulcer model the test drug was found both *precautionary and therapeutic in pre treated and Post treated test Groups at both dose level but the result was statistically non significant.* The histopathological findings are also in consonance. The findings indicate that the test drug does not possess anti anxiety properties and the same has not also been mentioned in Unani classics. However, it is clear from the result that the test drug has preventive & curative effect only at higher dose. Phytochemicals in *Rhus coraria* are ellagic acid, gallic acid isoquercitrin, myricitrin, myricetin, quercetin, quercitrin and tannic acid and flavinoids. Flavonoids protect the gastrointestinal mucosa from lesions produced by experimental ulcer models and different necrotic agents. Several mechanisms of action may be involved in this protective effect. Quercetin has an anti secretary mechanism of action. However, the most important mechanism of action responsible for the antiulcer activity of flavonoids is the antioxidant properties. Tannins are gastro protective which are present in the drug in sufficient amount[16,17].

As per the Unani theories it seems that the drug may have acted by temperament, as the *Mizaj* of the test drug is cold whereas that of diseases is hot[18,19,20]. But the anti ulcer mechanism cannot be just understood by *Mizaj* or Phytochemicals only as in case of herbal drugs only one or two or more Phytochemicals are not responsible for action. A number of chemicals and other interventions play the role in exerting actions and the ultimate effect is the cumulative effect. Further study is needed to establish the mechanism of anti ulcer effect of Post *Sumaq*.

Histopathological slides of different groups are shown below (Indomethacin ulcer model)




Group I : Normal mucosa inflammation and ulceration




Group II : Congestion, necrosis,


© 2014 Global Journals Inc. (US)




Group III : Oedema, necrosis, inflammation and ulceration




Group IV : Oedema, necrosis and ulceration




Group V : Inflammatory changes



Group VI : Inflammatory changes




Group VII : Oedema, necrosis, inflammation and ulceration




Group VIII : Inflammation and ulceration

Histopathological slides of different groups are shown below (Water immersion-induced restraint ulcer model)




Group I : Necrosis, inflammation and ulceration




Group II : Necrosis and inflammation


© 2014 Global Journals Inc. (US)




Group III : Necrosis and inflammation




Group IV : Necrosis and inflammation



Group V : Necrosis and inflammation



Group VI : Inflammatory changes



Group VII : Inflammatory changes

X. Conclusion

Results of different experimental models revealed Post *Sumaq* to be a promising antiulcerogenic drug. The test drug possesses curative effect at higher dose against Indomethacin induced gastric ulcer. In Water Immersion-induced restraint gastric ulcer model the effect was less prominent therefore it can be concluded the test drug does not possess anti anxiety effect as this model produces ulcer due to stress. This is also evident from the literature that Post *Sumaq* is not used as an anti anxiety. The preventive effect of the test drug was more pronounced. This also validates the claim that herbal drugs are more preventive in nature. The drug is more effective at higher dose; therefore the dose of Post *Sumaq* should be revised after toxicity studies.

XI. Acknowledgement

The authors are highly thankful to the authorities of NIUM for providing necessary facilities to carry out this research work.

Reference Références Referencias

1. Falcao HS, Leite JA, Barbosa-Filho JM, Athayde-Filho FP, and Chaves et al, Gastric duodenal antiulcer activity of alkaloids, Molecules., 2008; (13): 3198-3223.
2. Wilson and Ross, Anatomy and physiology in health and Illness,2003; (9): 323.
3. Calam J, Baron JH, Pathphysiology of duodenal and gastric ulcer and gastric cancer. *Brit Med. J,* 2001; (323):980-84.

© 2014 Global Journals Inc. (US)

4. Humes H, Kelley's essentials of internal medicine, 2nd ed. Lippincott William and Wilkins, 200l: 94-99.
5. Ledingham JG ,G Warrell D A, Concise Oxford Text book of Medicine, Oxford University Press, Inc New York, 2000: 530-33.
6. Bandyopadhyay SK, Satyesh C, Pakrashi A, The role of antioxidant activity of Phyllanthus emblica fruits on prevention from Indomethacin induced gastric ulcer, *Journal of Ethnopharmacology,* 2000: 171-176.
7. Raju. D, Ilango K, Chitra V, Ashish K, Evaluation of anti-ulcer activity of methanolic extract of Terminalia chebula fruits in experimental rats. *Journal Pharm. Sci.* & Res, 2009; (3): 101-110.
8. Dharmani P, Mishra PK, Maurya R, Chauhan VS and Palit G, Allophylus serratus; A plant with potential anti-ulcerogenic activity, *J. Ethno pharmacology,* 2005;(99): 361-366.
9. Dahnukar SA, Kulkarni RA, Rege NN, Pharmacology of Medicinal plants and natural products, *Indian Journal of Pharmacology,* 2000;(32): S81-S118.
10. Ghani N, Khazainul Advia; 3rd ed. Idarah Kitabul Shifa , New Delhi, 2011: 826.
11. Freirich EJ, Gehan EA, Rall DP, Schmidt LH, Skipper HE et al, Quantitative comprisom of toxicity of anti-cancer agents in mouse, rat, hamster, dog, monkey and man, *Cancer chemotherapy Report*,1966; 50 (4): 219-44.
12. Miller L C and Tainter M L, *Proc. Soc Expt. Biol, Med*, 1944: 57-261
13. Vogel HG, Drug discovery and evaluation pharmacological Assays, 2nd edi. Published by Springer-verlag Berlin Heidel berg, 2002: 869.
14. Hayaso M, Takeuchi K. Gastric acid secretion and lesion formation in rats under water immersion stress, *Dig Dis Sci* ,1986; (31): 166-71.
15. Brzozowski T, Konturek SJ, Kwiecien S, Pajdo R, Brzozowski I, Hahn EG et al, Involvement of endogenous cholecystokinin and somatostatin in gastro protection induced by intra duodenal fat, *J Clin Gastroenterol* , 1998; (27): 125-137.
16. Abu B, Ghaleb A, Dauod A, Safiya K, Advan AS, Antibacterial activity of Rhus coriaria extracts growing in Palestine, Journal *of the Islamic University of Gaza (Natural Science Series)*, 2005; 13(2): 147-153.
17. Duke JA, Bogenschutz GM, Ducellier J, Duke PAK, Handbook of Medical Plants, Boca Raton; CRC Press, 2003: 269-70.
18. Hubal I ,Kitabul Mukhtarat Fil Tibb (Urdu translation by CCRUM), Vol. 1st, 2nd, 3rd, New Delhi; Ministry of H and FW, Govt. of India, 2004: 129,187, 235.
19. Sina I, Al Qanoon (Urdu translation by Kantoori GH),Vol. 3rd. New Delhi; Idara Kitabush Shifa, New Delhi; 2007: 446-48.
20. Tabri R, Firdosul Hikmat (Urdu translation by Shah MA), Vol.1st and 2nd Deoband; Faisal Publications, 2000: 197-198.
21. Haqeeq A, Faiyaz A, Izharul H, & Shabbir A. Unani Description of Sumaq (Rhus Coriaria Linn.) *and its Scientific Report,* Global journal of medical research, 2013; 7 (13): 75-78
22. Haqeeq A, Wadud A, Nasreen J, Mudasir K, & Ghulamuddin S. *Evaluation of Anti-ulcer activity of hydro alcoholic extract of Post Sumaq (Rhus coriaria Linn.) in Ethanol induced Gastric ulcer in experimental Rats.* International Research Journal of Medical Sciences, 2013; 1(10): 7-12

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine
Volume 14 Issue 7 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Antidepressant and Anti-Inflammatory Activities of Cationic Amphiphilic Complexes of Sulphadiazine

By K.Hariprasath, I. Sudheer Babu, P.Venkatesh & U. Upendra Rao

Adarsa College of Pharmacy, India

Abstract- In our study we synthesized schiff' base of sulphadiazine on treating with aromatic aldehydes like para diethyl amino benzyldehyde and paradimethyl amino benzyldehyde. The synthesized schiff's bases were converted to its cationic amphiphilic bases by treating with methyl iodide. The cationic schiff bases were converted to metal complexes by treating with metals like copper chloride (CuCl2), zinc chloride (ZnCl2) and cadmium chloride (CdCl2). All the synthesized compounds were characterized by elemental analysis, IR and H1 NMR. Synthesized compounds were screened for anti-inflammatory and antidepressant activity. Copper metal complexes showed excellent anti-inflammatory activity and zinc metal complexes showed excellent anti-depressant activity.

Keywords: cationic amphiphilic bases, schiff' base, metal complexes, anti-inflammatory and antidepressant activity.

GJMR-B Classification : NLMC Code: QV 37.5



Strictly as per the compliance and regulations of:

© 2014. K.Hariprasath, I. Sudheer Babu, P.Venkatesh & U. Upendra Rao. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Antidepressant and Anti-Inflammatory Activities of Cationic Amphiphilic Complexes of Sulphadiazine

K.Hariprasath [α], I. Sudheer Babu [σ], P.Venkatesh [ρ] & U. Upendra Rao [ω]

Abstract- In our study we synthesized schiff' base of sulphadiazine on treating with aromatic aldehydes like para diethyl amino benzyldehyde and paradimethyl amino benzyldehyde. The synthesized schiff's bases were converted to its cationic amphiphilic bases by treating with methyl iodide. The cationic schiff bases were converted to metal complexes by treating with metals like copper chloride (CuCl2), zinc chloride (ZnCl2) and cadmium chloride (CdCl2). All the synthesized compounds were characterized by elemental analysis, IR and H1 NMR. Synthesized compounds were screened for anti-inflammatory and antidepressant activity. Copper metal complexes showed excellent anti-inflammatory activity and zinc metal complexes showed excellent anti-depressant activity.

Keywords: cationic amphiphilic bases, schiff' base, metal complexes, anti-inflammatory and antidepressant activity.

I. Introduction

An amphiphilic substance exhibits a double affinity, which can be defined from the physico-chemical point of view as a polar-apolar duality. When a single surfactant molecule exhibit both anionic and cationic dissociations it is called amphoteric or zwitterionic. Cationic amphiphilic drugs (CADs) are widely used in chronic pharmacotherapies in spite of frequently observed side effects connected with lysosomal phospholipid (PL) storage. Cationic amphiphilic drugs (CADs) represent compounds of different therapeutic classes such as antidepressants, neuroleptics, and antiarrhythmics. In acidic cellular compartments these drugs become efficiently protonated and thus trapped in, e.g. lysosomes. As a result of pHdependent ion trapping, total lysosomal drug concentrations may exceed extracellular levels by orders of magnitude. Lysosomotropic drugs may inhibit lysosomal phospholipid (PL) metabolism leading to the formation of dense cytoplasmic granules, i.e. lysosomes filled with undegraded PLs. The formation of drug-PL complexes further enhances intracellular accumulation of drugs.

Author α ω : Adarsa College Of Pharmacy, Kothapalli, Rajamundary, A.P, India. e-mail: hariprasath79@gmail.com

Author σ ρ : Sir C R Reddy College Of Pharmaceutical Sciences, Eluru, A.P, India.

We all require iron, copper and zinc for normal brain function but metal metabolism becomes dysregulated in a variety of neurodegenerative diseases. Metals accumulate in Alzheimer's dementia and Parkinson's disease and are deficient in Menkes disease. Transition metals perform a wide range of biological functions in the brain. A common feature is their ability to exist in a variety of oxidation states and participate in redox reactions; thus copper, iron, and manganese are all catalytically active metals in a class of enzymes that sequester free radicals. It is useful to look at the common and varying functions of transition metals in the brain to better understand what mechanisms are disrupted in metal dyshomeostasis and how this may lead to cell death in diseases of the CNS (Tyszka, 2014).

Metal complexes are also known as coordination compounds, which include all metal compounds. Metal complex is a structure consisting of a central atom (or) ion (metal) bonded with anions (ligands). Compounds that contain a coordination complex are called coordination compounds. The bonding characteristics of complexes and alteration in size of the metal ion are related to thermodynamic aspects. The stability constants for the complexes formed from various metal ions and one ligand have a particular sequence (Banerjee, 2009, Shi, 2007).

The parent sulpha drugs Sulphadiazine is a well known antibacterial in olden days. But owing to their narrow spectrum of activity and side effects, now a days their usage is limited. In our research we improved the biological activity by different synthetic modifications, In the first step by converting the parent sulpha drug in the form of Schiff base, there by generating a lone pair of electrons in a sp2 hybridized orbital in the structure leading to the derivation of and different biological properties. In the second step of synthesis the Schiff bases were converted in to their cationic amphiphiles, which may alter their pharmacokinetic profile like distribution and binding parameters. This step helped us to improve the spectrum activity from narrow spectrum to broad spectrum activity, increasing the permeability of drug molecule in brain to cross blood-brain barrier, which improves the antidepressant property, and release of cationic lipids into the macrophage cytoplasm is a necessary step for anti-inflammatory activity. In the


© 2014 Global Journals Inc. (US)

third step by deriving metal complexes of copper and zinc the biological properties like antidepressants and anti-inflammatory activities were strongly elucidated.

II. Materials and Methods

a) Synthesis of Schiff's Base

The Schiff's base has been synthesized by refluxing the reaction mixture of hot ethanolic solution (30 ml) of Sulphadiazine (0.01 mole) with hot ethanolic solution (30 ml) of different aromatic aldehyde (0.01 mole) for about 2-3 hours at 60-70^0 C (Fig-1). The mixture was allowed to stand over night. After that the colored solid product was filtered off, re-crystallized with ethanol and finally washed with petroleum ether. The final product was dried under reduced pressure over anhydrous calcium chloride (Panneerselvam, 2005).




Compound A R-C_2H_5, R_1-C_2H_5
Compound B R-CH_3, R_1-CH_3

Figure 1: Scheme of synthesis of Sulphadiazine Schiff base

b) Synthesis of cationic derivative of Schiff base

Cationic derivatives of Schiff bases were obtained by direct reaction between equimolar amount of the synthesized Schiff bases and methyl iodide in 50ml ethanol (Fig-2). The reaction mixture was refluxed for 8 hours and left overnight. The precipitated products were filtered and recrystallized (Negm, 2010)




Compound A R-C_2H_5, R_1-C_2H_5
Compound B R-CH_3, R_1-CH_3

Figure 2 : Scheme of synthesis of cationic Schiff base

© 2014 Global Journals Inc. (US)

c) Synthesis of transition metal complexes.

Metal ion solutions of anhydrous $CuCl_2$, $ZnCl_2$ and $CdCl_2$ (0.0005 mol) in 50ml ethanol was added with synthesized cationic derivative of Schiff base separately and refluxed for 6 hours (Fig-3). The reaction mixture was left overnight to complete the precipitation of the products. The products were recrystallized with ethanol to obtain pure products (Ibotomba, 2012, Ajaykumar, 2009).



Compound A R-C_2H_5, R_1-C_2H_5
Compound B R-CH_3, R_1-CH_3
MCl_2- $CuCl_2$, $Zncl_2$ and $CdCl_2$

Figure 3 : Scheme of synthesis of cationic Schiff base

d) Determination of median lethal doses (LD50)

Animal's Swiss albino mice (20-25gm) and Male Sprague - Dawley rats (160-180) were maintained at standard diet and *ad libido.* The experiment protocol was approved from institutional ethical committee. The test compounds were dissolved in 3 % DMSO administered orally to different groups with increasing doses. Six animals were taken in each group. Mortality was determined after 24 hours of treatment. The dose, at which the 50 % mice survived, was considered as LD50 value of the compound (OECD, 2002).

Male Sprague - Dawley rats weighing 160-180 grams were divided into eight groups of six animals each. The test groups received orally 20 mg/kg of each sample. The reference group received imipramine (5 mg/kg, p.o) while the control group received vehicle (1 % CMC). Naïve rats are individually forced to swim inside a vertical Plexiglas cylinder (height : 40 cm ; diameter : 18 cm ; containing 15 cm of water maintained at 25 oC). Floating behaviour during this 5 minutes period has been determined in different groups of rats. The percentage inhibition was calculated by the formula (Kulkarni, 2010).

e) Anti depressant activity

$$\text{Percentage inhibition} = \frac{\text{Before treatment} - \text{After treatment}}{\text{Before treatment}} \times 100$$

f) Anti inflammatory activity

Swiss albino mice were divided into eight groups of six animals each. The test groups received orally 20 mg/kg of each sample. The reference group received diclofenac sodium (10 mg/kg, p.o) while the control group received vehicle (1 % CMC). All the animals should make a mark on both hind paws just beyond tibiofasial junction, so that every time the paw is dipped in mercury column up to fixed mark to ensure constant paw volume. After drug administration inject 0.1ml white egg portion to the plantar region of left paw of control as well as treated group. The right paw serve as reference non inflamed paw for comparison. The inflammation was quantitated in terms of ml i.e.

© 2014 Global Journals Inc. (US)

replacement of water by edema using a Plethysmometer immediately before egg white injection and then 0, 1, 2 and 3 hours after egg white injection. The percent inhibition of edema as calculated for each group with respect to its vehicle treated control group. The anti-inflammatory activity was calculated by using the relation used by

$$\% \text{ inhibition} = (Vc\text{-}Vt/Vc) \times 100$$

Whereas Vc was the average inflammation (hind paw edema) of the control group of mice at a given time, Vt was the average inflammation of the drug treated (i,e sample or reference diclofenac sodium) mice at the same time (Sathe, 2011).

g) Statistical analysis

The data were expressed as mean ±SEM. Statistical analysis was performed one-way ANOVA followed by Dunnett‟s multiple comparison test using sigma stat software (version 2.0, Jandel Scientific Inc. USA

III. Results and Discussion

a) Characterization of synthesized compounds

A1- Copper metal complex of (E)- N-(4-(diethyl, methyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine.

M.F: $C_{22}H_{26}Cl_2CuI_2N_5O_2S$. M. wt: 812.8. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H^1 NMR ($CDCl_3$) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group, two triplet at 1.13 for two CH_3 groups in N-ethyl substitution, two quadret at 3.39 for two CH_2 group in N-ethyl substitution and singlet at 2.85 for N-methyl substitution. Elem Anal Calc: C, 32.51; H, 3.22; Cl, 8.72; Cu, 7.82; I, 31.2; N, 8.62; O, 3.94; S, 3.95. Elem Anal Found: C, 32.41; H, 3.32; Cl, 8.62; Cu, 7.92; I, 31.21; N, 8.60; O, 3.95; S, 3.96.

A2- Zinc metal complex of (E)- N-(4-(diethyl, methyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine.

M.F: $C_{22}H_{26}Cl_2CuI_2N_5O_2SZn$. M. wt: 814.6. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H^1 NMR (CDCl3) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group, two triplet at 1.13 for two CH3 groups in N-ethyl substitution, two quadret at 3.39 for two CH2 group in N-ethyl substitution and singlet at 2.85 for N-methyl substitution. Elem Anal Calc: C, 32.44; H, 3.22; Cl, 8.70; I, 31.16; N, 8.60; O, 3.93; S, 3.94; Zn, 8.03. Elem Anal Found: C, 32.45; H, 3.23; Cl, 8.66; I, 31.14; N, 8.62; O, 3.83; S, 3.98; Zn, 8.13.

A3- Cadmium metal complex of (E)- N-(4-(diethyl, methyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine.

M.F: $C_{22}H_{26}CdCl_2I_2N_5O_2S$. M.wt: 861.7. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H^1 NMR (CDCl3) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group, two triplet at 1.13 for two CH_3 groups in N-ethyl substitution, two quadret at 3.39 for two CH_2 group in N-ethyl substitution and singlet at 2.85 for N-methyl substitution. Elem Anal Calc: C, 30.67; H, 3.04; Cd, 13.05; Cl, 8.23; I, 29.46; N, 8.13; O, 3.71; S, 3.72. Elem Anal Found: C, 30.77; H, 3.14; Cd, 13.00; Cl, 8.13; I, 29.36; N, 8.03; O, 3.78; S, 3.74.

B1- Copper metal complex of (E)- N-(4-(trimethyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine

M.F: C20H22Cl2CuI2N5O2S. M.wt: 784.7. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H1 NMR (CDCl3) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group and three singlet at 2.85 for three N-methyl groups. Elem Anal Calc: C, 30.61; H, 2.83; Cl, 9.04; Cu, 8.10; I, 32.34; N, 8.92; O, 4.08; S, 4.09. Elem Anal Found: C, 30.59; H, 2.81; Cl, 9.14; Cu, 8.00; I, 32.14; N, 8.82; O, 4.18; S, 4.00.

B2- Zinc metal complex of (E)- N-(4-(trimethyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine

M.F: $C_{20}H_{22}Cl_2I_2N_5O_2SZn$. M.wt: 786.6. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H^1 NMR (CDCl3) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group and three singlet at 2.85 for three N-methyl groups. Elem Anal Calc: C, 30.54; H, 2.82; Cl, 9.01; I, 32.27; N, 8.90; O, 4.07; S, 4.08; Zn, 8.31. Elem Anal Found: C, 30.44; H, 2.92; Cl, 9.00; I, 32.26; N, 8.95; O, 4.17; S, 4.13; Zn, 8.21

B3- Cadmium metal complex of (E)- N-(4-(trimethyl amino) benzylidene)-4-(pyrimidin -2-sulfonamidyl) benzenamine.

M.F: $C_{20}H_{22}CdCl_2I_2N_5O_2S$. M.wt: 833.6. IR (KBr) cm-1: NH bond stretching at 3400 cm-1, C=N bond stretching at 1690 cm-1, S=O stretching at 1140 cm-1, C=C stretching at 1600 and 1475 cm-1. H^1 NMR ($CDCl_3$) δ values: Multiplet at 7.44-8.38 for aromatic nucleus, singlet at 8.39 for N=CH peak, singlet at 4.0 for aromatic NH group and three singlet at 2.85 for three N-methyl groups. Elem Anal Calc: C, 28.82; H, 2.66; Cd, 13.48; Cl, 8.51; I, 30.45; N, 8.40; O, 3.84; S, 3.85. Elem Anal Found: C, 28.92; H, 2.56; Cd, 13.42; Cl, 8.57; I, 30.40; N, 8.45; O, 3.82; S, 3.87

© 2014 Global Journals Inc. (US)

b) Anti inflammatory activity of synthesized compounds

Anti inflammatory activity was carried by paw oedema method using diclofenac sodium as standard. The results are given in the table-1. Currently used anti-inflammatory drugs are associated with some severe side effects. Therefore, the development of potent anti-inflammatory drugs with fewer side effects is necessary. A major factor limiting their use is gastrointestinal toxicity. In recent years, Schiff bases are widely used in formulating various types of drugs for their diversive biological activities. Metal complexes of Schiff bases have also been used as anti-inflammatory and antiarthritic agents. Anti- inflammatory of Zn (II) and Cu(II) complexes of indomethacin has been reported previously (Venugopala, 2003, Wei, 2006, Alam, 2012 & Sondhi, 2006), based on this we have evaluated anti-inflammatory activity of synthesized metal complexes. The down regulation of pro-inflammatory mediators through interaction of cationic lipids with the PKC pathway may explain this anti-inflammatory activity. Furthermore, since cationic lipids have intrinsic anti-inflammatory activity. Studies indicating that the release of cationic lipids into the macrophage cytoplasm is a necessary step for anti-inflammatory activity (Mario, 1997). Results revealed that the copper metal complexes (20mg/kg.b.wt) of Schiff bases of sulpha drugs A1, B1 showed excellent anti-inflammatory activity in carrageenan induced edema method by comparing with standard drug diclofenac sodium (10mg/kg.b.wt). It was thus confirmed that copper complexes, a unique class of potentially more therapeutically useful anti-inflammatory drugs. These results demonstrate that cationic lipids can be considered as novel anti-inflammatory agents.

Table1 : Results of *in vivo* anti inflammatory activity of metal complexes of schiff's base of Sulphadiazine

	Group	Dose	Paw volume (ml)				Difference Vc-Vt/Vc	Mean value	% anti-inflammatory activity
			0 hour	1 hour	2 hour	3 hour			
Group-I	Control	1% CMC	0.2	0.6	0.5	0.5	0.4	0.33	-
			0.2	0.6	0.6	0.7	0.4		
			0.2	0.5	0.6	0.6	0.3		
			0.2	0.5	0.6	0.7	0.3		
			0.2	0.5	0.5	0.8	0.3		
			0.2	0.5	0.7	0.8	0.3		
Group-II	Compound 1A1	(20mg/kg b.wt)	0.3	0.3	0.1	0.1	0	0.06	81***
			0.3	0.4	0.2	0.2	0.1		
			0.3	0.3	0.2	0.1	0		
			0.3	0.4	0.2	0.1	0.1		
			0.3	0.4	0.2	0.2	0.1		
			0.2	0.3	0.2	0.2	0.1		
Group-III	Compound 1A2	(20mg/kg b.wt)	0.3	0.5	0.4	0.3	0.2	0.20	39*
			0.3	0.6	0.4	0.3	0.3		
			0.3	0.5	0.4	0.3	0.2		
			0.2	0.4	0.2	0.2	0.2		
			0.3	0.4	0.3	0.2	0.1		
			0.3	0.5	0.4	0.4	0.2		
Group IV	Compound 1A3	(20mg/kg b.wt)	0.3	0.5	0.5	0.4	0.2	0.23	30*
			0.3	0.6	0.4	0.4	0.3		
			0.2	0.4	0.3	0.3	0.2		
			0.3	0.5	0.4	0.4	0.2		
			0.3	0.6	0.5	0.3	0.3		
			0.3	0.5	0.5	0.4	0.2		
Group V	Compound 1B1	(20mg/kg b.wt)	0.3	0.4	0.2	0.1	0.1	0.05	84***
			0.3	0.3	0.2	0.1	0		
			0.3	0.3	0.2	0.2	0		
			0.3	0.4	0.1	0.1	0.1		
			0.3	0.4	0.2	0.1	0.1		
			0.3	0.3	0.2	0.1	0		

© 2014 Global Journals Inc. (US)

Group VI	Compound 1B2	(20mg/kg b.wt)	0.3	0.5	0.4	0.4	0.2	0.18	45**
			0.3	0.6	0.5	0.2	0.3		
			0.3	0.5	0.4	0.3	0.2		
			0.3	0.4	0.2	0.1	0.1		
			0.3	0.4	0.3	0.2	0.1		
			0.3	0.5	0.4	0.4	0.2		
Group-VII	Compound 1B3	(20mg/kg b.wt)	0.3	0.6	0.4	0.4	0.3	0.26	21*
			0.3	0.7	0.5	0.5	0.4		
			0.3	0.5	0.5	0.1	0.2		
			0.3	0.4	0.7	0.2	0.1		
			0.3	0.7	0.8	0.7	0.4		
			0.3	0.5	0.7	0.4	0.2		
Group VIII	Standard Diclofenac sodium	10mg/kg	0.2	0.2	0.2	0.1	0	0.05	84***
			0.2	0.3	0.1	0.1	0.1		
			0.2	0.3	0.2	0.1	0.1		
			0.3	0.4	0.1	0.2	0.1		
			0.2	0.3	0.1	0.1	0.1		
			0.2	0.3	0.1	0.16	0.1		

n = 6. Values are expressed as ± S.E.M.

***P< 0.001, **P< 0.01, *P< 0.05, ns P > 0.05 Vs Control (One way ANOVA followed by Dunnett's test).

c) Antidepressant activity of synthesized compounds

Anti depressant activity was evaluated by force swim test method, the animals which are immobile for less time considered as active. The results are given the table-2. Results revealed that the zinc metal complexes (20mg/k.b.wt) of Schiff bases of sulpha drug A2, B2 showed excellent anti-depressant activity in reducing the duration of depressed behavior in animal models by despair swim test when compared with the standard drug Imipramine (5mg/kg.b.wt).

Table 2 : Results of *in vivo* anti depressant activity of metal complexes of schiff's base of Sulphadiazine

S.No	Treatment	Immobile response in 5 minutes		Percentage response (%)
		Before treatment	After treatment	
1	Group-I Control (1% CMC)	3.20	3.66	-
		2.86	3.14	
		3.12	3.00	
		3.56	3.66	
		3.88	3.22	
		2.88	2.86	
2	Group-II Standard (Imipramine 5mg/kg.b.wt)	3.28	1.48	65***
		3.86	1.76	
		3.42	1.34	
		3.66	0.68	
		3.18	0.86	
		3.88	1.26	
3	Group-III A1 (20mg/kg.b.wt)	3.88	1.88	51**
		4.66	1.86	
		4.12	1.96	
		2.44	1.44	
		3.72	1.63	
		3.34	1.86	
4	Group-IV A2 (20mg/kg.b.wt)	4.20	1.66	75***
		3.86	0.88	
		4.12	1.22	
		3.44	0.86	
		4.66	0.66	
		2.88	0.46	
5	Group-V A3 (20mg/kg.b.wt)	3.88	1.88	37**
		2.88	1.42	
		3.56	2.88	
		4.66	3.68	

© 2014 Global Journals Inc. (US)

		3.98	2.24	
		4.00	2.62	
6	Group-VI B1 (20mg/kg.b.wt)	4.20	2.61	34*
		2.86	2.02	
		3.72	1.68	
		3.48	2.82	
		3.88	2.86	
		4.88	2.92	
7	Group-VII B2 (20mg/kg.b.wt)	3.20	1.42	67***
		2.86	1.04	
		4.44	1.22	
		4.33	1.06	
		3.86	1.48	
		2.56	0.66	
8	Group-VIII B3 (20mg/kg.b.wt)	4.20	2.86	29*
		3.86	1.98	
		3.62	2.80	
		3.56	2.98	
		3.86	2.62	
		3.58	2.68	

n = 6. Values are expressed as ± S.E.M. ***P< 0.001, **P< 0.01, *P< 0.05, ns P > 0.05 Vs Control (One way ANOVA followed by Dunnett's test)

The most probable causes for depression are connected with the loss of homeostasis of the stress hormones, neurotransmitters, and disturbed trace elements levels. It has been reported that successful depression therapy can lead to zinc level normalization. It is reported that early life stress is a major risk factor for development of later depression due to affected neurogenesis in brain, especially in hippocampus. On the molecular level, these processes may be zinc-dependent via antioxidative activity changes and its influence on proper course of brain development process (Malgorzata, 2014, Chandramouli, 2012). There is an evidence for the role of mitochondrial dysfunction in the pathophysiology and treatment of neurodegenerative diseases, including mood disorders (Kato, 2000, Stork, 2005). Respiratory rate is a parameter characterizing functioning of the oxidative phosphorylation. The mitochondrial hypothesis states that impaired energy metabolism of brain cells is involved in the pathophysiology of antidepressants. The therapeutic or side effects of drugs administered in the treatment of depression may involve the targeted regulation of mitochondrial functions. In previous study it is reported that there is a direct mitochondrial targeting is involved in mechanisms of action of pharmacologically different antidepressants. Antidepressants are potent partial inhibitors of mitochondrial respiration (Jana, 2012).

IV. Conclusion

Cationic amphiphilic drugs (CADs) represent compounds of different therapeutic classes such as antidepressants, neuroleptics, and antiarrhythmics. In their neutral, lipophilic form cationic amphiphilic drug enter cells and their organelles. In acidic cellular compartments these drugs become efficiently protonated and thus trapped in, e.g. lysosomes. By increasing the permeability of drug molecule in brain to cross blood-brain barrier, which improves the antidepressant property and release of cationic lipids into the macrophage cytoplasm is a necessary step for anti-inflammatory activity.

V. Acknowledgement

The authors are thankful to the department of Biotechnology, Acharya Nagarjuna University, Guntur, A.P for providing necessary facilities to carryout this work.

Reference Références Referencias

1. Tyszka-Czochara M, Grzywacz A, Gdula-argasi G, Librowski S, Wili Ski B and Opoka W (2014). The Role of Zinc in the Pathogenesis and Treatment of Central Nervous System diseases. Implications of zinc Homeostasis for proper CNS Function. Acta Poloniae Pharmaceutica ñ Drug Research, 71: 369-377.
2. Banerjee M, Azam A and Sahu SK (2009). Synthesis, Characterization and Biological Evaluation of Schiff'sbase Transition Metal Complexes with Celecoxib and Sulfomethoxazole, Journal of Pharmacy Research, 2: 1155-1158.
3. Shi L, Ge HM, Tan SH, Li HQ, Song YC, Zhu HL (2007). Synthesis and antimicrobial activities of Schiff bases derived from 5-chloro-salicylaldehyde, Eur J Med Chem, 42: 558–564.
4. Panneerselvam P, Nair RR, Vijayalakshmi G, Subramanian EH, Sridhar SK (2005) . Synthesis of

© 2014 Global Journals Inc. (US)

Schiff bases of 4-(4-aminophenyl)-morpholine as potential antimicrobial agents., Eur J Med Chem, 40: 225–229.

5. Negm NA, Zaki MF, and Salem MA (2010). Cationic Schiff base amphiphiles and their metal complexes: Surface and biocidal activities against bacteria and fungi. Colloids and Surfaces B: Biointerfaces 77: 96-103.
6. Ibotomba Singh U, Bhubon Singha RK , Radhapiyari Devib W and Brajakisor Singh Ch (2012). Schiff base complexes of Copper (II) ions : Synthesis, Characterization and Antimicrobial studies Journal of Chemical and Pharmaceutical Research, 4: 1130-1135.
7. Ajaykumar D, Kulkarni L, Sangamesh A, Patil L, Prema S, Badami S. (2009) Electrochemical Properties of some Transition Metal Complexes: Synthesis, Characterization and Invitroantimicrobial studies of Co(II), Ni(II), Cu(II), Mn(II) and Fe(III) Complexes. Int. J. Electrochem. Sci, 4: 858-862.
8. OECD (2002). Acute oral toxicity – Acute oral toxicity class method guidelines 423 adopted 23.03.996. In: Eleventh Addendum to the OECD guidelines for the testing of chemicals, Organization for Economic co-operation and development, Paris, June 2000.
9. Kulkarni SK. Handbook of experimental pharmacology. 3rd ed. Delhi: Vallabh prakashan; 2010: 131 – 134.
10. Sathe BS, Jaychandran E, Jagtap VA, and Sreenivasa GM (2011). Synthesis characterization and anti-inflammatory evaluation of new fluorobenzothiazole schiff's bases, International Journal of Pharmaceutical Research and Development, 3: 164–169.
11. Wei D, Li N, Lu G, and Yao K (2006). "Synthesis, catalytic and biological activity of novel dinuclear copper complex with Schiff base," Science in China B, 49: 225–229.
12. Venugopala KN and Jayashree BS (2003). "Synthesis of carboxamides of 2′-amino-4′-(6-bromo-3-coumarinyl) thiazole as analgesic and antiinflammatory agents," Indian Journal of Heterocyclic Chemistry, 12, 4, 307–310.
13. Alam MS, Choi J, and Lee D (2012). "Synthesis of novel Schiff base analogues of 4-amino-1, 5-dimethyl-2- phenylpyrazol-3-one and their evaluation for antioxidant and anti-inflammatory activity," Bioorganic & Medicinal Chemistry 20: 4103–4108.
14. Sondhi SM, Singh N, Kumar A, Lozach O, and Meijer L (2006). "Synthesis, anti-inflammatory, analgesic and kinase (CDK-1, CDK-5 and GSK-3) inhibition activity evaluation of benzimidazole/benzoxazole derivatives and some Schiff's bases," Bioorganic and Medicinal Chemistry, 14: 3758–3765.
15. Mario C. Filion & Nigel C. Phillips. Anti-inflammatory activity of cationic lipids (1997). British Journal of Pharmacology 122: 551 – 557.
16. Malgorzata Tyszka-Czochara, Agata Grzywacz, Goanna Gdula-argasi SkaTadeusz Librowski1, Bogdan wili Ski and Wlodzimierz Opoka (2014). The Role of Zinc in the Pathogenesis and Treatment of Central Nervous System (CNS) diseases. Implications of zinc Homeostasis for proper CNS Function. Acta Poloniae Pharmaceutica ñ Drug Research, 71: 369-377.
17. Chandramouli C, Shivanand MR, Nayanbhai TB, Bheemachari B, and Udupi RH (2012). "Synthesis and biological screening of certain new triazole schiff bases and their derivatives bearing substituted benzothiazole moiety," Journal of Chemical and Pharmaceutical Research 4: 1151–1159.
18. Kato T., Kato N (2000). Mitochondrial dysfunction in bipolar disorder. Bipolar Disorders 2: 180–190.
19. Stork, C., Renshaw, P.F (2005). Mitochondrial dysfunction in bipolar disorder: evidence from magnetic resonance spectroscopy research. Molecular Psychiatry 10: 900–911.
20. Jana Hroudova, Zdenek Fisar (2012). In vitro inhibition of mitochondrial respiratory rate by antidepressants. Toxicology Letters 213: 345– 352.

© 2014 Global Journals Inc. (US)

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine
Volume 14 Issue 7 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

In Vivo Anti-Inflammatory and in Vitro Antioxidant Activities of Toona Ciliata Leaves Native to Bangladesh

By Hemayet Hossain, Proity Nayeeb Akbar, Shaikh Emdadur Rahman, Tanzir Ahmed Khan, Md. Mahfuzur Rahman & Ismet Ara Jahan

Khulna University, Bangladesh

Abstract- The study was performed to assess the anti-inflammatory, antioxidant activities and identify the polyphenols of *Toona ciliata* grown in Bangladesh. Anti-inflammatory activity was examined by the methods of carrageenan and histamine-induced paw edema. At the dose of 400 mg/kg, effective anti-inflammatory activity ($P<0.01$) was observed in rats for both the test models of carrageenan and histamine-induced paw edema, compared to indomethacin. In ABTS scavenging assay, IC50 value was found significant (5.50µg/ml) compared to ascorbic acid (12.01µg/ml). The maximum absorbance of reducing power was obtained 0.4939 at 250µg/ml relative to ascorbic acid (1.1115µg/ml). Total antioxidant capacity, total phenolic and flavonoid content were found to be 357.1 mg/g ascorbic acid, 239.2 mg/g gallic acid, and 98.36 mg/g quercetin equivalent, respectively.

Keywords: toona ciliata, free radical scavenging, anti-inflammatory, hplc, epicatechin, p-coumeric acid, rutin hydrate.

GJMR-B Classification : NLMC Code: QV 766



Strictly as per the compliance and regulations of:

© 2014. Hemayet Hossain, Proity Nayeeb Akbar, Shaikh Emdadur Rahman, Tanzir Ahmed Khan, Md. Mahfuzur Rahman & Ismet Ara Jahan. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

In Vivo Anti-Inflammatory and in Vitro Antioxidant Activities of Toona Ciliata Leaves Native to Bangladesh

Hemayet Hossain [α], Proity Nayeeb Akbar [σ], Shaikh Emdadur Rahman [ρ], Tanzir Ahmed Khan [ω], M d. Mahfuzur Rahman [¥] & Ismet Ara Jahan [§]

Abstract- The study was performed to assess the anti-inflammatory, antioxidant activities and identify the polyphenols of *Toona ciliata* grown in Bangladesh. Anti-inflammatory activity was examined by the methods of carrageenan and histamine-induced paw edema. At the dose of 400 mg/kg, effective anti-inflammatory activity ($P<0.01$) was observed in rats for both the test models of carrageenan and histamine-induced paw edema, compared to indomethacin. In ABTS scavenging assay, IC50 value was found significant (5.50µg/ml) compared to ascorbic acid (12.01µg/ml). The maximum absorbance of reducing power was obtained 0.4939 at 250µg/ml relative to ascorbic acid (1.1115µg/ml). Total antioxidant capacity, total phenolic and flavonoid content were found to be 357.1 mg/g ascorbic acid, 239.2 mg/g gallic acid, and 98.36 mg/g quercetin equivalent, respectively. During HPLC analysis, catechin and ellagic acid were determined in considerable amounts (825.95 and 416.70 mg/100g extract, respectively). The findings suggest that *Toona ciliata* could be a potential source of natural antioxidant.

Keywords: toona ciliata, free radical scavenging, anti-inflammatory, hplc, epicatechin, p-coumeric acid, rutin hydrate.

I. Introduction

Toona ciliata *(T. ciliata)*, also commonly known as the red cedar, toon or toona, Burma cedar, Indian cedar or Indian mahogany, is a forest tree in the mahogany family (Meliaceae). It grows widely in the regions of southern Asia and Australia. (1, 2) These are usually large plants that grow up to a height of 25 to 35 m and the leaves are alternate and pinnetely veined with assymetrical base and an acute apex. (3)

Studies on the transverse section of the bark of *T. ciliata* revealed the presence of periderm, cortex, sclerides, mednllary rays and phloem fiber. (4) The barks were also found to contain tetranortriterpenoids, including toonacilin and the leaves hold a considerable amount of aromatic compounds like coumarin, glycoside, tannins, flavonoids, phenolic compounds, triterpenoids and steroids. (5) In addition, three new norlimonoids, two new tirucallane-type triterpenoids, and a new pimaradiene-type diterpenoid, along with two known limonoids and eight known tirucallane-type triterpenoids, were isolated from the leaves and twigs of *T. ciliata.* (5, 6)

The plant *T. ciliata* possess many important biological properties that account for it's traditional uses in medicinal treatments, construction purpose, dye preparation, etc. (7) The flowers are used to produce dye, which are worn around Asia as color silk. *T. ciliata* barks are useful in chronic dysentery, ulcer, leprosy, fever, headache, blood complaints, etc. (8)

The plant has been reported to exhibit significant antibacterial, antifungal, anticancer, anti-ulcer, anti-tumor, analgesic, anti-microbial, gastro protective and cytotoxic activity. (9, 10, 11) The ethanol leaf extract of *T. ciliata* was studied for its inhibitive effects on protein non-enzymatic glycation. (12)

The aim of the present work is to determine the anti-inflammatory, antioxidant activities and identify the bioactive polyphenolic compounds by HPLC in the ethanol extract of Toona ciliata leaves grown in Bangladesh.

II. Materials and Methods

a) Plant material

Fresh leaves were collected in May 2013 from Khulna, Bangladesh. Leaves of *T. ciliata* were washed, dried in the shade to minimize loss of volatile constituents and reduced to powder with a grinder.

b) Extraction

Collected fresh leaves were separated from undesirable materials and washed with water before letting it stand under the sun for a week. The dried leaves were coarsely powdered with the help of a grinder (Capacitor start motor, Wuhu motor factory, China). About 400g of the powered material was taken in a clean, flat-bottomed glass container and soaked in 1000 ml of ethanol. The container along with its contents

Author α σ §: BCSIR Laboratories, Dhaka, Bangladesh Council of Scientific and Industrial Research, Dr. Qudrat-E-Khuda Road, Dhaka-1205, Bangladesh. e-mail: hemayethossain02@yahoo.com
Author ρ: Pharmacy Discipline, Life Science School, Khulna University, Khulna-9208, Bangladesh.
Author ω ¥: Institute of Food Science and Technology, Dhaka, Bangladesh Council of Scientific and Industrial Research, Dr. Qudrat-E-Khuda Road, Dhaka-1205, Bangladesh.

© 2014 Global Journals Inc. (US)

was sealed and left to stand for a period of 7 days with continuous stirring by an orbital shaker. The mixture was first filtered in a clean cotton plug to remove any plant debris, and then through Whatman filter paper no. 1 (Bibby RE200, Sterilin Ltd., UK). The filtrate was concentrated by rotary vacuum evaporator (R-210, Buchi, Switzerland) and dried. The sample rendered 51g of greenish gummy concentrate (12.75%) and was designated as the crude ethanol extract.

c) Chemicals

Gallic acid (GA), (+)-catechin hydrate (CH), vanillic acid (VA), caffeic acid (CA), (-)-epicatechin (EC), p-coumaric acid (PCA), rutin hydrate (RH), ellagic acid (EA), quercetin (QU), ascorbic acid, ABTS, folin-ciocalteu's phenol reagent, carrageenan and histamine were purchased from Sigma–Aldrich (St. Louis, MO, USA). Acetonitrile (HPLC), methanol (HPLC), acetic acid (HPLC), ethanol, trichloroacetic acid (TCA), phosphate buffer (pH 6.6), potassium ferricyanide [K3Fe(CN)6], ferric chloride (FeCl3), sodium phosphate, EDTA, ammonium molybdate and sodium carbonate were of analytical grade and purchased from Merck (Darmstadt, Germany).

d) Test animals & drugs

For the screening of in vivo anti-inflammatory activity, male rats of Wister strain weighing 175-205 g were used. The animals were housed under standard laboratory conditions maintained at 25 ± 1°C and under 12/12 h light/dark cycle and feed with Balanced Trusty Chunts and water *ad libitum*. All experimental protocols were in compliance with Bangladesh Council of Scientific and Industrial Research (BCSIR) ethics committee on Research in animals as well as internationally accepted principles for laboratory animal use and care.

The standard drug, Indomethacin was used for this study and purchased from Square Pharmaceuticals Ltd, Bangladesh.

III. Anti-Inflammatory Activity Test

a) Carrageenan-induced oedema

The activity of *T. ciliata* ethanol leaf extract was evaluated using the carrageenan induced hind paw edema model. (13) The rats were divided into four groups (five rats per group). Group I (control) was given 1% tween 80 in normal saline (10 ml/kg), while Group II (positive control) received 10 mg/kg body wt. of indomethacin orally. Group III and IV were injected with 200 and 400 mg/kg body wt. of *T. ciliata* orally, respectively. Acute inflammation was induced in all the four groups by sub plantar injection of 0.1 ml of its suspension of carrageenan with 1% tween 80 in normal saline in the right paw of the rats, 1 h after the oral administration of the tested materials. The paw volume was measured with a micrometer screw gause at 1-hour interval after the administration of the drug and the extract. The percentage inhibition of inflammatory effect of the extract was calculated using the following expression:

Percentage inhibition of inflammation = [(Vc-Vt)/Vc] × 100, where Vc is the average degree of inflammation by the control group and Vt is the average degree of inflammation by the test group.

b) Histamine-induced oedema

The activity of the *T. ciliata* extract was evaluated with histamine-induced paw edema model. (14) The paw oedema was generated by injecting 0.1% histamine solution sub-plantarly into the left hind paw of each mice at a dose of 0.1 ml. Twenty rats were divided into four groups of five animals each. Group I (control) was supplied with 1% tween 80 in normal saline (10 ml/kg). Group II (positive control) received 10 mg/kg body wt. of indomethacin orally. Group III and IV were given 200 and 400 mg/kg body wt. of *T. ciliata* orally, respectively. Acute inflammation was induced in all the four groups by sub plantar injection of 0.1 ml of histamine with 1% tween 80 in normal saline in the right hind paw of the rats, 1 h after the oral administration of the tested materials. The paw volume was measured with a micrometer screw gause at 1, 2, 3 and 4 h after the administration of the drug and the extract. The percentage inhibition of inflammatory effect of the extract was calculated using the same formula as for calculating the carrageenan-induced paw oedema.

IV. Antioxidant Activity Test

a) ABTS radical scavenging activity test

The method of decolourisation of free radical $ABTS^+$ was performed according to Fan et al. with some modifications. (15) ABTS radical cation was prepared by mixing 7 mM ABTS solution with 2.45 mM potassium persulfate. The mixture was allowed to stand for 12-16 h at room temperature in the dark until reaching a stable oxidative state. The ABTS solution was diluted with ethanol to an absorbance of 0.70 ± 0.02 with pH 7.4 phosphate buffered saline (PBS) solution at 734 nm, before use. The reaction mixture was allowed to stand at room temperature for 6 min and the absorbance at 734 nm was immediately recorded. The ABTS scavenging activity was calculated as follows:

$$\text{ABTS scavenging effect} = I\ (\%) = (A_o - A_s / A_o) \times 100$$

Where, A_o = Absorbance of control and A_s = Absorbance of sample

© 2014 Global Journals Inc. (US)

b) Reducing power assay

The reducing power of *T. ciliata* was studied using the method of Hemayet et al. and Dehpour et al. [16, 17] The extract at different concentrations was mixed with 1 ml ethanol, 2.5 ml phosphate buffer (0.2 M, pH 6.6), and 2.5 ml potassium ferricyanide $[K_3F_e(CN)_6]$ (1%). The sample solutions were next incubated at 50°C for 20 min and a 10% solution of trichloroacetic acid (2.5 ml) was added to them. They were then centrifuged at 3000 rpm for 10 min. The top layer of the mixture (2.5 ml) was mixed with 2.5 ml distilled water and 0.5 ml of 0.1% $FeCl_3$. The absorbance was measured at 700 nm with a spectrophotometer. All determinations were carried out in triplicate.

c) Total antioxidant capacity

The total antioxidant capacity was measured by the method of Prieto et al. [18] The ethanol extract was prepared in its respective solvent and mixed with 1 ml of the reagent solution (0.6M H_2SO_4, 28 mM sodium phosphate, 4 mM ammonium molybdate mixture). The tubes were incubated for 90 min at 95°C. The mixture was cooled to room temperature and the absorbance was read at 695 nm against a blank sample. Ascorbic acid equivalents were calculated using the standard graph for ascorbic acid. The experiment was conducted in triplicates and values were expressed as equivalents of ascorbic acid in mg per gram of extract.

D. Total phenolic content

Total phenolic content of the extract was determined using the modified Folin-Ciocaltu method. [19, 20] After reacting 0.5 ml of extract (1 mg/ml), 5 ml Folin-Ciocaltu reagent (1:10 v/v distilled water) and 4 ml (75 g/l) of sodium carbonate, the sample solutions were mixed and left to stand at 40°C for the next 30 min for color development. The absorbance was read at 765 nm. The total phenolic content was calculated and expressed as mg of gallic acid equivalent per gram using the equation obtained from the standard gallic acid calibration curve, y = 6.993x + 0.0379, R2 = 0.9995.

d) E. Total flavonoid content

The total flavonoid content was determined by reactions of the aluminium chloride colorimetric method with some modifications. [21, 22] The absorbance of the reaction mixture was measured at 430 nm with a double beam Analykjena UV/Visible spectrophotometer (Model 205, Jena, Germany). Quercetin was used for calibration of a standard curve ($y = 6.2548x + 0.0925$; $R^2 = 0.998$) and the results were expressed as mg of quercetin equivalent per gram dry weight of sample.

V. HPLC Detection of Polyphenolics

Detection of selected polyphenolic compounds in the extract was carried out by HPLC as described by Ismet et al. [23] The analysis was performed on a Dionex UltiMate 3000 system equipped with quaternary rapid separation pump (LPG-3400RS) and photodiode array detector (DAD-3000RS). Separation was done using Acclaim® C_{18} (5μm) Dionex column (4.6 x 250 mm) at 30 °C with a flow rate of 1 ml/min and an injection volume of 20 μl. The mobile phase consisted of acetonitrile (solvent A), acetic acid solution pH 3.0 (solvent B), and methanol (solvent C) with the gradient elution program of 10%A/80%B/10%C (0-9 min), 20%A/60%B/20%C (10-19 min) and 100%A (20-30 min) with post run equilibration of the system with 5%A/95%B (5 min). The UV detector was set to 280 nm for 18.0 min, changed to 320 nm for 6 min, and finally to 380 nm and held for the rest of the analysis period while the diode array detector was set at an acquisition range from 200 nm to 700 nm. For the preparation of calibration curve, a standard stock solution was prepared in methanol containing gallic acid (GA), vanillic acid (VA), (+)-catechin (CH), (-)-epicatechin (EC), *p*-coumaric acid (PCA), rutin (R), ellagic acid (EA) (20 μg/ml each), caffeic acid (CA) (8 μg/ml) and quercetin (QU) (6 μg/ml). A solution of *T. ciliata* leaf extract was prepared in ethanol at 5 mg/ml. All solutions were filtered through 0.20 μm nylon syringe filter (Sartorius, Germany) and degassed in an ultrasonic water bath (Hwashin, Korea) for 15 min. Data acquisition, peak integration, and calibrations were performed with Dionex Chromeleon software (Version 6.80 RS 10).

VI. Statistical Aanalysis

Data were presented as mean ± Standard deviation (S.D). Statistical analysis for animal experiment was carried out using one-way ANOVA followed by Dunnet's multiple comparisons using SPSS Data Editor for Windows, Version 11.5.0 (SPSS Inc., U.S.A.). The results obtained were compared with the control group. *P* values < 0.05 were considered to be statistically significant.

VII. Results

a) Carrageenan-induced paw edema

The anti-inflammatory effect of the *T. ciliata* using carrageenan induced oedema test is expressed in Table 1. The paw edema was highly reduced by indomethacin ($p<0.05$; $p<0.01$) between the first and forth hour (50.48% to 64.46% inhibition). A maximum edema paw volume of 1.66 ± 0.08 mm was observed in the control group, four hours after the carrageenan injection. Rats which received 400 mg/kg body weight of the extract were observed to significantly decrease ($p<0.05$; $p<0.01$) the carrageenan-induced oedema paw volume between the 1 to 4 hour time interval, in comparison to that of the standard drug, indomethacin, at a dose of 10 mg/kg body weight. The highest reduction in the paw volume by the 400 mg/kg body weight of the extract at 4 h was 55.42%, while that by indomethacin was 64.46%, respectively.

© 2014 Global Journals Inc. (US)

Table 1 : Effect of ethanol extract of *Toona ciliata* leaves and indomethacin on carrageenan-induced oedema paw volume in male wistar rats

Treatment Groups	Doses (mg/kg body weight)	Right hind paw volume (mm)			
		1 h	2 h	3 h	4 h
Control	2 ml/kg	1.03 ± 0.07	1.30 ± 0.09	1.50 ± 0.04	1.66 ± 0.08
Positive Control (Indomethacin)	10	0.51 ± 0.05* (50.48)	0.63 ± 0.06** (51.54)	0.70 ± 0.07* (53.33)	0.59 ± 0.03** (64.46)
Extract	200	0.95 ± 0.04* (7.77)	1.09 ± 0.05* (16.15)	1.16 ± 0.08** (22.67)	1.18 ± 0.09* (28.92)
Extract	400	0.59 ± 0.06** (42.72)	0.70 ± 0.03* (46.15)	0.76 ± 0.06** (49.33)	0.74 ± 0.04** (55.42)

Values in brackets denote percentage inhibition of the oedema paw volume.

Values are expressed as mean±SD; Values are calculated as compared to control using one way-ANOVA followed by Dunnet's Test; * indicates $P < 0.05$; ** indicates $P < 0.01$ vs. control; $n = 5$.

b) Histamine-induced paw edema

Table 2 gives information on the effect of *T. ciliata* extract on acute inflammation using histamine-induced paw edema test. A maximum edema paw volume of 1.59 ± 0.08 mm was observed in the control group at 4 h after histamine was injected. Rats that were pre-treated with 400 mg/kg body weight of the extract significantly compressed ($p<0.05$; $p<0.01$) the histamine-induced edema paw volume, in comparison to that by indomethacin. The percentage inhibition of the edema paw volume at 1, 2 and 3 h by the 400 mg/kg body weight of the extract was also found effective ($p<0.05$; $p<0.01$). The maximum reduction in the paw volume by the 400 mg/kg body weight of *T. ciliata* at 4 h was 56.60%, while that by the indomethacin declined to 65.41%, respectively.

Table 2 : Effect of ethanol extract of *Toona ciliata* leaves and indomethacin on histamine-induced oedema paw volume in male wistar rats

Treatment Groups	Doses (mg/kg body weight)	Right hind paw volume (mm)			
		1 h	2 h	3 h	4 h
Control	2 ml/kg	1.09 ± 0.06	1.29 ± 0.07	1.40 ± 0.05	1.59 ± 0.08
Positive Control (Indomethacin)	10	0.48 ± 0.08** (55.96)	0.56 ± 0.03* (56.59)	0.53 ± 0.08* (62.14)	0.55 ± 0.03** (65.41)
Extract	200	0.84 ± 0.05* (22.94)	0.87 ± 0.09* (32.56)	0.92 ± 0.07** (34.29)	0.99 ± 0.06* (37.73)
Extract	400	0.60 ± 0.04* (44.95)	0.67 ± 0.07** (48.06)	0.71 ± 0.05* (49.29)	0.69 ± 0.05** (56.60)

Values in brackets denote percentage inhibition of the oedema paw volume.

Values are expressed as mean±SD; Values are calculated as compared to control using one way-ANOVA followed by Dunnet's Test; * indicates $P < 0.05$; ** indicates $P < 0.01$ vs. control; $n = 5$.

c) ABTS radical scavenging activity

At minimum concentration (10 µg/ml), the highest activity obtained by the extract of *T. ciliata* was 98.22 ± 0.04 µg/ml (Table 3). The IC50 value of the extract was found to be 5.50 ± 0.16 µg/ml, which was similar to that of the ascorbic acid (12.01 ± 0.12 µg/ml).

Table 3 : ABTS radical scavenging activity of *T. ciliata* leaf extract with standard ascorbic acid

Concentration (µg/ml)	*T. ciliata* leaf extract	Ascorbic acid
10	98.22 ± 0.04	48.60 ± 0.17
20	98.60 ± 0.12	85.79 ± 0.25

© 2014 Global Journals Inc. (US)

40	98.73 ± 0.08	99.19 ± 0.21
60	98.87 ± 0.41	99.25 ± 0.29
80	99.24 ± 0.10	99.53 ± 0.24
100	99.31 ± 0.13	95.58 ± 0.18
250	99.34 ± 0.11	99.85 ± 0.27
IC_{50}	5.50 ± 0.13	12.01 ± 0.12

The values are expressed as mean ± standard deviation (n=3).

d) Reducing power assay

The reducing power assay was determined based on the relative maximum absorbance of the extract of *T. ciliata* and was observed to increase with an increase in concentration (Table 4). At 250 µg/ml, the maximum absorbance for the ethanolic leaf extract of *T. ciliata* was found to be 0.4939 ± 0.029, while the standard ascorbic acid showed an absorbance of 1.1115 ± 0.009.

Table 4 : Reducing power assay of *T. ciliata* leaf extract with standard ascorbic acid

Concentration (µg/ml)	*T. ciliata* leaf extract	Ascorbic acid
10	0.0190 ± 0.028	0.3801 ± 0.012
20	0.0695 ± 0.071	0.4577 ± 0.017
40	0.0819 ± 0.017	0.5398 ± 0.023
60	0.1056 ± 0.041	0.6345 ± 0.037
80	0.1699 ± 0.062	0.7125 ± 0.013
100	0.1986 ± 0.041	0.7811 ± 0.029
250	0.4939 ± 0.029	1.1115 ± 0.009

The values are expressed as mean ± standard deviation (n=3).

e) Total antioxidant capacity

The ethanol extract of *T. ciliata* possessed a high total antioxidant capacity (Table 5). The total antioxidant capacity of the extract was obtained in significant quantity relative to the standard ascorbic acid per gram of extract (357.10 ± 2.02).

Table 5 : Total antioxidant capacity of ethanolic leaf extract of *T. ciliata*

Extract	Total antioxidant capacity
	mg of ascorbic acid equivalent (AAE) per g of dry extract
T. ciliata leaf extract	357.1 ± 2.02

The values are expressed as mean ± standard deviation (n=3).

f) Total phenolic content

Table 6 demonstrates the total phenolic content in the ethanol leaf extract of *T. ciliata.* High phenolic content was determined in the extract (239.2 ± 2.53 mg/g of gallic acid equivalent).

Table 6 : Total phenolic and flavonoid content of ethanolic leaf extract of *T. ciliata*

Extract	Total phenolic content	Total flavonoid content
	mg of gallic acid equivalent (GAE) per g of dry extract	mg of quercetin equivalent (QE) per g of dry extract
T. ciliata leaf extract	239.2 ± 2.53	98.36 ± 1.07

The values are expressed as mean ± standard deviation (n=3).

g) Total flavonoid content

Table 6 demonstrates the total flavonoid content in the leaf extract of T. ciliata. A considerably large amount of flavonoid was observed in the extract (98.36 ± 1.07 mg/g of quercetin).

h) HPLC assay of T. ciliata

The contents of the phenolic compounds in the leaf extract of *T. ciliata* were analyzed by RP-HPLC (Table 7). Based on the comparison of the retention times with those of the standard peaks, seven phenolic

© 2014 Global Journals Inc. (US)

compounds: (+) catechin, vanillic acid, epicatechin, p-coumeric acid, rutin hydrate, ellagic acid and quercetin were identified, respectively (Figure 1). The most abundant phenolic compound obtained from the extract of *T. ciliata* was catechin (825.95 ± 5.39 mg/100 g dry extract) followed by ellagic acid (416.70 ± 3.58 mg/100 g dry extract). Next, there was epi-catechin, p-coumeric acid and rutin hydrate, which were also obtained in significant quantities, but in lower amounts than that of the first two (211.7 ± 2.36, 102.20 ± 1.87 and 77.57 ± 1.49 mg/100 g dry extract, respectively). Other polyphenolic compounds like vanillic acid and quercetin were also obtained in similar concentrations (34.05 ± 0.83 and 29.13 ± 0.65 mg/100 g dry extract).

Table 7 : Contents of polyphenolic compounds in the ethanolic leaf extract of *T. ciliata* (n=3)

Polyphenolic compound	Ethanol extract of *T. ciliata* leaf	
	Content (mg/100 g of dry extract)	% RSD
CH	825.95	5.39
VA	34.05	0.83
EC	211.7	2.36
PCA	102.2	1.87
RH	77.57	1.49
EA	416.7	3.58
QU	29.13	0.65




Figure 1 : HPLC chromatogram of ethanol leaf extract of *T. ciliata*. Peaks: 1, catechin hydrate; 2, vanillic acid; 3, epi-catechin; 4, *p*-coumeric acid; 5, rutin hydrate; 6, ellagic acid; 7, quercetin

VIII. Discussion

Carrageenan and histamine induced paw oedema were evaluated for their anti-inflammatory effect in *T. ciliata*. The carrageenan induced inflammatory response in rats is a biphasic response, which causes marked oedema formation that results from the rapid production of several inflammatory mediators such as histamine, serotonin, and bradykinins. The second step is the release of prostaglandins and nitric oxide with a peak at 3 h, which is produced by an inducible is of orm of cyclooxygenase (COX-2) and nitric oxide synthase

© 2014 Global Journals Inc. (US)

(iNOS). [24] The present investigation was carried out in an attempt to reduce the oedomatogenic response in rats evoked by carrageenan. Results show that pre-treated oral administration of the extract was effective in the reduction of the response. Thus, a relationship can be inferred between the anti-inflammatory properties of the extract and the inhibition of intracellular signalling pathways in inflammatory mediators.

On injection, histamine acts as an inflammation mediator. [25] The liquid spreads out inside the body of the rat like a wheal and increases the permeability of the host capillary venules in the skin. Substances that inhibit the activity of histamine receptors shrink that particular area where the wheal was formed. This could be because the anti-inflammatory activity of the extract is supported by its anti-histamine activity. The antihistaminic effect of the extract increases with the concentration of the extract. The extract inhibits the formation and action of the inflammatory mediators, effectively suppressing the production of oedema by histamine. This study shows that *T. ciliata* has significant anti-oedematogenic effect ($P<0.01$) on paw oedema in rats induced by both carrageenan and histamine.

When subjected to reducing power assay, the extract causes the oxidation of ferricyanide complex to its ferrous form. This results in the extract to donate a hydrogen atom, which in turn helps to break the free radical chain and exert an antioxidant response. [26] The high phenolic content in the ethanol leaf extract of *T. ciliata* might be a reason for this reduction of Fe3+ to Fe2+, exhibiting stronger reducing power ability.

The total antioxidant capacity depends on the reduction of Mo (VI) to Mo (V) by the extract and the subsequent formation of green phosphate/Mo (V) complex at an acidic pH. [27])

Studies on the antioxidant activities of the leaf and flower extracts of *Toona ciliata* have previously been carried out using several different solvents i.e. petroleum ether, chloroform, ethyl acetate and methanol. Based on the results, all the extracts showed significant DPPH and ABTS radical scavenging activity in comparison with the standard, BHT (butylatedhydroxytoluene). The petroleum ether, chloroform, ethyl acetate and methanol extracts of *Toona ciliata* showed DPPH and ABTS significant activity with IC50 value of 150, 135.5, 105 and 92.5 µgml-1 for DPPH and 145, 120, 120.5 and 95 µgml-1 for ABTS scavenging activity, respectively, in comparison to standard BHT with an IC50 of 8 µgml-1 and 11.5 µgml-1, respectively. (12) Better results for the ABTS radical scavenging activity of *T. ciliata* ethanol extract relative to ascorbic acid were obtained from the present investigation (IC50 5.50 ± 0.130). Another study on the methanol and hexane fractions of *T. ciliata* leaves compared the plant's reducing power activity, which suggests a better activity for the methanol extract with a maximum relative absorbance of 0.440 ± 0.0001, which was very close to the absorbance of the standard drug, ascorbic acid (0.461 ± 0.0030). [28]

HPLC analysis of the ethanol leaf extract of *T. ciliata* was used to determine and quantify the phenolic compounds present in the extract. Several studies have shown that vanillic acid and quercetin possess antioxidant properties. In addition, catechin and ellagic acid compounds have been found to play a role in the anti-inflammatory activity [29, 30, 31] and rutin hydrate and quercetin are known to demonstrate good anti-inflammatory properties. [32, 33, 34] HPLC studies confirm the presence of relatively high concentration of these antioxidant chemicals in *T. ciliata*, which helps to explain the significant anti-inflammatory and antioxidant activity of this plant extract.

IX. Conclusion

The study demonstrates significant antioxidant and anti-inflammatory activity of the ethanol leaf extract of T. ciliata. Moreover phenolic compounds were detected with HPLC and a correlation can be suggested between the plant's antioxidant and anti-inflammatory properties and the high level of polyphenolic compounds present in its extract. Nevertheless, activity varies depending on several conditions including the plant type, its biome, growing conditions, etc. However, based on the results obtained, it can be asserted that the plant *T. ciliata,* grown in Bangladesh can be of great medicinal value in physiological processes and other cures, relieves and preventions.

X. Conflict of Interest

We declare that we have no conflict of interest.

Reference Références Referencias

1. The Ayurvedic pharmacopeia of India part -1. Vol.5. Govt of India Ministry of Health and Family welfare Department of Ayush, 179.
2. Chopra RN, Nayar SC, Chopra IC. Glossary of Indian Medicinal Plants.Council of Industrial and Scientific Research. New Delhi, 1986.
3. Da Saliva M, Fatima Das GF, Agasinho SMM, De Paula JR, Neto JO, Castro-Gamboa LF, Rodrigues FE, Fernandes JB, Vieira PC. Chemistry of *Tonna ciliata* & Cedrela odorata graft (Meliaceae): Chemosystematic & ecological significance. Pure Applied Chemistry. 1999; 71: 1083-7.
4. Gautam A, Jhade D, Ahirwar D, Sujane M, Sharma GN. Pharmacognostic evaluation of *Toona ciliata* bark. Journal of Advanced Pharmaceutical Technology and Research. 2010; 1(2): 216-20.
5. Sharma P, Yadav A, Ghule S, Malik P, Singh S. Anti-oxidant study of *Toona ciliata*. Pharmaceutical Research. 2009; 1: 114-7.
6. Hua-Dong C, Sheng-Ping Y, Yan W, Lei D, Jian-Min Y. Terpenoids from *Toona ciliate.* Journal of Natural Products. 2009; 72(4): 685-9.

© 2014 Global Journals Inc. (US)

7. Negi SJ, Bisht VK, Bhandari KA, Bharti MK, Sundriyal RC. Chemical and pharmacological aspects of *Toona* (Meliaceae). Research Journal of Phytochemistry. 2011; 5: 14-21.
8. Kiritikar KR, Basu BD. Indian Medicinal Plants. International Book distributors, Dehradhun 248001,1995; 562.
9. Malairajan P, Gopalakrishnan G, Narasimhan S, Jessi KVK. Analgesic activity of some Indian medicinal plants. Journal of Ethnopharmacology. 2006; 106: 425-8.
10. Chowdhury R, Hasan CM, Rashid MA. Antimicrobial activity of *Toona ciliata* and *Amoora rohituka*. Fitoterapia. 2003; 74: 155-8.
11. Shaohong C, Pengkang R, Yuntao Z. Inhibitory effects of ethanol extract from *Toona sinensis* leaves on the formation of protein non-enzymatic. Journal of Anhui Agricultural Science. 2010; 11: 5642.
12. Kumara SK, Sreedharamurthy S. Evaluation of antimicrobial and antioxidant activities from *Toona ciliata* Roemer. Journal of Analytical Science and Technology. 2013; 4: 23.
13. Lanhers MC, Fleurentin J, Dorfman P, Motrier F, PEAV JM. Analgesic, antipyretic and anti-inflammatory properties of *Euphorbia hirta.* Planta Medica. 1991; 57: 225-31.
14. Perianayagam JB, Sharma SK, Pillai KK. Anti-inflammatory activity of *Trichodesma indicum* extract in experimental animals. Journal of Ethnopharmacology. 2006; 104: 410-4.
15. Fan YJ, He XJ, Zhou SD, Luo AX, He T, Chun Z. Composition analysis and antioxidant activity of polysaccharide from *Dendrobium denneanum.* International Journal of Biological Macromolecules 2009; 45: 169-73.
16. Hemayet H, Ismet AJ, Sariful H, Jamil AS, Shubhra KD, Arpona H. Anti-inflammatory and antioxidant activities of ethanolic leaf extract of *Brownlowia tersa* (L.) Kosterm. Oriental Pharmacy and Experimental Medicine. 2013; 13: 181-9.
17. Dehpour AA, Ebrahimzadeh MA, Nabavi SF, Nabavi SM. Antioxidant activity of methanol extract of *Ferula assafoetida* and its essential oil composition. Grasas Aceites. 2009; 60(4): 405-12.
18. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. Analytical Biochemistry. 1999; 269: 337–41.
19. Hemayet H, Shahid-Ud-Daula AFM, Ismet AJ, Tarek A, Subrata B, Utpal K. Antinociceptive and antioxidant potentials of crude ethanol extract of the leaves of *Ageratum conyzoides* grown in Bangladesh. Pharmaceutical Biology. 2013; 51(7): 893-8.
20. Wootton-Beard PC, Moran A, Ryan L. Stability of the total antioxidant capacity and total polyphenol content of 23 commercially available vegetable juices before and after in vitro digestion measured by FRAP, DPPH, ABTS and Folin-Ciocalteu methods. Food Research International. 2011; 44(1): 217-24.
21. Hemayet H, Ismet AJ, Sariful IH, Jamil AS, Shubhra KD, Arpona H. Anti-inflammatory and antioxidant activities of ethanolic leaf extract of *Brownlowia tersa* (L.) Kosterm. Oriental Pharmacy Experimental Medicine. 2013; 13: 181-9.
22. Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. Journal of Food Drug Analaysis. 2002; 10: 178-82.
23. Ismet AJ, Proity NA, Nasima K, Tanzir AK, Mohammad MR, Arpona H, Hemayet H. Comparative study of anti-nociceptive activity and phenolic content of the ethanol extracts of *Piper nigrum* and *Piper longum fruits*. International Journal of Pharmaceutical Science Review and Research. 2014; 7: 47-52.
24. Seibert K, Zhang Y, Leahy K, Hauser S, Masferrer J, Perkins W, Isakson P. Pharmacological and biochemical demonstration of the role of cyclooxygenase-2 in inflammation and pain. Proceedings of the National Academy of Science of the United States of America. 1994; 91: 12013-7.
25. Cuman RKN, Bersani-Amadio CA, Fortes ZB. Influence of type 2 diabetes on the inflammatory response in rat. Inflammatory Research. 2001; 50: 460-5.
26. Hatano T, Edamatsu R, Hiramatsu M, Mori A, Fujita Y. Effects of the interaction of tannins with co-existing substances. VI: Effects of tannins and related polyphenols on superoxide anion radical and on 1, 1-diphenyl-2-picrylhydrazyl radical. Chemical and Pharmaceutical Bulletin. 1989; 37: 2016-21.
27. Singleton VL, Rossi JA. Calorimetry of total phenolics with phosphomolybdic acid-phosphotungstic acid reagents. American Journal of Enology and Viticulture. 1965; 16: 144-58.
28. Vinodhini V, Lokeswari TS. Antioxidant activity of the isolated compounds, methanolic and hexane extracts of *Toona ciliata* leaves. International journal of engineering and technology. 2014; 4(3): 135-8.
29. Guruvayoorappan C, Kuttan G. (+)-Catechin inhibits tumour angiogenesis and regulates the production of nitric oxide and TNF-α in LPS-stimulated macrophages. Journal of Innate Immunity. 2008; 14(3): 160-74.
30. Iñiguez-Franco F, Soto-Valdez H, Peralta E, Ayala-Zavala JF, Auras R, Gámez-Meza N. Antioxidant activity and diffusion of catechin and epicatechin from antioxidant active films made of poly (L-lactic acid). Journal of Agricultural Food Chemistry. 2012; 60(26): 6515-23.

© 2014 Global Journals Inc. (US)

31. Gainok J, Daniels R, Golembiowski D, Kindred P, Post L, Strickland R, Garrett N. Investigation of the anti-inflammatory, antinociceptive effect of ellagic acid as measured by digital paw pressure via the Randall-Selitto meter in male Sprague-Dawley rats. AANA Journal. 2011; 79: 28-34.
32. Kroes BH, Van den Berg AJ, Quarles van Ufford HC, van Dijk H, Labadie RP. Anti-inflammatory activity of gallic acid. Planta Medica. 1992; 58: 499-504.
33. Selloum L, Bouriche H, Tigrine C, Boudoukha C. Anti-inflammatory effect of rutin on rat paw oedema, and on neutrophils chemotaxis and degranulation. Experimental and Toxicologic Pathology. 2003; 54(4): 313-8.
34. Kleemann R, Verschuren L, Morrison M, Zadelaar S, van Erk MJ, Wielinga PY, Kooistra T. Anti-inflammatory, anti-proliferative and anti-atherosclerotic effects of quercetin in human in vitro and in vivo models. Atherosclerosis. 2011; 218(1): 44-52.

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine
Volume 14 Issue 7 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Glycemic Control and Self-Care Practice among Ambulatory Diabetic Patients in Ambo General Hospital, West Showa, Ethiopia

By Gobezie Temesgen Tegegne, Awoke Shiferaw, Belayneh Kefale Gelaw, Amsalu Degu Defersha, Minyahil Alebachew Woldu & Jimma Likisa Linjesa

Ambo University, Oromia region, Ethiopia

Abstract- *Background:* The main goal in diabetes care is to improve the patient's quality of life, to maintain satisfactory metabolic control and to retain minimal complications caused by diabetes mellitus (DM). To accomplish these goals, self-care has a great role. However, most patients don't control their blood sugar label. The objective of this study was to investigate status of glycemic control and self-care practice among Ambulatory diabetes patients in Ambo General Hospital.

Method: Cross-sectional study was conducted from 01 February to 30 May 30/ 2014. Sample population includes all diabetes patients who will come for checkup during data collection period. The data was collected with structured questionnaires and check list by trained data collectors. Data was cleaned and analyzed using SPSS version 20.

Keywords: *diabetes, self-care, fasting blood sugar, glycemic control, ambo general hospital.*

GJMR-B Classification : *NLMC Code: WD 200*



Strictly as per the compliance and regulations of:

© 2014. Gobezie Temesgen Tegegne, Awoke Shiferaw, Belayneh Kefale Gelaw, Amsalu Degu Defersha, Minyahil Alebachew Woldu & Jimma Likisa Linjesa. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Glycemic Control and Self-Care Practice among Ambulatory Diabetic Patients in Ambo General Hospital, West Showa, Ethiopia

Gobezie Temesgen Tegegne [α], Awoke Shiferaw [σ], Belayneh Kefale Gelaw [ρ], Amsalu Degu Defersha [ω], Minyahil Alebachew Woldu [¥] & Jimma Likisa Linjesa [§]

Abstract- Background: The main goal in diabetes care is to improve the patient's quality of life, to maintain satisfactory metabolic control and to retain minimal complications caused by diabetes mellitus (DM). To accomplish these goals, self-care has a great role. However, most patients don't control their blood sugar label. The objective of this study was to investigate status of glycemic control and self-care practice among Ambulatory diabetes patients in Ambo General Hospital.

Method: Cross-sectional study was conducted from 01 February to 30 May 30/ 2014. Sample population includes all diabetes patients who will come for checkup during data collection period. The data was collected with structured questionnaires and check list by trained data collectors. Data was cleaned and analyzed using SPSS version 20.

Results: The study showed that 58.5% of the respondents had type 1 diabetes and the remaining 41.5% were type2 DM. Mean FBS of three consecutive month was183.28 mg/dL. Only23.4%, 34.2% and 28.8% of the respondents were able to control their Fasting Blood Sugar (FBS) to level below 126 mg/dL during their last first, second and third visit to the hospital. 20(18.01%) of the respondents do exercise daily and attend their follow up program as scheduled respectively.

Conclusion: The present study illustrates that the level of knowledge about diabetes and self care practices amongst diabetic patients was low. In addition, it showed that respondents' level of physical activity, their educational status and time of insulin injection was low. Type 1 DM is the most prevalent type identified during the study period. In general, self-care practice was inadequate, especially in terms of physical self-care activity and a deficit in terms of knowledge related to diabetes; this could be explained by factors such as limited education and low levels of economic status

Recommendation: Health care providers should educate and promote health to address the lack of information on a healthy diet, benefits of exercise and how exercise should be undertaken for ambulatory DM patients when they come for regular follow up regularly at the hospital. Primary care physician must interpret (preferably in patient's language), for each diabetic patient, the short and long-term benefits of adhering to diet and exercise recommendations, insulin storage site in the hospital as well as at home must be given special emphasis.

Author α σ ρ ω ¥ §: Department of Pharmacy, College of Medicine and Health Science, Ambo University, Oromia region, Ethiopia.
e-mail: gob.ju2006@gmail.com

Keywords: diabetes, self-care, fasting blood sugar, glycemic control, ambo general hospital.

I. Introduction

a) Background

Diabetes mellitus is not a single disease entity but rather a group of metabolic disorders sharing the common underlying feature of hyperglycemia. Hyperglycemia in diabetes results from defects in insulin secretion, insulin action, or, most commonly, both. The chronic hyperglycemia and attendant metabolic dysregulation of diabetes mellitus may be associated with secondary damage in multiple organ systems, especially the kidneys, eyes, nerves, and blood vessels. It also greatly increases the risk of developing coronary artery disease and cerebrovascular disease. In concert with great technologic advances, there have been pronounced changes in human behavior, with increasingly sedentary life styles and poor eating habits. This has contributed to the simultaneous escalation of diabetes and obesity worldwide, which some have termed the "diabesity" epidemic. (6)

Prevalence of both type 1 and type 2 DM is increasing worldwide, type 2 DM is rising much more rapidly, presumably because of increasing obesity, reduced activity levels as countries become more industrialized, and the aging of the population [1].

According to IDF diabetes Atlas, 5th edition 2012 report, currently, more than 80% of people with diabetes live in low and middle income countries. The African region is expected to experience the highest increase in coming years withestimated increase in prevalence rates of 98% for sub-Saharan Africa, and 94% for North Africa and the Middle East (10, 11, 12, 13).It also said regional prevalence of 3.8%. This would rise to 4.3% 2030.Based on the IDF Atlas 5th edition, 2012 report, number of cases of diabetes in Ethiopia to be estimated about 1.4 million in 2011 (7, 8).

The diagnosis involves evaluation of blood glucose levels which are normally maintained in a very narrow range; usually 70 to 120 mg/dl that is established by elevation of blood glucose by any one of the following three criteria (1, 3)

© 2014 Global Journals Inc. (US)

A random blood glucose concentration of 200 mg/dL or higher, with classical signs and symptoms; A fasting glucose concentration of 126 mg/dL or higher on more than one occasion, or; An abnormal oral glucose tolerance test (OGTT), in which the glucose concentration is 200 mg/dL or higher 2 hours after a standard carbohydrate load (75 gm of glucose). (2)

b) Statement Of The Problem

DM is a life-long challenge that requires behavioral change and adequate self-care practices for better glycemic control. In the absence of appropriate self-care practice, the desired therapy targets are difficult, or even impossible, to achieve.

Thus, the aim of the present study was to assess self-care practices and glycemic control among diabetes patients in Ambo general hospital.

Non- communicable diseases including diabetes account for 60% of all deaths worldwide andmore than 80% of diabetes deaths occur in low- and middle-income countries. According to IDF Atlas 5th edition2012 report, Diabetes caused 4.6 million deaths in 2011 globally. World Health Organization projects that diabetesdeaths will double between 2005 and 2030. Statistics for medical complications from diabetes are also concerning. Proportions of patients with diabetic complications in sub Saharan region ranged from 7-63% for retinopathy, 27-66%for neuropathy and 10-83% for nephropathy. Diabetes is likely to increase the risk of several important infections inthe region, including tuberculosis, pneumonia and sepsis (14).Diabetes being a chronic illness requires continuous self-management practices bysufferers so that they can contribute meaningfully in the management of their lives. A situation where diabetes patients visit clinics regularly and their blood glucose levels still remain high despite the treatment they receive is problem that calls for attention. This is a very common observation in many diabetes patients. Severe complications, like gangrene that may lead to amputation and possible premature death, this might be because of lack of appropriate self-management practices (4).

Despite the benefits of engaging in a recommended self-management practice, research remains limited on determining recommended self-care practices level and its associated factors among diabetes patients. Researchers have suggested that self-care activities vary extensively according to the nature of the activity itself, with taking of medication often occurring as recommended and exercise frequently falling below recommended levels. For example, results from one study showed that 97% of respondents' with diabetes always or usually took their medication, whereas only 41% always or usually exercised, as cited by NancyE. Schoenberg (14). Furthermore, we currently lack an in-depth understanding of level and associated factors of type2 diabetes patients to ward diabetes self-care practices.

Although all forms of diabetes mellitus share hyperglycemia as a common feature, the underlying causes of hyperglycemia vary widely. The vast majority of cases of diabetes fall into one of two broad classes: (1)

Type1diabetes is characterized by an absolute deficiency of insulin secretion caused by pancreatic β-cell destruction, usually resulting from an autoimmune attack. It accounts for approximately 10% of all cases (5).

c) Significance Of The Study

The major problematic condition about diabetes self-care practices is that there is limited research findings on diabetic patients in our country, even there is no enough published material and little research is done on this areas. To address these deficits, this research explores for diabetes self-management practices. So the findings of this research can help diabetic patients to know how to control their sugar level and improve their self-care practice. Based on the findings of the research, patients of DM at Ambo hospital will be advised to strengthen those positive practices and will also be advised to practice appropriately by discouraging improper practices. The study can also serve as starting material for those who want to undertake further research on this area.

II. Objectives of The Study

a) General Objectives

- The aim of this study was to assess factors prevalence of glycemic control and self-management practices among ambulatory diabetic patients in Ambo general Hospital

b) Specific Objectives

- To determine prevalence of poor glycemic control.
- To assess dietary self-care practices among diabetic patients at Ambo General Hospital.
- To assess the physical activity and foot care practices among diabetic patients at Ambo
- General hospital.
- To assess blood glucose monitoring practice among the diabetic patients within three months.

III. Methods and Participants

a) Study area and period

This study was conducted in Ambo General Hospital, which is found in Ambo town, west showa, Ethiopia. Ambo has a total population of 260, 193 of whom 131, 922 are men. It is located in the west Showa zone of the Oromia region, 114 km West of Addis Ababa, the town has a latitude and longitude of 8.983O N 37.850O E and an elevation of 2101m. The town has an annual rain fall of 1012 mm with 18 CO average temperatures. The Hospital is found in 01 kebele of Ambo town near West Shewa zone prison House. The hospital give serves for about 10,000 People. The study was conducted from February 1/2014 to .May/2014

© 2014 Global Journals Inc. (US)

b) Study design

Cross-sectional study was conducted from 01 February to May 30, 2014.

c) Source population

Source population includes all diabetes patients that attend their follow up schedule in Ambo general hospital.

d) Study population

Study population includes all diabetes patients that follow their diseases status in Ambo general Hospital for the last three months.

e) Sampling size Determination

All DM patients who came to Ambo general hospital for follow up during data collection period were considered to be included in the sample. Accordingly 111 patients fulfilled the inclusion criteria and were included in the study. Six of the patients that didn't fulfilled the inclusion criteria were excluded from the study.

f) Inclusion and exclusion criteria

i. *Inclusion criteria*

Patient that has been part of a follow-up program for at least three follow at Ambo General Hospital was included in the study.

ii. *Exclusion criteria*

Patients with mental health problems, hearing impairments or any other serious health problems and those patients who were unable to provide the appropriate information were excluded.

g) Study Variables

i. *Dependent variables*

Self-care practice among diabetic patients, glycemic control

ii. *Independent variables*

Socio demographic characteristics of study population

h) Data collection procedure and Instrument

Patients were interviewed using structured questionnaires and check lists was used to gain information from their card. The data collection was conducted by the joint collaboration of the investigator, nurses and health care professionals that are involved in delivery of care to the specified patients in the hospital by orienting them on how to collect the data. Questionnaires were prepared in English and translated into Amharic and Afan Oromo (local languages) and translated back into English to check its consistency. To identify the patterns of glycemic control, patients' charts was reviewed, retrospectively; the last three successive FBS or RBS results was recorded from the patient's card.

i) Data quality control

Pretest was done in 10 patients at Ambo Hospital to assure validity of the check list and questionnaire. Language experts who were qualified with second degree with linguistic and are Ambo University stuff members were used to translate the questionnaires from English to Amharic and Afan Oromo version. The questionnaires were revised for its completeness and consistency.

j) Data analysis

The data was cleaned, coded, entered and analyzed using SPSS version 20. Categorical variables were described by frequencies and percentages, and continuous variables were described by means and standard deviations. Figures and tables were used to summarize the results.

k) Ethical consideration

Formal letter was obtained from Research Ethics Committee of Ambo University and submitted to Ambo General Hospital, so the letter was given to the hospitals and they allowed us to do the research. Verbal consent was taken from the patient.

l) Operational definition

The level of glycemic control was indicated as 'adequate glycemic control' when FBS results were less than 126 mg/dL (7 mm/L) (i.e. an average of three visits), or when RBS results were less than 200/dL; 'inadequate glycemic control' takes place when a parameter is beyond the criteria of adequate glycemic.

i. *Knowledge*

Knowledge of patients' relating to diabetes and self-care practice was assessed by making use of 'yes/no' questions. A correct answer will be coded as '1' and an incorrect answer as '0'; the score is then computed. Respondents are labeled as having knowledge of diabetes and self-care practices if he or she scored ≥ the mean value, and having poor knowledge if he or she scored less than the mean.

ii. *Physical activity*

The levels of physical activity of the patients were classified into three levels based on their physical activities as light, moderate and heavy.

1. Light activity: Patients are in a sitting position most of the time, less than half of the time they are standing or walking, they seldom carry heavy things, and travel by car or motorbike.
2. Moderate activity: Patients are sitting, standing and walking about half of their time. They spend some time carrying heavy things and use public transport during non-leisure hours.
3. Heavy activity: Patients spend almost none of their time sitting and almost all of their time standing or walking, most of the time carrying heavy things, and they use public transport, cycle or walk everywhere.

Self-care

Self-care means looking after yourself in health way. This includes changes to your diet, different types

© 2014 Global Journals Inc. (US)

of exercise or different types of medication you may need to make.

m) Limitations of the study

The findings from this situational analysis may not be generalized to the total population because of the lower sample size than the expected.

IV. Results

a) Socio-demographic characteristics of the participants

A total of 111diabetic patients were participated in the study giving a response rate of 100%. From total respondents 63 (56.75%) were male, regarding the age of participants, 33 (29.7%) of them were younger than 30 years of age and the remaining 78 respondents (70.27%) were above age of 30. Most of them completed grade 7-12 and 29 patients had monthly income of 500-800 birr.

Table 1 : socio-demographic characteristic of the patients in ambulatory diabetic patients

Characteristic type	Characteristic	Frequency (%)
Sex	Male	63(56.75)
	Female	48(43.25)
Age	1-29	33(29.7)
	30-44	31(27.9)
	45-64	37(33.3)
	Above 65	10(9)
Marital status	Single	42(37.8)
	Married	65(58.5)
	Widowed	4(3.3)
Religion	Orthodox	64(57.6)
	Muslim	8(7.2)
	Protestant	37(33.3)
	Others	2(1.8)
Ethnicity	Oromo	96(86.7)
	Amhara	14(12.6)
	Tigre	1(0.9)
Occupation	Farmer	38(34.2)
	Merchant	20(18)
	Civil servant	28(25.2)
	Others	25(22.5_)
Education level	Illiterate & non formal edu.	18(16.2)
	Grade 1-6	26(23.4)
	Grade 7-12	44(39.6)
	Above grade 12	23(20.7)
Income	<500	51(45.9)
	500-800	29(26.1)
	801-1500	16(14.4)
	>1500	15(13.5)

b) Clinical characteristics of the patients

Regarding the clinical characteristics of the patients, from the total of 111 patients 65 of them were diagnosed for type 1 DM (58.5%) and the remaining 46 of them (41.5%) were type 2 DM.70.27% say no family member with DM and the rest 29.73% say there is a family member with DM. Concerning the presence of other co morbidities 17. 11% have hypertension and 6.3% of the patients have CKD. When we look their respective drug use, 35 patients which contribute 31.53% use glabinclamide, 22 patients which contribute about19.81% use metformin, 6 patients which contribute 5.1% use both insulin & metformin, and 65 patients which contribute about 58.55% use insulin. Out of the 65 insulin users, 10 patients (15.38%) use refrigerator, 55(84.61%) use home prepared cool sites.

Of the patients who were using insulin, only 23 (20.7%) took meals 30 minutes after each insulin injection, and the remaining patients were used to eat before injection while 70.2% took meals after one hour of taking an injection. The majority of the respondents (104, 93.7%) follow their medication strictly to avoid

© 2014 Global Journals Inc. (US)

raise in blood sugar level. From the total respondents, 96 of them (86.5%) attend their follow up program as per the schedule.10.8% patients have habit of alcohol or smoking and the remaining 99 respondents (89.2%) do not have any habit.

Means of communication was assessed to know how diabetic patients can obtain information or education from Medias and from newspapers as well as to know whether they have phone in case they face emergency conditions(hypoglycemia or hyperglycemia) to obtain health services. The results shows the majority of the patients have access to radio, TV and phone services and 91 respondents (82%) do not have the chance to get newspapers.

Table 2 : clinical characteristics of the patients

Variables	*Frequency (%)*
Type of DM	
Type1 DM	65(58.55)
Type 2 DM	46(41.44)
Time since DM d	
<2000	6(5.4)
2000-2002	33(29.7)]
2003-2006	72(64.89)
Hx of DM in the family	
Yes	33(29.73)
No	78(70.27)
Co morbidities	
CKD	7(6.3)
HTN	19(17.11)
CHF	1(0.9)
Stroke	0(0)
Others	13(11.71)
Pattern of drugs for DM	
Insulin	65(58.55)
Oral hypoglycemic agents	
Metformin	22(19.81)
Glabineclamide	35(31.53)
Insulin &metformin	6(5.4)
Source of information for DM	
Radio Yes	83(74.8)
No	28(25.2)
TV Yes	60(54)
No	51(46)
Phone Yes	58(52.2)
No	53(47.8)
Newspaper yes	20(18)
No	91(82)

c) Knowledge about diabetes

Participants were asked whether DM is a chronic disease or a curable disease and whether it is possible to control it by interventions, such as a healthy diet, exercise, and administering insulin and hypoglycemic drugs. Accordingly, 72 respondents (64.86%) responded that it is chronic disease 39 respondents (35.14%) said that DM is curable and 96respondents (86.48%) reported that it is possible to control diabetes. Furthermore, the majority (83.78%) of the respondents knows the sign and symptoms of DM and the remaining (16.22%) do not know the signs and symptoms of DM.

d) Self- care practice

i. *Dietary self -care practice*

Concerning to food items that they consume to control their sugar level, the majority of the respondents answered that injera (i.e. a stable food diet in Ethiopia made of Teff cereal), barely and kocho (i.e. a traditional staple food made of a false banana plant called enset or Ensete Scitamineae) have a low glycemic index and could be eaten freely by diabetic patients; Only 74 respondents (66.7%) stated that fibrous food (e.g.whole grain cereals) has a high glycemic index and similarly 29 respondents (26.1%) do not know the sign and symptoms of hypoglycemia.From those who know the sign and symptoms of hypoglycemia, 50% uses candy to control their sugar, 46% uses table sugar and the remaining uses soft drinks. 59.5% have regular time for meal whereas the remaining 41.5% do not have regular meal time. The majority of the respondents (103, 92.8%) eat three times per day.

© 2014 Global Journals Inc. (US)

Table 3 : Type of food consumed by diabetic patients

Type of food they consume *Frequency (%)*	
Type of food they consume	**Frequency (%)**
Injera (teff)	79(71.17)
Barely	73(65.76)
Kocho	13(11.71)
Others	17(15.31)
Food they take during hypoglycemia	
Table sugar	51(46)
Candy	55(49.54)
Soft drink	31(27.92)
Others	11(9.9)

ii. *Physical activity self- care practice*

The physical activity that the respondent performs was summarized in the table 4 below. Accordingly, 49.54%practice hard work such as farming, daily laborer, 21.62% perform aerobic exercise, and the rest 23.42%were involved other activities. Most of the respondents, 45.05% do physical exercise sometimes whereas 18.01% do exercise daily and the rest 36.94% never do physical exercise. With regard to practice of walking, 91 respondents (81.98%) do walk and the rest 20 respondents (18.02%) do not practice walking. 95.49 %) of patients has shoe wearing practice whereas the rest 5 do not. There is 110 (99.09 %) daily and 1(0.91%) twice a week foot washing practice among the respondents. Of the total respondents 99 (89.19%) have not sustained foot injury. Of those sustained foot injury 9get health care's and the rest 3get to traditional healers for support.

Table 4 : Physical activity self-care among DM patients of Ambo general Hospital, WestShoa, Ethiopia April –May 2004

Question	Response	Number of respondents (%)
1 What type of physical activity do you practice for your health	Hard work	55(49.54
	Aerobic exercise	24(21.62)
	Swimming	4(3.60)
	Driving bicycle	2(1.80)
	Others	26(23.42)
2 How often do you perform physical exercise	Daily	20(18.01)
	Sometimes	50(45.05)
	Never	41(36.94)
3 Practice of Walking on foot	Yes	91(81.98)
	No	20(18.02)
5 Do you wear shoe for your foot care	Yes	106(95.49)
	No	5(4.51)
6 How often do you wash your feet	Daily	110(99.09)
	Twice a week	1(0.91)
	Weekly	0(0.00)
	Do not wash	0(0.00)
7 Sustaining of foot injury	Yes	12(10.81)
	No	99(89.19)
8Actions taken in case of foot injury	Get health care	9(8.3)
	traditional healers	3(2.7)
	Heal spontaneously	0(0.00)

iii. *Self-Blood glucose monitoring, medication self-care, and Foot care practice*

There were 102 patients practicing blood glucose monitoring at home. 79.27% of the respondents do not forget to use their medication, whereas 98(88.28%) do not miss their medication intentionally. Most patients (106, 95.49%) do not interrupt their medication due to side effect or when feeling free of the disease but the rest 4.51 % (5 patients) interrupt their medication due to side effect or when feeling free of the disease. 99 of the respondents (89.19%) do not forget their medication while travelling but the rest 12(10.81%) do. With regard to the presence of health problem other than DM, 82 respondents (73.87%) say no and the rest 29 patients (26.13%) say yes. 78 patients(70.27%) say no family member with DM and the rest 33(29.73%) say there is a family member with DM.

© 2014 Global Journals Inc. (US)


When we look their respective drug use, 35 patients which contribute 31.53% use glabinclamide, 22 patients which contribute about19.81% use metformin, 6 patients which contribute 5.1% use both insulin &metformin, and 63 patients which contribute about 56.76% use insulin. Out of the 69 insulin users, 10 patients (14.49%) use refrigerator, 53 patients (76.81%) use home prepared cool sites and 6 patients (8.70%) use other methods for storage of the drug at appropriate temperature.

Table 5 : Self- Blood glucose monitoring, medication self-care and Foot care practice among DM patients of Ambo general Hospital in West Shoa, Ethiopia April –May 2004

Question	Response	Number of respondents(%)
1-Blood glucose monitoring practice at home	Yes	9(8.11)
	No	102(91.89)
2-Forgetting medication use	Yes	23(20.73)
	No	88(79.27)
4- Medication missing for a reason other than forgetting	Yes	13(11.72)
	No	98(88.28)
6- Medication interruption due to side effect	Yes	5(4.51)
	No	106(95.49)
7- Forgetting medicine while traveling	Yes	12(10.81)
	No	99(89.19)
8-Interruption of medication when feeling free of the disease	Yes	5(4.51)
	No	106(95.49)

iv. *Glycemic control among respondents*

Check list was used to assess patients card to know their year of diagnosis, type of DM and their fasting blood sugar level for the last three months and the data obtained shows from the total of 111 patients 65 of them were diagnosed for type 1 DM (58.5%) and the remaining 46 of them (41.5%) were type 2 DM. Concerning the year of first diagnosis 72 of them (64.8%) were diagnosed after 2003, 33 of them (29.7%) were diagnosed from 2000-2002 and the remaining before 2000.




Figuer 1 : Year of first diagnosis of DM patients at Ambo General Hospital West Shoa, Ethiopia

© 2014 Global Journals Inc. (US)

To know the sugar level of the patients each patient's card was revised to obtain their FBS level during their last three visits to Ambo Hospital and the result obtained was presented in the chart below

The chart below shows that the mean FBS of the respondents during the last visit to the hospital was 183.28. Out of the total 111 patients who visited Ambo Hospital during the third month 62 of them (55.8%) were tested to have FBS level of above 126, 13 (11.7%) have IFG level,32(28.8%) of them maintained their normal blood sugar level and 4 patients fall in hypoglycemia. During their second visit 38 patients (34.2%) maintained the FBS level within the normal range and 54 of them (48.6%) were in hyperglycemia. Similarly the FBS of the patients during their first visit shows 74 of them (66.65%) were tested to have FBS of above 126, 9 of them (8.1%) impaired sugar level,26 of them(23.4%) maintained their normal FBS.



Figuer 2 : FBS measurements for the last three months at Ambo general hospital West Shoa, Ethiopia April –May 2004

V. Discussion

The present study attempted to assess diabetes mellitus Patients' knowledge and self-care practices in terms of living with the disease. According to the findings of the study, most of the respondents know that DM is chronic disease. But 35.14% of the respondents think DM is curable and 16.22% of the respondents even do not know the sign and symptoms of DM. This is in agreement with the study conducted in Jima University [9] in which 34.21% of the patients responded DM is curable. The majority of the respondents consume teff, barely and kocho which they think have low glycemic index and could be eaten freely by diabetic patients. Diabetes self-management behaviors such as diet and exercise involve and depend on guidance from a health care provider, meal preparation in a family context and exercising with a partner [1].

The study also shows physical activity level of the patients were sub optimal and 41(36.42%) respondents never do any physical activity. This is in line with the study conducted at JimmaUniversity [8] in which most of the respondents has sub optimal physical exercise which could be because of having inadequate knowledge in terms of the benefits of regular physical exercise and a fear of hypoglycemia. Exercise has multiple positive benefits including cardiovascular risk reduction, reduced blood pressure, maintenance of muscle mass, reduction in body fat, and weight loss. For individuals with type 1 or type 2 DM, exercise is also useful for lowering plasma glucose (during and following exercise) and increasing insulin sensitivity [1]. In patients with diabetes, the ADA recommends 150 min/week (distributed over at least 3 days) of moderate aerobic physical activity. The exercise regimen should also include resistance training.

The study also showed that the majority of the respondents 95.49% wear shoe and 99.09% of the respondents wash their foot daily which is very important to decrease the complications associated with the disease. DM is the leading cause of non-traumatic lower extremity amputation. Patient education should emphasize careful selection of footwear, daily inspection of the feet to detect early signs of poor-fitting

© 2014 Global Journals Inc. (US)

footwear or minor trauma, daily foot hygiene to keep the skin clean and moist, avoidance of self-treatment of foot abnormalities and high-risk behavior (e.g., walking barefoot), and prompt consultation with a health care provider if an abnormality arises (1).

Of the patients who were using insulin, only 23 (20.7%) took meals 30 minutes after each insulin injection, and the majority 70.2% of the patients were used to eat before injection. Regular insulin is given 30–45 min prior to a meal [1]. 6.3% of the respondents miss their medication and 13.5% do not attend their follow up as scheduled for them. More than half of the respondents 65(58.5%) were type 1 DM. Diabetes is managed via a regimen of control. Physicians advise adults living with type 2 diabetes to control blood sugar levels by controlling diet, maintaining regular exercise, and complying with medication. The extent to which individuals are able to adhere to such recommendations varies (5).

Regarding the FBS of the patients for the last three months the mean FBS of the last visit was found to be 183.28mg/dl which are far from the normal glucose homeostasis. Generally the results of their last three visit shows only, 32(28.8%) of them during the third month, 38 patients (34.2%) during the second month and 23.4% of them during the first visit maintained their sugar level within the normal range. This is consistent with other studies [10] who reported adequate glycemic control in 43.8% of type -2 diabetic patients. This indicates that most of the patients were not controlling their blood glucose level, despite most of them taking medication provided for them. Despite the increasing prevalence of diabetes, improved understanding of the disease, and a variety of new medications, glycemic control does not appear to be improving. Self-monitoring of blood glucose (SMBG) is one strategy for improving glycemic control; however, patient's adherence is suboptimal and a proper education and follow-up are crucial. SMBG should include post-prandial monitoring to identify glycemic excursions after meals, to indicate the need for lifestyle adjustments, and to provide patients' feedback on dietary choices (12).

Discussing the adequacy of glycemic control will be a handicap without mentioning glycosylated hemoglobin (HbAlc) determination; the ADA recommends that a patient should have glycosylated hemoglobin determination at least twice yearly. In addition, a study conducted in the United States of America (USA) showed that at least 77% of diabetic patients had at least one glycosylated hemoglobin determination in the two years preceding the study. However, none of the patients had glycosylated hemoglobin. Our findings, therefore, suggest that monitoring of glycemic control among DM patients at Ambo Hospital west Shoa Ethiopia may be less than optimal and this may be a probable contributory factor to late detection of patients at risk of complications and death from poorly controlled diabetes.

To know and follow self care practices anthropometric measurements are also important parameters for diabetic patents. But none of them have anthropometric measurement data.

Self-care activity in diabetic management includes medication self-care, dietary self-care, physical activity self-care and self-monitoring of blood glucose levels.

VI. Conclusion and Recommendation

a) *Conclusion*

In general, self-care practice was inadequate, especially in terms of physical self-care activity and a deficit in terms of knowledge related to diabetes. In some cases the patients do not attend their follow up strictly and sometimes miss taking their medications. Almost all the patients have good foot care hygiene. Regarding insulin injection time with respect to meal, majority of them take their medication after eating. The mean fasting blood sugar of the patients during the last visit was not controlled.

b) *Reccomandations*

The following recommendations are forwarded.

Health care providers: should educate and promote health to address the lack of information on a healthy diet, benefits of exercise and how exercise should be undertaken for ambulatory DM patients when they come for regular follow up regularly at the hospital

Primary care physician must interpret (preferably in patient's language), for each diabetic patient, the short and long-term benefits of adhering to diet and exercise recommendations, Insulin storage site in the hospital as well as at home must be given special emphasis.

Patients should learn to follow their medication strictly including the site of injection care for their diet with regular meal time and understand the benefit of physical exercise.

The hospital follow up schedule must not be longer than one month and patients must adapt to frequently visit the hospital for regular checkup. Patients can learn to control their blood sugar level at home by using glucometer.

Generally self -care practice including blood glucose monitoring is the back bone to control DM, hence further investigation by researchers is needed to strengthen diabetic self-care practice and promote health.

VII. List of Abbreviations

ADA = American Diabetic Association

ATP = Adenosine Triphosphate

DM = Diabetes Mellitus

DNA = DeoxyriboNucleic Acid

© 2014 Global Journals Inc. (US)

FBS = Fasting Blood Sugar

HbA1c = Glycosylated Hemoglobin

OGTT = Oral Glucose Tolerance Test

RBS = Random Blood Sugar

IFG = Impaired fasting glucose

CKD = Chronic kidney disease

CHF = Congestive heart failure

VIII. Acknowledgements

We are very grateful to our college staff members for unreserved guidance and constructive suggestions and comments from the stage of proposal development to this end. We would like to thank Ambo University for supporting the budget which required for this research. Finally our deepest gratitude goes to Ambo General Hospital staff workers who help and allow us in conducting this research.

Conflict of interest : none declared

Reference Références Referencias

1. Kasper L.Harrison's Principle of Internal medicine.18thed.Graw hill company,2012;p.2275-2311
2. Hennekens H. Increasingburde of cardio vascular disease. J Am Heart Association 1998;p.1095-1102
3. World Health organization. Definition, diagnosis and classification of Diabetes Mellitus and its complication. Geneva; 2007
4. American Diabetes Association. Standards of medical care in diabetes .DIABETES CARE.2008; 31(1):43.
5. Sigurdardottor A. Self-care in diabetes: Models of factors affecting self-care.Jclin Nur.2005;14:301-14.http://dx.doi.org/10.1111/j.13652702.2004.01043.x,pmid:15707440
6. Hilary G.magazine of WHO. Geneva; 1998.
7. Lin C, Anderson R, Hagerty B, Lee B. Diabetesse lf management experience among Taiwanese patients with type two DM.J Clin Nurs.2007;252-267.
8. Tan M, Magarty J. self-care practice of Malaysian adults with diabetes and sub optimal glycemic control patients. Educ counseling 2008; 72:252-276.
9. Eriksson J, Lindstorm J, Valle T, Anulo, Hamalaines H, Hannep. prevention of type II diabetes in subjects with impaired glucose tolerance. Diabetologia. 1999;42(7):793-801.htt://dx.doi.org.
10. http://emedicine.medscape.com/article/117853workup. Accessed in 10/02/2014
11. (http://www.bettermedicine.com/article/diabetes.Accessed in 02/02/2014
12. http://www.worlddiabetesfoundation.org/. Accessed in 05/02/2014
13. http://www.worlddiabetesfoundation.org/media.Accessed in11/03/2014
14. http://www.who.int/mediacentre/factsheets/fs312/en/index.html. Accessed in11/03/2014

© 2014 Global Journals Inc. (US)

Global Journal of Medical Research: B
Pharma, Drug Discovery, Toxicology and Medicine
Volume 14 Issue 7 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Assessment of Knowledge, Attitude and Practices Regarding Life Style Modification among Type 2diabetic Mellitus Patients Attending Adama Hospital Medical College, Oromia Region, Ethiopia

By Abdulkadir Mustefa Adem, Esayas Tadesse Gebremariam, Belayneh Kefale Gelaw, Mustefa Ahmed, Muluneh FromsaSeifu & Dr. Thirumurugan G.

Ambo University, Ethiopia

Abstract- *Background:* The person with diabetes mellitus has a chronic lifelong disease, the person must be knowledgeable to coordinate life style modification in to a daily routine of work to achieve and maintain normal physiological blood glucose level. The objective of this study was to determine the knowledge, attitude and practices of LSM management of type 2 diabetic patients in Adama Medical college Hospital.

Methodology: Across-sectional study was conducted to assess knowledge, attitude and practice of type 2 diabetic patients towards LSM management of DM in Adama Medical College Hospital. Data was cleaned and analysed by using SPSS version 16.0 and presented descriptively and analytically.

Keywords: *type 2 diabetes mellitus, life style modification, knowledge, attitude, practice.*

GJMR-B Classification : *NLMC Code: WD 200*



Strictly as per the compliance and regulations of:

© 2014. Abdulkadir Mustefa Adem, Esayas Tadesse Gebremariam, Belayneh Kefale Gelaw, Mustefa Ahmed, Muluneh FromsaSeifu & Dr. Thirumurugan G. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Assessment of Knowledge, Attitude and Practices Regarding Life Style Modification among Type 2diabetic Mellitus Patients Attending Adama Hospital Medical College, Oromia Region, Ethiopia

Abdulkadir Mustefa Adem [α], Esayas Tadesse Gebremariam [σ], Belayneh Kefale Gelaw [ρ], Mustefa Ahmed [¥], Muluneh FromsaSeifu [§] & Dr. Thirumurugan G. [χ]

Abstract- Background: The person with diabetes mellitus has a chronic lifelong disease, the person must be knowledgeable to coordinate life style modification in to a daily routine of work to achieve and maintain normal physiological blood glucose level. The objective of this study was to determine the knowledge, attitude and practices of LSM management of type 2 diabetic patients in Adama Medical college Hospital.

Methodology: Across-sectional study was conducted to assess knowledge, attitude and practice of type 2 diabetic patients towards LSM management of DM in Adama Medical College Hospital. Data was cleaned and analysed by using SPSS version 16.0 and presented descriptively and analytically.

Result: Concerning knowledge of the patients towards LSM management of diabetic; majority of the patients were knowledgeable which accounts 90(77.59%) followed by 13(11.21%) patients fairly knowledgeable and the other 13(11.21%) patients were poorly knowledgeable. Regarding attitude of the patients 95(81.89%) patients had positive attitude and the other 21(18.11%) had fair attitude. In another way almost half of the patients 57(49.1%) had good practice. The other 39(33.62%) and 20(17.24%) have poor and average practice respectively.

Conclusion and Recommendation: The result of this study showed, majority of type2 DM patients had good knowledge, positive attitude and good practices towards LSM. The researcher recommend all stake holders (Ministry of health, Diabetic associations, Health institution, health professionals, caregivers and NGO) found around this area must cooperate to improve KAP of the patients towards LSM. Especial attention should be given to the practices of the patients and further research should also be done on this topic.

Keywords: type 2 diabetes mellitus, life style modification, knowledge, attitude, practice.

Author α σ ρ Ω ¥ § χ: Department of Pharmacy, College of Medicine and Health Science, Ambo University, Ambo, Ethiopia.
e-mail: abdulkadir.mustefa@yahoo.com

I. Introduction

a) Background

Diabetes mellitus is a syndrome characterized by chronic hyperglycaemia, due to absolute or relative deficiency or diminished effectiveness of circulating insulin. It is the most common serious metabolic disease. Diabetes mellitus has been recognized as a clinical syndrome since ancient times and remains a crippling global health problem today [1]. Its clinical diagnosis indicated by presence of symptoms such as polyurea, polydipsia and unexplained weight loss; and confirmed when one of the following abnormal glucose measured: fasting plasma glucose (FPG) value of >= 7.0 mmol l-1 (126 mg dl-1), or the casual plasma glucose value >= 11.1 mmol l-1 (200 mg dl-1), or if the plasma glucose value 2 hours after a 75g oral load of glucose is>= 11.1 mmol l-1 (200 mg dl-1). However, in asymptomatic subjects, the test should be performed more than one occasion to confirm the diagnosis and treat the subject [2].

There are different categories of DM; but the two broad categories are type1 and type2.Both types of diabetes are preceded by a period of abnormal glucose classified as impaired fasting glucose (IFG) or impaired glucose tolerance (IGT). Type 1 DM is the result of complete or near-total insulin deficiency. Type 2 DM is a heterogeneous group of disorders characterized by variable degrees of insulin resistance, impaired insulin secretion, and increased glucose production [3].

The prevalence of type 2 diabetes is increasing in the world at high rate. For example according to the study of Diabetes Screening in Canada (DIASCAN) in the year 2000; the prevalence of Type 2 diabetes in individuals 40 years of age and older who see a general practitioner was 16.4% in Canada and nearly 20% in Québec. It was predicted that the number of Canadian diabetics will double within the next 15 years. The impact of diabetes on the health of populations and

© 2014 Global Journals Inc. (US)

individuals afflicted with the disease is primarily related to late-stage complications. Diabetes is the primary cause of terminal kidney failure, blindness before the age of 65 and amputations [4].

The burden of diabetes is increasing in the worldwide including developing countries like Ethiopia. International diabetic federation association reported Ethiopia to be ranked 3rd in Africa with 1.4 million DM and prevalence of 3.32 by year 2012.The study done to determine prevalence of un diagnosized DM & its risk factor in institution of Bishoftu town, East Shoa at 95% confidence interval with (p=0.05 and margin of error 5%) in the year 2012/2013 reported the prevalence of undiagnosed DM to be 5% [5].

Ancient civilizations in Egypt, Greece, Rome and India recognized diabetes mellitus and recommended dietary modifications. During previous century's recommendation about dietary carbohydrates from diabetic individuals were based on theory rather than scientific facts. Prior to discovery of insulin, diabetes was treated with low carbohydrate, semi starvation diet. Even after insulin was discovered in 1921, most eastern diabetes specialists used low carbohydrates, high fat diets to treat diabetic individuals [6].

Today ;essential components of the treatment for diabetes include diabetes self management education, lifestyle interventions, and goal setting glucose management and pharmacologic management of hypertension and hyperlipidemia[7]. Life style modifications are considered the corner stone of management of diabetes mellitus and include the prescription of healthy diet, regular exercise, and avoidance of tobacco [8] .

Education regarding diabetes is very important to improve the life style of the patients which would be helpful in maintaining (controlling) blood glucose. Observation has been reported that improper guidance and communication can lead to poor compliance to both medication and life style [9].

Dietary recommendations for DM patients focus on the reduction of fat intake and increase of vegetable consumption with moderate calorie restriction and it should be individualized according to the patients' physical activity, co morbid condition and personal preferences. Da Qing study in China showed diet intervention alone associated with 31% reduction in the risk of developing type2 DM [10].

In the other way alcohol intake exacerbate neuropathy, dyslipidemia and obesity. Therefore it should be prohibited, if used it must be in moderation .Similarly smoking should be prohibited as it increases risk of complication [7].

Physical activity (exercise) also reduces the risk of developing DM type2 by 30-50%.Physical activity improves insulin sensitivity .eg It can reduces free fatty acid load to the liver; there by reduces hepatic insulin resistance .Moderate exercise as little up to 30min/day or 150min/week can show the differences [11].

b) Statement of the problem

Despite the availability of different treatment modalities, diabetes has remained a major cause of death. Now a day's 3.8 million deaths are attributable to the diabetes in each year worldwide. It is the 4th leading cause of global death. In 2005 there were 246 million people worldwide were affected with diabetes and are expected to affect 380 million by 2025, over a seven fold increase just over 20 years [12]. This indicates high burden of DM in today world and developing countries account for a substantially high proportion [13].

In Ethiopia; from hospital based studies; it was observed that the prevalence of diabetes admission had increased from 1.9% in 1970 to 9.5% in 1999 of all medical admissions [14]. World Health Organization (WHO) also estimated the number of diabetics in Ethiopia to be about 800,000 cases by the year 2000, and the number is expected to increase to 1.8 million by 2030 [15].

In another way, according to the 2011 report of the International Diabetes Federation (IDF), the number of adults living with diabetes in Ethiopia was 3.5%. A study done on urban Commercial Bank employees in Ethiopia showed a 6.5% prevalence of DM [16] which indicated the significance of lifestyle for DM aetiology and its burden to our country too.

Although, the diabetes is causing high wastage of life and resources the management is still low .Its management include pharmacologic and life style modification. However, LSM is ignored by many of the patients and care givers. For example Study done in Omani on type 2 DM patients in the year 2013 reported only less than 40% of the patients participate on regular exercise [17]. The other study done at USA in the year 2008 showed among 69 patients classified as elevated risk of diabetes only 17%,32% and 30% had received advice for weight loss ,exercise & diet modification [18].

Another similar cross sectional study done in four provinces of Kenya in the year 2010 found that; of the people participated in the study 75% had poor dietary practices, while 72%didn't participate in regular exercise & about 80% didn't monitor their body weight [19].

In another way, study done at Jimma University Specialised Hospital in the year 2011to assess quality of care given to diabetic patients showed that there was no attention given to diabetic education in Ethiopia. There were no diabetes nurse educators and diabetes dietician in the country and those who provide health services for diabetes had no special training for diabetes care [20].

In AMCH the diabetes patients appointed to follow up clinic according their disease status. Most patients come to the Hospital every month or every two month and refill their medication. They check up their

© 2014 Global Journals Inc. (US)

blood glucose only when they come according to their follow up appointment and contact the physician for less than five minutes for refill. Most of the time they asked only for adherence to medication and do not checked awareness for life style intervention. In certain circumstances both health care giver and patients do not raise all about non pharmacologic management of diabetes including new diagnosed type2 DM patients. These observations raised the researcher' concern about the knowledge, attitude and practice of lifestyle modifications among diabetic patients at AMCH, which this study seeks to explore and document. Since there was no similar study in the area, this information gap about of LSM among AMCH type 2 DM patients could be answered by this study.

c) Significance of the study

It is obvious that there is no adequate information on knowledge, attitude and practice of non pharmacologic management of diabetes in our country; Ethiopia. Most of literatures are drafted from developed countries like USA, Russia and European countries.

Majority of the studies were done to evaluate the knowledge, attitudes and practices about diabetes mellitus among diabetics rather than on lifestyle modifications.

The result of this current study could be useful as a base line in implementing a community based awareness programme which will promote the importance of lifestyle modifications in the prevention and control of non-communicable diseases, particularly diabetes mellitus. Therefore it is helpful for all stake holders that involves in this areas i.e patients, caregivers, health care providers, health institution and policy makers (MOH, NGO, diabetic DM association)

The other benefit of this study is that the result of this study helps health care provider to give education and awareness on LSM management of DM. In addition DM patients' and family members realize the benefit of non pharmacologic management hand by hand with pharmacologic management for the success control of the disease and its complication. Thus the patients' care will improve and the patients' will be benefited.

Finally, Adama Medical College Hospital also uses this result for improving its services for DM patients and other chronic disease patients. It also helps to encourage good services of the clinics and modify the possible problems of between hospital mission and actual services given by diabetic follow up clinic .The hospital management can use this result to expand its services especially, this can help AMCH and diabetic association as feedback for the services they are giving for the patients

II. Objectives

a) General objective

- To assess knowledge, attitude and practice of Life Style Modification management of type 2 DM in Adama Medical College Hospital patients following diabetic clinic.

b) Specific objectives

To determine the knowledge, attitudes and practices of patients in relation to type 2 diabetes mellitus with reference to:

- Diet modification
- Importance of regular exercise
- To determine the demographic characteristics of type 2 diabetes mellitus patients attending diabetic clinic at Adama Hospital.

III. Method and Materials

a) Study area and period

The study was conducted in Adama Medical College Hospital (AMCH) located 99 km south east of Addis Ababa, Ethiopia, Oromiya region, East Shoa zone. It was established in 1946 by Italian Missionaries and named as HailemariamMammo memorial hospital little bit after establishment but its name was changed to Adama Referral Hospital in mean time and now it renamed AMC H by Oromiya regional state health bureau after it start to teach accelerated medicine, emergency surgery and anaesthesia nurses. The hospital gives services for about 5 million people east and southern parts of Oromia, Afar, Somali and Southern Nation Nationalities and People (SNNP).

Now the hospital has 465 different workers to give different services, of which 194 are administration workers. The other 271 workers are health professionals. There are specialist in different field(23),general practitioners(GP)36,Nurses(116),laboratory workers(20),x-ray(5), physiotherapy(2), sanitarians(2), Biomedical(1), Midwifery(16), Anaesthesia(9), Health officers(9),psychiatry nurses(3) and masters in different fields.

AMCH has different departments in it: Outpatient department (OPD) team case, internal medicine, dermatology, paediatrics, gynaecology/obstetric, surgery, dentistry, psychiatry, physiotherapy, ophthalmology, hospital pharmacy, anti retroviral therapy (ART) and tuberculoses (TB) Clinic. The study was conducted in OPD case team from April 1 to May 1.

The data from hospital management 2012/13 show the top 10 leading causes of outpatient visit were: Trauma, Pneumonia, Dyspepsia, Acute Upper Respiratory Tract Infection, Other or unspecified diseases of the eye and Adnexa, Urinary tract infection, Diarrhoea (non-bloody), Dental and gum diseases, Acute febrile Illness and Helmenthiasis. But the top 10 causes of admissions were: Other delivery , Trauma (Injury, Fracture etc), Pneumonia, AIDS and related diseases , Appendicitis ,Medical Abortion without complication (safe abortion), Other or unspecified obstetrics condition, Diarrhoea with dehydration, Diabetes mellitus and Severe acute malnutrition.DM

© 2014 Global Journals Inc. (US)

ranked 19th from outpatient visit but 9th causes of hospital admission[24].

b) Study design

This study was a descriptive cross-sectional study

IV. Populations

a) Source population

All diabetic patients attending the diabetic clinic follow up were used as source population.

b) Study population

The study population consisted of all type 2 diabetes mellitus patients, aged 30 years and above, attending Adama Hospital for regular follow-ups from the 1st April 2014to the 1st May 2014. Age of 30 years was chosen as the cut-off age. One hundred sixteen (116) type 2 diabetes mellitus patients attended the diabetic clinic of AMCH during this period.

All diabetic patients who visited the follow up clinic within the limit of study period were included in the study.

c) Sample size and Sampling technique

The size of study population was limited by the number of diabetic patients visiting the clinic during the study. All diabetic patients those visited the hospital during study period were included depending on their consent.

d) Inclusion and Exclusion criteria

i. *Inclusion criteria:* Type 2 diabetes mellitus patients aged 30 and above attending the diabetic clinic at Adama Medical College Hospital for their regular follow up visits and had willing to participate in the study were included in the study.
ii. *Exclusion criteria:* Patients with type 1 diabetes mellitus, gestational diabetes, other specific types of diabetes mellitus and diabetes insipidus were excluded from the study. In addition, all type 2 diabetes mellitus patients with impaired memory or cognitive functions and those younger than 30 years were also excluded.

e) Variables

i. *Dependent variables:* knowledge, attitude and practice
ii. *Independent variables:* age, religion, Sex, educational status, marital status, ethnicity, monthly income, duration of disease.

V. Data Collection Process and Collection Technique

In this study a face-to-face interview using a structured questionnaire was carried out for data collection. The data was collected by the researcher and his assistant colleagues using a structured questionnaire. The researcher trained two qualified graduating students of clinical pharmacy, proficient in the local language (Amharic &Afan Oromo) as research assistants. They assisted him throughout the data collection processes through a face-to-face interview. During and after data collection principal investigator checked the consistence and completeness of the data.

a) Data Quality Assurance

In order to assure the quality of the data the following measures was under taken:

Data was collected by three of graduating class of clinical pharmacy students

- The data collectors were taken the training to check completeness of the data during the data collection and appropriate recording.
- The body mass index (BMI) of each participant was calculated by the researcher and his assistants using the formula BMI = Weight (kg)/Height (m2) after the weight and height were measured using a calibrated beam scale with height rod graduated in centimetres and participants were classified according to the WHO international classification of adult weight.
- The principal investigator supervised all field work, check for completeness and accuracy of data collection daily.

b) Data processing and analyzing

The collected data was coded and checked for completeness. Once data coded and checked for completeness, data processing was done by SPSS version 16.1 and, presentation of the data was done by using frequency distribution, percentage and rate.

c) Ethical consideration

Ethical clearance letter was written by Ambo University department of Pharmacy after approved the proposal of the study to request AMCH for the permission. AMCH management and research office approved the letter of study and requested different hospital departments and any help during the study. The purpose of the study was explained to the respondent and data was collected after ensuring their willingness to give their response. Confidentiality of participants maintained at all time .Participants were informed that the participation were voluntary.

d) Operational definitions

LSM- Non pharmacological management such as diet modification, and exercise design to treat problem of type 2 DM patients.

Knowledge- Understanding lifestyle modification in glycemic control and management of type 2 DM patients

Attitude- A patients' positive or negative feeling towards performing the defined behaviour i.e. (LSM)

Practice - Is a previous utilization of any of LSM

© 2014 Global Journals Inc. (US)

VI. Results

a) Demographic Characteristics of Respondents

From total number of 116 type2 DM patients participated in the study 44(37.9%) male and 72(61.1%) were female. Regarding marital status of the patients; 16(13.8%) single, 80(69%) married, 10(8.6%) widow and 10(8.6%) patients were divorced. Concerning age of the clients 14(12.1%), and 52(44.83%) were in age group of 30-40, and >=61 respectively. The other 50 were in 1:1 ratio of 41-50 and 51-60 ages. Ethnicity of the patients participated in the study include Oromo (33.6%), Amhara (37.9%), Tigre (5.2%) and 27(23.3%) were other ethnic groups.

On the other hand 38(32.8%) patients do not read and write, and 24(20.7) patients read and write only. The other 28(24.1%), 23(19.8%), and 3 (2.6%) patients attended grade 1-8, grade9-12 and college respectively. When we consider employment status of the study participants; 27(23.8%) patients were unemployed (private employee) and other 11 (9.5%) were unable to do because of old age. Furthermore, 3(2.6%) were government employee, 17(14.7%) farmers, 34(29.3%) house wife and other 22(19%) had different jobs. In another way, about 41(35.3%) patients had monthly income less than 500 birr and 60(51.71%) patients had monthly income of 501-1500 birr. Only 14(12.1%) and 1(.9%) patients had monthly income of 1500-2500 and 2500-3000 birr per month respectively.

Table1. Distributions of socio demographic characteristics of diabetic patients at diabetic follow up clinic in AMCH in April, 2014

Socio demographic characteristics or variables		Number	Percent
Sex	Male	44	37.9
	Female	72	62.1
Age	30-40	14	12.1
	41-50	25	21.55
	51-60	25	21.55
	=>61	52	44.83
Marital status	Single	16	13.8
	Married	80	69
	Widow	10	8.6
	Divorced	10	8.6
Religion	Muslim	25	21.6
	Orthodox	65	56
	Protestant	18	15.5
	Other	8	6.9
Ethnicity	Oromo	39	33.6
	Amhara	44	37.9
	Tigre	6	5.2
	Other	27	23.3
Educational status	Not read write	38	32.8
	Read and write only	24	20.7
	Grade 1-8	28	24.1
	Grade 9-12	23	19.8
	College graduate	3	2.6
Employment status	Gov. Employee	3	2.1
	Un employed (Private employee	27	23.28
	Farmer	17	14.66
	House wife	34	29.31
	Not able to work	13	11.21
	Other	22	18.99
Monthly income in birr	<500 birr	41	35.3
	501-1500	60	51.7
	1501-2500	14	12.1
	2501-3000	1	0.9

b) Anthropometric Characteristics of Respondents

Among the study population only 5 (4.3%) patients were obese class I .The other 59(50.9%) patients and 52(44.8%) were in the range of normal weight and overweight respectively.

© 2014 Global Journals Inc. (US)




Figure1 : Distribution of type2 DM patients by their body weight in AMCH in April, 2014

c) Anthropometric Characteristics of Respondents

Regarding diabetic education such as attending group classes or having meeting with health professional half of the patients didn't get the opportunities while the other half got the chance one or more till now.

On the other hand, majority of the patients 59(50.9%) were 1-5 years with the disease followed by 6-10 years with the disease which were 42(36.2%).only one patient with disease for 25 years.




*Figure 2 :*Duration of disease of type 2 DM patients in AMCH in April, 2014.

d) Sources of life style modification

Majority of the patients 68(58.62%) following the diabetic clinic get the information from health personnel followed by those who get information from more than one source 25(21.55%).From the whole participants 12 patients reported that they have no information about LSM management of diabetes. See table 2 below

© 2014 Global Journals Inc. (US)

Table 2 : Source of LSM Management of Type2 Dm Patients in AMCH in April,2014

Source of information	Frequency	Percent	Valid Percent
NO INFORMATION	12	10.3	10.3
HEALTH PERSONNEL	68	58.6	58.6
MEDIA	4	3.4	3.4
RELATIVE	2	1.7	1.7
FRIENDS	3	2.6	2.6
MORE THAN 1 OPTION	25	21.6	21.6
OTHER	2	1.7	1.7
Total	116	100.0	100.0

e) Types of treatment

When this study conducted; almost all patients were taking more than one types of treatment modalities. All patients attending the clinic were at least on one type of pharmacologic treatment and in parallel practice LSM partially or completely. Only 2 patients were taking one type of treatment modalities.




Figure 3 : Frequency distribution of type of treatment of diabetes patients in AMCH in April; 2014

f) Dietary habit

Regarding selected dietary practices of cigarette smoking, alcohol consumption, fruit and vegetable, sweet sugar, buttery and fatty meal; almost all patients do not smoke and consume alcohol always while 67(57.8%) patients never take sweaty food, 71 patients never take buttery and fatty meal, while only 3(2.6) patients never take fruit and vegetable because of lack of resource. About 54(46.6%) and 37(30.21%) patients take eggs usually and never respectively. Surprisingly only one patients use herbal medicine to control her diabetic symptom with her medication.

Table 3 : Selected Dietary Practice of Type2 Dm Patients at AMCH in April, 2014

Type of diet	Always	Some time	Occasionally	Never
Smoke	0	2	6	108
Alcohol	0	6	17	93
Chat	4	4	3	105
Fruit &vegetable	30	52	31	3
Sugar	2	13	34	67
Egg	0	27	54	35
Buttery and fatty meal	0	4	41	71

© 2014 Global Journals Inc. (US)

g) Physical Activity

Regarding physical activity, 77(66.4%) patients exercise on regular programme or physical active and 39(33.6%) do not exercise on regular programme and were not physically active due to difference reasons. Of those who exercise regularly 19(16.4%) exercise less than 15 minute perday,21(18.1%) 15-30minute per day,14(12.1%) patients 30 minute per day ,11(9.5%) 30-45 minute per day ,2(1.7%) 46-1hour per day and only 8(6.9%) patients exercise for greater than 1 hour per day.

Table 4 : Frequency distribution of respondents with regard to physical activity

Variable	Number	Frequency (%)
Exercising regularly		
Yes	77	66.4
No	39	33.6
Total	116	100
If yes, how often		
Less than 15 min/day	21	18.1
Less than 30 min/day or less than 150 min/week	40	34.5
More than 30 min/day or more than 150 min/week	21	18.1

h) Knowledge Assessment of Respondents

Concerning knowledge of the patients towards LSM management of diabetic; majority of the patients were knowledgeable which accounts 90(77.59%). Of the left 26 patients, 13(11.21%) fairly knowledgeable and the other 13(11.21%) were poorly knowledgeable. Figure1 below summarize status of the knowledge of the patients.




Figure 4 : Distribution of respondents according to their knowledge regarding the benefit of life style modification at diabetic follow up clinic AMCH April, 2014

i) Attitude Assessment of Respondents

Majority of the patients 95(81.89%) had positive (good) attitude toward s LSM management of diabetes. No patient had negative attitude to LSM but, 21(18.1%) patients had neutral attitude.

Table 5 : attitude status of the diabetic patients to ward LSM in AMCH April, 2014.

Attitude level	Number	Percent %
Positive	95	81.9
Neutral	21	18.1
Negative	0	0

© 2014 Global Journals Inc. (US)

Regarding practice of LSM, almost half of the patients 57(49.1%) had good practice. The other 39(33.62%) and 20(17.24%) had low and average practice respectively. Table 3 below summarize the results.

Table 6 .Frequency distribution of diabetic patients by their practice of LSM management of diabetes AMCH April, 2014.

Level practice	Frequency	Percent
Good practice	57	49.13
Average practice	20	17.24
Low practice	39	33.62
Total	116	99.99

j) Correlation between Knowledge, Attitudes and Practices

Table 7 : Correlations between knowledge, attitude and practice level regarding lifestyle modifications

		knowledge of clients	level of attitude	level of practice
knowledge of clients	Pearson Correlation	1	.098	.184*
	Sig. (2-tailed)		.293	.048
	N	116	116	116
level of attitude	Pearson Correlation	.098	1	.517**
	Sig. (2-tailed)	.293		.000
	N	116	116	116
level of practice	Pearson Correlation	.184*	.517**	1
	Sig. (2-tailed)	.048	.000	
	N	116	116	116

*. Correlation is significant at the 0.05 level (2-tailed).

**. Correlation is significant at the 0.01 level (2-tailed).

Table7: shows the correlation of level of knowledge, attitude and practice towards LSM. It shows that there is positive Pearson correlation between level of knowledge and attitude(r=0.098, p=0.293), positive Pearson correlation of 0.184(p=0.048) between knowledge and practice, and positive Pearson correlation of 0.517(p=0.000) between level of attitude and practice level.

VII. Discussion

a) Demographic Characteristics of Respondents

There were more females (62.1%) than males (37.1%) participated in this study, a reflection of the gender ratio attendance of patients in the diabetic clinic at AHMC hospital. More recent reports from developing countries have found that DM and its risk factors are more common in women This finding is in keeping with the results from a study conducted in South Africa at Mamelodi Hospital in which 81.1 % were female and 18.9 % were male [25].

Majority of respondents in this study came from the age groups 41-50 years, 51-60 years and >61 years with 21.55%, 21.55% and 44.83% of respondents respectively, which add up to 87.93 % of respondents.. This is reflective of the fact that the ethnology of type 2 diabetes mellitus usually at old age [3, 17, and 25].

In this study respondents with no formal education consists (53.5%) and only (2.6%) respondents with higher education. This indicates that most respondents had little or no education. This result may be the direct consequence of scarcity of higher education system in Ethiopia in the past [26]. Additionally the results of this study reported less than half (45.7%) of the participants got diabetic education such as attending meeting with health professionals. This confirmed with the study conducted in 2011 at Jimma University Specialised Hospital which reported as there was no attention given to diabetic education in Ethiopia [20].

Majority of respondents in this study had income between 501 birr and 1500 birr (51.7%) followed by respondents in the less than 500 birr income (35.3%). This low income amongst majority of respondents could limit their accessibility and affordability of a well-balanced diet and healthy food and it was considered as the main factors (barrier) to their practice of life style modification especially diet modification.

This finding was in keeping with the results from a cross sectional study of Adherence to Diabetes Self-Management Practices among Type Ii Diabetic Patients in Ethiopia; in which majority of the study participants 139 (43%) had very low monthly income [27].

© 2014 Global Journals Inc. (US)

Majority of respondents (50.9%) had normal weight, followed by 44.8% with overweight and only 4.3% had class I obesity. In the study class 2 obesity and class 3 (morbid obesity) were not found. This study had just demonstrated that lack of physical activities and poor dieting habit among respondents, seem to contribute to the development of type 2 DM rather than obesity. But about 45% of the patients in the study were overweight which increases the risk of obesity.

This finding is in contrast with many studies done on this area in which obesity was common in the representative sample of type 2 diabetes patients attending a diabetes clinic [28].

b) Knowledge Assessment of Respondents

In this study (77.59%) of respondent had adequate knowledge, (11.21%) of respondents had fair (average) knowledge and 11.21% of respondents had poor knowledge of the benefits of exercise, and healthy diet. Large numbers of the participants (68%) got information from health professionals may have contributed to this result. This relatively revealed similar result with study done in Nigeria at Kaduna in the year 2012 on 347 patients; 230 non diabetic and 117 diabetic patients' .The study recorded 56.4% score of knowledge for diabetic participants [23].

In contrast to this finding, IKOMBELE found in his study that no respondent had good knowledge and 92.6% of respondents had poor knowledge of the benefits of exercise, weight loss and healthy diet [25].

c) Attitude Assessment of Respondents

The majority of respondents (81.89%) had positive attitude towards lifestyle modifications, followed by 18.1% of respondents who had neutral attitude, and no respondents had a negative attitude. This revealed relatively similar results with study conducted on 100 patients attending diabetic clinic at kinikkashitan Seri Manjung which recorded 99% of patients answered greater than or equal to 50% of attitude question [29].
This finding is similar to those of studies done in South Africa at Mamelodi Hospital in which the majority of respondents (92.7% and 51.6% respectively) had positive attitude towards lifestyle modifications [25].

d) Practice Assessment of Respondents

The proportion of respondents with good practice (49.1%) and those with average practice (17.24%) and poor practice (33.62%).About one third, patients had poor practice of LSM and the result was not satisfactory as that of knowledge and attitude. That could be due to majority of the patients had limited resources and low income which limit their affordability for a well balanced dieting and necessary equipment to exercise. This result was similar with study conducted in Qatar and Omani which reported 49.5% of the respondents were not exercise regularly and 48% of the participants were not practices recommended diet [21] and less than 40% exercise regularly and only 56% of the patients were adhere to recommended diet respectively [19].

Regarding consumption of alcohol ,chat and smoking the results of this study is promising as 93.1%,90.51% and 80.2% patients never smoke, chew chat and take any type of alcohol respectively. This result is similar with study conducted in Western Nigeria in 2012 which reported all study patients neither consume alcohol nor smoke cigarette [22]. The other similar study conducted in Omani in the year 2013 also reported out of 106 study patients only 10.6% were smoker.

Additionally the result of this study showed as only one patient use herbal medicine. Surprisingly study in Omani also reported only 2 patient use herbal medicine regularly [17].

Correlation between Knowledge, Attitudes and Practices

There was a weak, non-significant positive correlation ($r = 0.098$, $p = 0.293$) between knowledge level and practice level of respondents. This means that being knowledgeable did not necessarily willingness to observe healthy lifestyle habits. In other way, there was a significant positive correlation ($r = 0.184$, $p = 0.048$) between the knowledge level and the practice level of respondents in this study. This means that the better respondents were knowledgeable, the better they were practice healthy lifestyle.

Finally, there was a very significant positive correlation ($r = 0.517$, $p = 0.000$) between attitude level and practice level. This means the better the patients had positive attitude toward LSM, the better they were practices healthy life style modification.

The result found in this study were opposite to the study conducted in South Africa at Mamelodi Hospital which reported significant positive correlation ($r = 0.171$, $p = 0.012$) between knowledge and attitude level, and weak positive non significant correlation ($r = 0.037$, $p = 0.587$) between attitude and practice level. In addition that study also reported weak positive non significant correlation($r = 0.102$, $p = 132$) between attitude and practice level [25].

VIII. Conclusionand Recommendation

a) Conclusion

The discussion on the findings of this study shows that the knowledge and attitude levels of lifestyle modifications among type 2 diabetes mellitus patients attending Adama Medical College Hospital were generally high. However practice of the patients regarding LSM still not sufficient as more than half of the patients had no good practices. The study also found out that there was significant positive correlation ($r = 0.0184$, $p = 0.048$) between knowledge and practice, very significant positive correlation ($r = 0.517$, $p = 0.000$) between attitude and practice and a weak non-significant correlation($r = 0.098$, $p = 0.293$) between knowledge and attitude.

© 2014 Global Journals Inc. (US)

b) Recommendation

Lifestyle modification has important roles in prevention and management of chronic diseases like type 2 DM patients. But its prevalence is increasing worldwide at an alarming rate especially in developing countries due to different factors like sedentary life style (westernization), and deficits in the knowledge and practice of LSM.

Based on these facts and on our research findings, it was recommended that:

- ✓ Health education about life style modification (importance of exercise, physical exercise and weight loss) to the general society should be implemented by the responsible body.
- ✓ Medical nutrition intervention program should be implemented with a multidisciplinary team (Doctor, dietician, pharmacists…) and all stake holders (health institution, MOH, diabetic association other responsible nongovernmental organization) should work cooperatively on this issues
- ✓ Empower and train Adama hospital healthcare workers about this issue in order to promote behavioural change and adoption of healthy lifestyle practices by patients.
- ✓ Further research should be done on this area

IX. Acknowledgements

We are very grateful to our college staff members for unreserved guidance and constructive suggestions and comments from the stage of proposal development to this end. We would like to thank Ambo University for supporting the budget which required for this research. Finally our deepest gratitude goes to Adama Hospital Medical College staff workers who help and allow us in conducting this research.

X. Acronyms and Abbreviations

AHMC = Adama Hospital Medical College

ART = Anti retroviral therapy

BMI = Body mass index

DM = diabetes mellitus

IDF = international diabetic federation

KAP = Knowledge, attitude and practice

LSM = Life style modification

OPD = outpatient department

OHG = Oral Hypoglycaemic

USA = United States of America

WHO = world health organization

References Références Referencias

1. Kumar V, TripathiKM, Chauhan K.P, Singh K.P. Different non-pharmacological approaches for management of type 2 diabetes. joudibet. 2013; 1:6
2. World Health Organization. Screening for Type2 Diabetes. Reports of World Health Organization and International Diabetic Federation meeting. Geneva: World Health Organization, 2003.
3. Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, Jameson JL, et al., editors. Harrison's principles of internal medicine. 18th e. New York: McGraw Hill; 2012.
4. Jean-Pierre Hallé. The Management and Treatment of Type 2 Diabetes. Canadian Journal of CME .June 2001.p65-77.
5. MegersaYc,et al. Prevalence of undiagnosed DM and its risk factors in selected institutions of at Bishoftu town. journal of diabetes,2013.s12:068
6. Shills ME. Modern nutrition in health and disease, 8th e. USA Waiver,1995. pp-722-724
7. ICMR Guidelines for Management of Type 2 Diabetes. Non-Pharmacological Management of Diabetes. ICMR GUIDE LINES, 2005.
8. Kisokanth G, Prathapan S, Indrakumar3 J, Joseph J. Factors influencing self-management of Diabetes Mellitus; a review article. J diabet Oct,2013;3:1
9. Badrudin N, Basit A, Hydrie MZI, Hakeem R. Knowledge, Attitude and Practices of patient visiting diabetes care unit. Pak J Nutrition. 2002;1:99-102
10. K. G. M. M. Alberti, P. Zimmet, J. Shaw. International Diabetes Federation: a consensus on Type 2 diabetes prevention. Diabetes UK,2007; 24:451–463
11. Binu.M.G, Manoj.P, Bhuvaneszwari.S. Non-Pharmacological management of type 2 DM; Where do we stand?. International Journal of Clinical Cases and Investigations 2011. Volume 2 (Issue 6), 27:34, 6th July, 2011.p27-34
12. Seyoum B, Abdulkadir J, Gebregziabher F, Alemayehu B. Analysis of diabetic patients admitted to TikurAnbessa Hospital over eight years period. Ethiop J Health Dev. 1999; 13:9-13.
13. Gning SB, Thiam M, Fall F, Ba-Fall K, Mbaye PS, Fourcade L: Diabetes mellitus in sub-Saharan Africa: epidemiological aspects and management issues. Med Trop (Mars) 2007, 67(6):607–611.
14. Feleke F, Enquselassie F. An assessment of the health care system for diabetes in Addis Ababa. Ethiop.J.Health Dev. 2005; 19(3) P203-210.
15. Worku D. et al. Patterns of Diabetic Complications at Jimma University Specialized Hospital, Southwest Ethiopia. Ethiop J Health Sci. March 2010, Vol. 20, No. 1,P33-40
16. Abebe et al. Diabetes mellitus in North West Ethiopia: a community based study. BMC Public Health 2014,14:97
17. Al Bimani, Z.S. et al., Evaluation of T2DM related knowledge and practices of Omani patients. Saud PharmJ(2014),http://dx.doi.org/10.1016/j.jsps.2013.12.006C.
18. ROSAL, M. BENJAMIN, S. PEKOW, C. LEMON. Opportunities and Challenges for Diabetes

© 2014 Global Journals Inc. (US)

Prevention at Two Community Health Centres. DIABETES CARE. 2008; 31: 2, P247-254
19. Wk. Maina et al. Knowledge, attitude and practices related to diabetes among community members in four provinces in Kenya: a cross-sectional study.panAf j med.October 2010.
20. Gudina et al. Assessment of quality of care given to diabetic patients at Jimma University Specialized Hospital diabetes follow-up clinic, Jimma, Ethiopia. BMC Endocrine Disorders 2011; 11:19.
21. Abyad. Knowledge and Practice of Type 2Diabetic Patients Attending Primary Health Care in Qatar. J fam med.2011,9(4)
22. Oguntibeju OO, Odunaiya N, Oladipo B, Truter EJ. Health Behaviour and Quality of Life of Patients with Type 2 Diabetes Attending Selected Hospitals in South Western Nigeria. West Indian Med J. 2012; 61 (6): 619-626.
23. HamoudNehad , Al Ayoubi Dh, VanamaJ ,Yahaya H, Usman FH. Assessment of Knowledge and Awareness of Diabetic and Non-Diabetic Population towards Diabetes Mellitus in Kaduna, Nigeria. J AdvSci Res, 2012;3(3): 46-50
24. Adama Medical college Hospital Management Office
25. Ikombele JB. Knowledge, Attitudes and Practices Regarding Lifestyle Modifications among Type 2 Diabetic Patients Attending Mamelodi Hospital, Pretoria, Gauteng. University of Limpopo; 2011.
26. Ashcroft K. Analysis and discussion of curriculum, resources and organizational issues. Ethio J High Edu, 2005.
27. Berhe KK, KahsayBA,Gebru BH. Adherence to Diabetes Self-Management Practices among Type Ii Diabetic Patients in Ethiopia; A Cross Sectional Study. Green J Med Sci .2013;3(6):211-221
28. Shivapaksh et al. Body mass index waist curcumferunce in Type 2 diabets mellitus patients attending a diabetes clinic. Int J Biol Med Res. 2011; 2(3):636-638.
29. The Uk's expert provider of custom essays. Review literature on lifestyle modification by diabetes sufferers.http://www.ukessays.com/essays/nursing/reviewliteratureonlifestylemodifictionbydiabetessufferersnursingessay.php.UK'sessays,1999.

© 2014 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2014

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSM" title to individuals.The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSM" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall,Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.




© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSM, you will be given a renowned, secure and free professional email addres with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.



The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of you credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.



The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.

MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSM" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.

MARSM accrediting is an honor. It authenticates your research activities. Afterbecoming MARSM,you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.



MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you willbe given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook



We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.





Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

AUXILIARY MEMBERSHIPS

Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:



Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

INDEX

A

Acclimatized · 3
Amphiphiles · 13, 24
Analykjena · 29
Antiarrhythmics · 13, 23
Antiulcerogenic · 9

B

Benzothiazole · 24
Brownlowia · 38, 39
Butylatedhydroxy · 36

C

Carrageenan · 20, 25, 27, 28, 30, 31, 34, 36
Cotherapies · 13
Cyclooxygenase · 35, 39
Czochara · 23, 24

D

Darmstadt, · 27
Dendrobium · 38
Deoxyribo · 51
Diterpenoid, · 25
Dyshomeo · 13

E

Elucidated · 15
Epicatechin · 25, 27, 30, 34, 39
Equimolar · 15
Ethanolic · 15, 32, 34, 38, 39

F

Ferricyanide · 27, 29, 36
Fluoroben · 24
Fuzlat · 2

G

Glabinclamide, · 46, 49

H

Hailemariam · 57
Helmenthiasis. · 57
Heterocyclic · 24
Histopathological · 5, 6, 8
Homeostasis · 23, 24

I

Increasingburde · 52

K

Kinikkashitan · 64
Kruskall · 5, 6

L

Ledingham · 10

M

Methanolic · 10, 39
Mitochondrial · 24
Molyb · 27

N

Norlimonoids · 25

O

Oedematogenic · 36
Omeprazole · 3, 5, 6

P

Pcoumeric · 25
Phosphomolybdenum · 38
Pimaradiene · 25

Q

Qarah · 2

R

Rapeutically · 20

S

Scitamineae · 47
Sigurdardottor · 52
Submucosa · 1
Sulfonamidyl · 18

T

Tainter · 3, 10
Tibiofasial · 17
Tirucallane · 25
Trichodesma · 38
Triterpenoids, · 25
Tyszka · 13, 23, 24

U

Ulcerogenic · 5, 10

Z

Zothiazole · 24

save our planet



Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896

© Global Journals