



GLOBAL JOURNAL OF MEDICAL RESEARCH  
PHARMA, DRUG DISCOVERY, TOXICOLOGY AND MEDICINE  
Volume 13 Issue 4 Version 1.0 Year 2013  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

## Effect of Oral Administration of Chloramphenicol on Hematological Profile of Male Charles Foster Rats

By P. Shukla & R. K. Singh

*CSIR- Central Drug Research Institute, India*

**Abstract** - For a given organism, relevant information about the internal environment can be easily accessed by its hematological profile. Chloramphenicol being a potent broad spectrum antibiotic is used readily in eyed drop formulations and is also in food industry. In the present study, varying doses (750, 1500 and 2250 mg/kg B.Wt) of Chloramphenicol (CAP) was administered orally as single daily dosage for 24 days to Male Charles Foster rats, to assess the hematological changes associated with oral exposure to the drug. The results showed a significant ( $p < 0.05$ ) dose dependent decrease in Red Blood Cells (RBC) count, Hemoglobin (Hgb), Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin Concentration (MCHC) and increase in Hematocrit (Hct), White Blood Cells (WBC) and Platelets compared to the initial blood profile. The results recorded in this present study suggested that exposure to CAP results in Hematotoxicity. Hence, the potential of CAP to cause hematotoxicity is reported in the study.

**Keywords** : CAP, hematotoxicity, blood cells, CF rats.

**GJMR-B Classification** : NLMC Code: WB 350



*Strictly as per the compliance and regulations of:*



# Effect of Oral Administration of Chloramphenicol on Hematological Profile of Male Charles Foster Rats

P. Shukla<sup>α</sup> & R. K. Singh<sup>σ</sup>

**Abstract** - For a given organism, relevant information about the internal environment can be easily accessed by its hematological profile. Chloramphenicol being a potent broad spectrum antibiotic is used readily in eyed drop formulations and is also in food industry. In the present study, varying doses (750, 1500 and 2250 mg/kg B.Wt) of Chloramphenicol (CAP) were administered orally as single daily dosage for 24 days to Male Charles Foster rats, to assess the hematological changes associated with oral exposure to the drug. The results showed a significant ( $p < 0.05$ ) dose dependent decrease in Red Blood Cells (RBC) count, Hemoglobin (Hgb), Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin Concentration (MCHC) and increase in Hematocrit (Hct), White Blood Cells (WBC) and Platelets compared to the initial blood profile. The results recorded in this present study suggested that exposure to CAP results in Hematotoxicity. Hence, the potential of CAP to cause hematotoxicity is reported in the study.

**Keywords** : CAP, hematotoxicity, blood cells, CF rats.

## I. INTRODUCTION

Chloramphenicol (CAP) is a broad spectrum antibiotic, was first quarantined from bacterium *Streptomyces venezualae* in the year 1947. It was available by the trade name of Chloromycetin by Parke Davis & Co. It was prescribed in mass in 1948 in USA following an outburst of enteric fever. In 1949 it was cleared from Federal Food and Drug, since then it has been used and worked upon extensively being a potent inhibitor of protein synthesis (E. Cundliffe and K. McQuillen, 1967).

Some studies suggest the use of CAP in food. Although, most countries have banned CAP from animal food production, still traces of it have been detected in shrimp and other aquaculture products. According to regulations promulgated in 1980's and 1990's, use of CAP in food was banned and countries have established a zero tolerance policy. In Japan, zero tolerance thresholds for CAP is 50 ppb which in USA is 5 ppb. Meat and offal from treated animals contained CAP and its non – genotoxic metabolites (G. Milhaud, 1993).

Even being a potent antibiotic with a broad range of spectrum, the use of CAP is limited due to its

association with aplastic anaemia (AA) (M. L. Rich, et al., 1950) and bone marrow suppression (C. E. Amberkar, et al., 2000). AA is a rare, dose independent, irreversible, idiosyncratic, manifestation of CAP which in most cases is seen years after the treatment (A. A. Younis, 1989 (a) and is fatal (A. A. Turton, et al., 2002) risk of developing AA after CAP administration is 1:30000 to 1:5000015 (C. H. Li, et al., 2010). Only orally administered CAP leads to AA (R. Holt, 1967; R. A. Gleckman, 1975). This has made the CAP to be prescribed parenterally by many physicians. It is not known whether this lowers the incidence of AA or not but the risk is obviously lowered. Other than oral and parenterally absorbed CAP, it is also used as ophthalmic preparations where AA is also very rare (R. L. Rosenthal and A. Blackman, 1965; G. Carpenter, 1975; S. M. Abrams, et al., 1980)

Blood or hematological parameters are probably the more rapid and detectable variations under stress and are fuel in assessing different health conditions (V. Hymavathi and L. M. Rao, 2000). Hence, the significance of hematological parameters in clinical and experimental studies in life sciences cannot be overemphasized. Particularly, literature reports have proved that the alterations in the hematological parameters, from normal state/levels, may be used as valuable indicators of disease, or stress in different animal species (A. K. Solanke and V. Singh, 2000; B. K. Das and S. C. Mukherjee, 2003; L. H. Jee, et al., 2005; M. F. Rahman and M. K. Siddiqui, 2006; F. E. Uboh, et al., 2005).

Literature reports that hematological profile of different species of animals may be influenced adversely by phenylhydrazine (F. S. Sanni, et al., 2005; J. Berger 2005; S. K. Jain and D. Subrahmanyam, 1978), some antiretroviral drugs (A. A. A. Kayode, et al., 2011), Paclitaxel (J. A. Juaristi, et al., 2001), Carbamazepine (S. Thakur, et al., 2012), Doxorubicin (D. A. Eppstein, et al., 1989), Tetrachloroethylene (A. M. Emara et al., 2010), Phenacetin (C. B. Jensen and D. J. Jollow 1991) and Benzene (A. Beamonte et al., 2005; R. Synder and C. C. Hedli, 1996). Plants also have been shown to have ameliorative potential in reference to drug induced hematotoxicity (F. S. Sanni et al., 2005; E. E. Edet et al., 2011; E. V. Ikpeme et al., 2011; S. O. Kolawole et al., 2011; G. Prasad and G. L. Priyanka, 2011). Different

extracts (methanolic, ethanolic, water, chloroform, hexane) of some plants namely, *Hibiscus cannabinus* (G. A. Agbor, et al., 2005), *Brillantaisia nitens* (P. A. Akah, et al., 2009(a), 2010(b)), *Hibiscus sabdariffa* (A. Ologundudu, et al., 2010), *Zingiber officinale* (A. M. M. Attia, et al., 2013), *Ocimum basilicum* (S. Saha. et al., 2012) and *Ocimum gratissimum* (A. W. Obianime, et al., 2011) have been reported to express a positive impact on the hematological profile of several animal species. Assessment of hematological parameters can therefore be useful in determining the extent of deleterious effects of foreign substances on the blood parameters of an animal. The present investigation was therefore aimed at assessing the effect of Chloramphenicol on the hematological profile in Charles Foster male rats.

## II. MATERIALS AND METHODS

### a) Administration of Material

The chloramphenicol Capsules IP manufactured by Piramal Health Care Limited (Batch No-9BE012) were used for the study. Freshly prepared chloramphenicol suspension was administered orally by cannula for 24 days.

### b) Animals

Albino rats of *Charles Foster* strain were used in the study. IAEC approval number was taken from the Institutional Animal House Facility which is affiliated to and works under the guidelines of CPCSEA (No. 36/11/Toxicol/IAEC). Rats weighed between 120- 150 grams and were housed in polypropylene, autoclavable cages (dimensions: 43x27x15 cm) with steel wire-mesh lid having provisions for attaching water bottle and for keeping food pellets. Animals had continuous access to food and water during the entire period of experimentation. They were examined routinely for their body weights and hematological parameters.

### c) Experimental Design

20 rats showing evidences of good health were selected on the basis of findings of their initial health check-up and body weight recordings. They were

randomly assigned to four treatment groups, each group consisting of five male animals and one group comprising of an equal number of animals served as control.

Group I: Control (Distilled water)

Group II: Low Dose (750 mg/ kg B.wt CAP)

Group III: Mid Dose (1500 mg/ kg B.wt CAP)

Group IV: High Dose (2250 mg/ kg B.wt CAP)

### d) Hematological Investigations

Blood collected from the caudal vein of experimental animals was assessed for all hematological parameters RBC (Red Blood Cell), Hgb (Hemoglobin), MCV (Mean Corpuscular Volume), MCHC (Mean Corpuscular Hemoglobin Concentration), Hematocrit (Hct), White Blood Cells (WBC) and Platelets. Blood analysis was performed at regular time intervals using fully automatic hematology analyzer MS-9. (Make/Model of Analyzer: MS-9 (Mellet Schloesing). Standard chemicals and reagents supplied by company were used.

### e) Statistical Analysis

All data was analyzed by applying One way ANOVA with the *p* value limits of 0.05. Software used for the purpose was PRISM.

## III. RESULTS

The result of this study, on the effect of oral dosing of CAP on the hematological parameters in rats is presented in Table 1 and 2. The results showed that the hemoglobin (Hgb), Red Blood Cells (RBC) count, Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin Concentration (MCHC) obtained for rats administered with CAP orally were significantly ( $p < 0.05$ ) lower in a dose-dependent pattern, compared to the control (Tables 1 and 2). On the contrary, the total White Blood Cells (WBC), platelets and Hematocrit (Hct) levels obtained for rats administered with CAP orally following the same pattern, were significantly ( $p < 0.05$ ) higher, compared to the control.

Table 1 : Initial Hematological Profile of CF rats

Gp	Treatment	Hgb	RBC	Hct	MCV	MCHC	WBC	Platelets
Gp. I	Control, D.W.	11.88±0.51	7.09±0.54	44.44±2.68	62.80±2.29	26.70±0.67	7.02±1.15	530.00±112.84
Gp. II	750 mg/g b.wt	12.30±0.51	6.97±0.69	46.22±5.57	66.28±3.87	26.78±2.64	9.93±2.05	331.00±145.74
Gp. III	1500 mg/kg b.wt	12.92±0.26	7.81±0.75	52.18±3.44	66.96±2.88	24.74±1.13	10.70±2.79	339.00±162.06
Gp. IV	2250 mg/kg b.wt	12.52±0.63	7.65±0.68	48.82±3.99	63.88±1.65	23.98±5.35	10.40±3.60	391.00±52.72

Data are presented as Mean±S.D., n=5,  $p < 0.05$  compared to control.

Table 2 : Hematological Profile of CF rats after oral administration of Chloramphenicol for 24 days

Gp	Treatment	Hgb	RBC	Hct	MCV	MCHC	WBC	Platelets
Gp. I	Control, D.W.	10.42±0.30	6.96±0.38	50.48±2.53	56.42±2.27	24.62±1.39	17.94±3.62	560.20±126.80
Gp. II	750 mg/kg b.wt	11.62±0.57	5.67±0.57	55.82±5.14	58.48±1.73	24.98±0.92	16.92±3.81	339.40±28.38
Gp. III	1500 mg/kg b.wt	10.12±1.29	7.20±0.75	55.28±5.14	60.12±2.37	22.96±1.63	17.58±5.27	419.40±127.18
Gp. IV	2250 mg/kg b.wt	11.26±0.40	6.24±0.75	51.98±3.73	56.32±2.24	23.64±1.11	17.12±3.34	397.40±9.63

Data are presented as Mean±S.D., n=5, p< 0.05 compared to control.

#### IV. DISCUSSION

Hematological profiles are known to provide important information about the internal environment of a given organism. The results of this present investigation showed that oral exposure to CAP caused a significant decrease in Hgb, RBC, MCV and MCHC, whereas increase in Hct, WBC and Platelets. Similar effects on hematological parameters have been reported for such other drugs as Chlorpyrifos (Y. Savithri et al., 2010), Thiodan 35 E.0 (A. K. Solanke and V. H. Singh 2000), Chloropharm (T. Fujitani et al., 2001), Endosulfan (N. Choudhary and S. C. Joshi, 2002) and Lindane (M. D. A. Baig, 2007) and Deltamethrin (S. H. Kowalczyk-Bronisz, et al., 1990). The hematotoxic condition may results from different mechanisms, including decrease in the rate of blood cells synthesis and/or increase in the rate of blood cells destruction. The observed decrease in RBC count, Hgb, MCH and MCHC may therefore, may assumed to be associated with retarded hemopoiesis, destruction and shrinkage of RBC.

Increase in total white blood cells and platelets, as well as increase in Hct, is also reported in this study. The increase in total white blood cells and lymphocyte observed in this work may be suggested to be due to stimulated lymphopoiesis and/or enhanced release of lymphocytes from lymph myeloid tissue (B. K. Das and S. C. Mukherjee, 2003). This lymphocyte response might be a direct stimulatory effect of toxic substances on lymphoid tissues. Alternatively, this response may be assumed to be associated with the drug induced tissue damage and disturbance of the non-specific immune system leading to increased production of leukocytes.

Researchers have reported that CAP induce and enhances some defects which results in damage to undifferentiated marrow stem cells (E. P. Cronkite, 1964). Other researchers suggested that certain enteric bacteria can produce a specific enzyme that degrades CAP to a toxic product (R. Holt, 1967). This was suggested by further studies, which suggests that the metabolites of CAP generated by intestinal bacteria undergo further metabolic transformations in system with *in situ* production of toxic intermediate (A. A. Yunis,

1989 (a)). In a study (A. A. Yunis, 1973 (b)) it was actually revealed that the p-nitrosulfathiazole group is responsible for CAP induced hematotoxicity by inhibiting DNA synthesis in marrow stem cells. This theory was based on the observation that thiamphenicol which is a CAP derivative, does not have a p-nitrosulfathiazole group and does not cause hematotoxicity and thus, extensively used in Europe. This theory was further supported by studies indicating CAP reduced to p-nitrosulfathiazole which is a short lived reduction intermediate and leads to helix destabilization and strand breakage (M. Irena, et al., 1983) except than being unstable these intermediates are highly toxic (P. Eyer, et al., 1984). At a concentration of 2000-4000 µg/ml CAP depressed phagocytosis and burst activity of neutrophils (M. J. Paape, et al., 1990). Other studies suggests that CAP directly induce apoptosis in hematopoietic stem cells, directly leading to hematotoxicity (C. I. Kong, et al., 2000).

#### V. CONCLUSION

In conclusion, significant adverse changes in hematological parameters are reported to be associated with exposure to CAP, in this present study. This therefore suggest that exposure to CAP may be considered to be among the risk factors for the development of anaemic condition. Hence, exposure to this drug should be minimized.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. A. A. A. Kayode, O. T. Kayode, O. A. Aroyeun and M. C. Stephen (2011) Haematologic and hepatic enzyme alterations associated with acute administration of Antiretroviral drugs. J. Pharmacol. Toxicol., 6: 293-302.
2. A. A. Turton, C. M. Andrews, A. C. Harvard, T. C. Williams (2002) Studies on haematotoxicity of chloramphenicol succinate in Dunkin Hartley guinea pig. Int. J. Exp. Pathol., 5:225 -238.
3. A. A. Yunis (1989 (a) Chloramphenicol toxicity: 25 years of research. Am. J. Med., 3 N: 44N – 48N.
4. A. A. Yunis (1973 (b) Chloramphenicol – induced bone marrow suppression. Semin. Hematol., 10: 225 – 234.



5. A. Beamonte, F. Goldfain-Blanc, N. Casadevall, D. Bazot, H. Bertheux, N. Claude (2005) A case of drug-induced hematotoxicity: from in vivo to in vitro assessment. *Comp Clin. Path.*, 14: 61–65.
6. A. K. Solanke and V.H. Singh (2000) Haematological changes in rat, *Rattus rattus* after repeated exposure to thiodan 35. *EC. Environ. Ecol.*, 18: 529-531.
7. A. M. Emara., M. M. A. El-Noor, N. A. Hassan, A. A. Wagih (2010) Immunotoxicity and hematotoxicity induced by tetrachloroethylene in egyptian dry cleaning workers. *Inhal. Toxicol.*, 22 (2): 117 – 124.
8. A. M. M. Attia, F. A. A. Ibrahim, G. M. Nabil and S. W. Aziz (2013) Antioxidant effects of whole ginger (*Zingiber officinale* Roscoe) against lead acetate-induced hematotoxicity in rats. *J. of Med. Plants Res.*, 7(17): 1108-1113.
9. A. Ologundudu, A. O. Ologundudu, O. M. Oluba, I. O. Omotuyi and F. O. Obi. (2010) Effect of *Hibiscus sabdariffa* anthocyanins on 2, 4-dinitrophenylhydrazine-induced tissue damage in rabbits. *J. Toxicol. Environ. Health*, (1): 1-6.
10. A. W. Obianime, J. S. Aprioku and C. Esomonu (2011) the effects of aqueous *Ocimum gratissimum* leaf extract on some biochemical and hematological parameters in male mice. *Asian J. Biol. Sci.*, 4: 44-52.
11. B. K. Das and S. C. Mukherjee (2003) Toxicity of cypermethrin in *Labeo rohita* fingerlings: Biochemical enzymatic and haematological consequence. *Comp. Biochem. Physiol. Toxicol. Pharmacol.*, 134: 109-121.
12. C. B. Jensen and D. J. Jollow (1991) the role of N-hydroxyphenetidine in phenacetin-induced hemolytic anemia. *Toxicol. Appl. Pharmacol.*, 111(1):1-12.
13. C. E. Ambeker, B. Cheung, J. Lee, L. C. Chan, R. Liang, C. R. Kumana (2000) Metabolism of chloramphenicol succinate in human bone marrow. *Eur. J. Clin. Pharmacol.*, 56:405 – 409.
14. C. H. Li, Y. W. Cheng, P. L. Liao (2010) Chloramphenicol causes mitochondrial stress , decreases ATP biosynthesis, induces matrix metalloproteinase – 13 expression, and solid tumor cell invasion. *Toxicol Sci.*, 116 (1): 140 – 150.
15. C. I. Kong, D. E. Holt, S. K. Ma, A. K. Lie, L. C. Chan (2000) Effects of antioxidants and a caspase inhibitor on chloramphenicol induced toxicity on human bone marrow and HL 60 cells. *Hum. Exp. Toxicol.*, 19(9):503 – 510.
16. D. A. Eppstein, C. G. Kurahara, N. A. Bruno and T. G. Terrell (1989) Prevention of Doxorubicin-induced Hematotoxicity in Mice by Interleukin 1. *Cancer Res.* 49: 3955-3960.
17. E. Cundliffe and K. McQuillen (1967) Bacterial protein synthesis: the effects of antibiotics. *J. Mol. Biol.*, 30: 137 – 146.
18. E. E. Edet, M. I. Akpanabiatu, F. E. Uboh, T. E. Edet, A. E. Eno, E. H. Itam and I. B. Umoh (2011) *Gongronema latifolium* crude leaf extract reverses alterations in haematological indices and weight-loss in diabetic rats. *J. Pharmacol. Toxicol.*, 6: 174-181.
19. E. P. Cronkite (1964) Enigmas underlying study of haemopoietic cell proliferation. *Fad. Proc.*, 23: 649 – 661.
20. E.V. Ikpeme, U. B. Ekaluo, M. E. Kooffreh and O. Udensi (2011) Phytochemistry and haematological potential of ethanol seed, leaf and pulp extracts of *Carica papaya* (Linn.). *Pak. J. Biol. Sci.*, 14: 408-411.
21. F. E. Uboh, P. E. Ebong, O. U. Eka, E. U. Eyong, M.I. Akpanabiatu (2005) Effect of inhalation exposure to kerosene and petrol fumes on some anaemia-diagnostic indices in rats. *Global J. Environ. Sci.*, 3: 59-63.
22. F. S. Sanni, S. Ibrahim, K. A. N. Esievo and S. Sanni (2005) Effect of oral administration of aqueous extract of *Khaya senegalensis* stem bark on phenylhydrazine-induced anaemia in rats. *Pak. J. Biol. Sci.* 8: 255-258.
23. G. A. Agbor, J. E. Oben, J. Y. Ngogang (2005) Haematinic activity of *Hibiscus cannabinus*. *Afr. J. Biotech.*, 4 (8): 833-837.
24. G. Carpenter (1975) Chloramphenicol eye drops and marrow aplasia [letter]. *Lancet*, 2:326 – 327.
25. G. Milhaud (1983) Metabolic study discussion on chloramphenicol. WHO report.
26. G. Prasad and G. L. Priyanka (2011) Effect of rind extract of *Garcinia gummi-gutta* on haematology and plasma biochemistry of Catfish, *Pangasianodon hypophthalmus*. *Asian J. Biochem.*, 6: 240-251.
27. J. A. Juaristi, M. V. Aguirre, R. J. Carmuega, M. Romero-Benitez, M. A. Alvarez, N. C. Brandan (2001) Hematotoxicity induced by paclitaxel: in vitro and in vivo assays during normal murine hematopoietic recovery. *Methods Find Exp Clin Pharmacol.* 23(4):161-167.
28. J. Berger (2007) Phenylhydrazine Haematotoxicity. *J. Appl. Biomed.*, 5: 125–130.
29. L. H. Jee, F. Masroor and J.C. Kang (2005) Responses of cypermethrin-induced stress in haematological parameters of Korean rockfish, *Sebastes schlegeli* (Hilgendorf). *Aquacult. Res.*, 36: 898-905.
30. M. D. A. Baig, (2007) Pesticidal residue analysis of organochlorine residues in different milk samples from Chittoor district in Andhra Pradesh, India. Final Report of UGC Minor Research Project during the Period from 2004-2006.
31. M. F. Rahman and M. K. Siddiqui (2006) Hematological and clinical chemistry changes induced by subchronic dosing of a novel

- phosphorothion ate (RPR-V) in Wistar male and female rats. *Drug Chem. Toxicol.*, 29: 95-110.
32. M. Irena, Skolimowski, R. C. Knight, D. I. Edwards (1983) Molecular basis of chloramphenicol and thiamphenicol toxicity to DNA in vitro. *J. Antimicrob. Chemother.*, 12(6): 534 – 542.
  33. M. J. Paape, S. C. Nickerson, G. Ziv (1990) In vivo effects of chloramphenicol, tetracycline, and gentamicin on bovine neutrophil function and morphologic features. *Am. J. Vet. Res.*, 51: 1055 – 1061.
  34. M. L. Rich, R. J. Ritterhoff, R. J. Hoffman (1950) A fatal case of aplastic anemia following chloramphenicol (chloromycetin) therapy. *Ann. Inter. Med.*, 33: 1459 – 1467.
  35. N. Choudhary and S. C. Joshi (2002) Effect of short term endosulfan on hematology and serum analysis of male rat. *Indian J. Toxicol.*, 9: 83-87.
  36. P. A. Akah, C. E. Okolo and A. C. Ezike (2009) the haematinic activity of the methanol leaf extract of *Brillantasia nitens* Lindau (Acanthaceae) in rats. *Afr. J. Biotech.*, 8 (10): 2389-2393.
  37. P. A. Akah, C. E. Okolo, T. C. Okoye and N. V. Offiah (2010) Aqueous extract and methanol fractions of the leaves of *Brillantasia nitens* Lindau. Reverses phenylhydrazine – induced anaemia in rats. *J. Med. Plants Res.*, 4 (3): 271 – 277.
  38. P. Eyer, E. Lierheimer, M. Schneller (1984) Reactions of nitrochloramphenicol in blood. *Biochem. Pharmacol.*, 33:2299 – 2308.
  39. R. A. Gleckman (1975) Warning-chloramphenicol may be good for your health. *Arch. Intern. Med.*, 135:1125 – 1126.
  40. R. Holt (1967) the bacterial degradation of chloramphenicol. *Lancet*, 1:1259-1260.
  41. R. L. Rosenthal and A. Blackman (1965) Bone marrow hypoplasia following use of chloramphenicol eyedrops. *J. A. M. A.*, 191: 136 – 173.
  42. R. Synder and C. C. Hedli (1996) an overview of benzene metabolism (Review). *Environ. Health Perspect.*, 104: 1165-1171.
  43. S. H. Kowalczyk – Bronisz, J. Gieldanowaki, B. Bubak (1990) Immunological profile of animals exposed to pesticide--deltamethrin. *Arch. Immunol. Ther. Exp (Warsz)*, 38 (3 - 4): 229 – 38.
  44. S. K. Jain and D. Subrahmanyam (1978) on the mechanism of phenylhydrazine-induced hemolytic anemia. *Biochem. Bioph. Res. Co.*, 82 :(4) 1320 – 1324.
  45. S. M. Abrams, T. J. Degnan, V. Vinciguerra (1980) Marrow aplasia following topical application of chloramphenicol eye ointment. *Arch. Intern. Med.*, 140:576 – 577.
  46. S. O. Kolawole, O. T. Kolawole and M. A. Akanji (2011) Effect of aqueous extract of *Kkaya senegalensis* stem bark on biochemical and haematological parameters in rats. *J. Pharmacol. Toxicol.*, 6: 602-607.
  47. S. Saha, M. K. Mukhopadhyay, P. D. Ghosh and D. Nath (2012) Effect of Methanolic Leaf Extract of *Ocimum basilicum* L. on Benzene-Induced Hematotoxicity in Mice. *Evid. Based Complement. Alternat. Med.*, doi:10.1155/2012/176385.
  48. S. Thakur, M. Eswaran, S. G. Rajalakshmi. (2012) Amelioration of carbamazepine induced oxidative stress and hematotoxicity by vitamin C. *Spatula DD*, 2(3):173-180.
  49. T. Fujitani, Y. Tada, A. T. Noguchi and M. Yoneyama (2001) Effect of chlorpropham (CIPC) on the hemopoietic system of rats. *Food Chem. Toxicol.*, 39: 253-259.
  50. V. Hymavathi and L.M. Rao (2000) Effect of sublethal concentrations of lead on the haematology and biochemical constituents of *Channa punctatus*. *Bulletin Pure Applied Sci.*, 19: 1-5.
  51. Y. Savithri, P. R. Sekhar and P. J. Doss (2010) Changes in hematological profiles of albino rats under chlorpyrifos toxicity. *Int. J. Pharma. Bio. Sci.*, 1: 1-7.

