

Impact of Socioeconomic Status on Development of Incarcerated Inguinal Hernia in Children Treated in Gezira National Center for Paediatrics Surgery (GNCPS)

Mogahid Mahmoud Mohammed Ali¹

¹ M.B.S.S university of Gezira Registrar of general surgery

Received: 5 April 2013 Accepted: 1 May 2013 Published: 15 May 2013

Abstract

Background: Inguinal hernia is common surgical problem in paediatrics and incarceration is the most common subsequent complication if left untreated. The fact that there is effect of the socioeconomic factors on development of incarceration in form of delayed presentation. Objectives: To determine the relation between socioeconomic status and development of incarcerated inguinal hernia among children presented to (GNCPS). Results: 43 case presented with incarcerated inguinal hernia within the study period those <5Yrs were 40 (93,0

Index terms— incarcerated inguinal hernia, manual reduction, socioeconomic effects.

1 Introduction

studies on incarceration and its socioeconomic implications have yielded interesting results. The underdeveloped countries have increase in the risk of incarceration by three-folds, while reduction has been made at a low rate of 34% because these patients attended the hospital late. (1)(2) Classically, the child would present with painful and persistent inguinal swelling. There may be redness and tenderness on examination. Abdominal distension and signs of intestinal obstruction may set in if diagnosis is not made early. Unless there is clear peritonitis or bowel compromise, attempt should be made to manually reduce the incarcerated hernias by using a technique called taxis. In this technique, with the infant relaxed (using sedation if necessary), gentle inferolateral pressure is applied to the incarcerated hernia with some pressure from above to straighten the canal. Approximately 80% of incarcerated inguinal hernias can be reduced using this technique (3) . Because of the high rate of early recurrent incarceration, it is recommended to admit these children and then do the surgery 24 to 48 hours later after the edema has subsided. Any child with an incarcerated hernia that cannot be reduced must undergo immediate operative repair. It is important not to reduce the hernia under anesthesia before the incision in order to inspect the incarcerated bowel for evidence of strangulation. In girls, an incarcerated ovary may be present in the hernia sac. If reduction is unsuccessful, there is a risk of vascular compromise from ovarian torsion occurring in as many as 33% of cases. Immediate repair can prevent this complication and is recommended by multiple authorities (3,4,5) .

2 II.

3 Patient & methods

The author will use non-probability-based, purposeful-convenient sampling. All subjects presented to the National center of paediatrics surgery with incarcerated inguinal hernia. Some candidates were eliminated, and only those who fulfilled the inclusion criteria were considered as the final study sample.

Patient presented with incarcerated inguinal hernia with obstructive symptoms will be admitted, kept fasting, fluid therapy, NG. tube, catheter. Investigations were request in form of CBC, UG, RFT and electrolytes they candidate for General anesthesia with halothane. all candidates will receive injectable antibiotics. supine position,

42 scrubbing with Antiseptic solution. incision will be transverse inguinal. Exploration of the contents will be done
43 then managed accordingly. hernia sac will be excised and closure of the wound. post operative follow up in form
44 of recording the vital signs, injectable antibiotics intravenous fluids and paracetamol suppositories as form of
45 analgesia. They discharged to be seen in referred clinic after a week for assessment of general condition and the
46 wound.

47 For those patients presented early with no features of intestinal obstruction we will apply manual reduction(
48 MR).

4 Volume XIII Issue II Version I

5 Statistical analysis

51 Data will be coded and fed in a computer to handle statistical and mathematical procedure, to display the
52 analyzed data and present them graphically using SPSS software (statistical package for social sciences).

53 IV.

6 Results

54 The overall number of patients seen with inguinal hernia in (GNCPs) during the study period(from Jan.2011-
55 Jan2013) were 722 cases. from this number there were 43 case presented with incarcerated inguinal hernia.
56 Regarding age distribution < 5 Yrs were 40 cases (93%). from 5-10 yrs were 2 cases (5,4%) while 1case (2,3%)
57 was more than 10 yrs figure (1). The males were 42 cases (97,7%) while the females was one case(2,3%), the
58 waiting time for surgery was 3 months in 14 cases (32,5%) while it was 3-6 months in 10 cases (23,3%) and more
59 than 6 months in 19 cases (44,2%) Table 1. (6). patients from Gezira state were 42 cases (97.7%) and there was
60 1 case (2.3%) from out Gezira state. within the Gezira state We noticed that most of our candidate came from
61 rural areas 31 cases (72,1%) while 12 cases (27,9%) from urban areas.

62 All candidates 43 cases (100%) presented with irreducible inguinal lump with positive silk sign. The affected
63 site was right in 27 cases (62,8%) while it was left in 15 cases (34.9%) and bilateral in 1case (2.3%).

64 28 cases (65.1%) present with obstructive symptoms while not in 15 cases (34.9%). Intra operatively transverse
65 inguinal incision was applied, the contents were small bowel in 24 cases (55,8%) while in 2cases (4,7%) it was
66 large bowel and testes in 5 cases (11,6%).the condition of the contents were viable in 31 cases (72,1%) while dusky
67 in 2 cases (4,7%) and gangrenous bowel in 1 case (2,3%) that required resection & anastomosis. simple reduction
68 was performed in 40cases (93%) while hot wrapping done in 2 cases (4,7%) and resection & anastomosis done in
69 1 case (2,3%). Regarding scrotal swelling post operatively was 27 cases (62,8%) while was not found in 7 cases
70 (16,3%) Table (1).

71 Figure 1

7 Discussion

72 The total number of patients seen with inguinal hernia in (GNCPs) during the study period were 722. Out of
73 this number there were 43 (5.9%) cases presenting with incarcerated inguinal hernia.

74 In this study, 40 patients (93%) were under 5 year of age and 2 cases (4,7%) under 10 year of age and 1 case
75 (2.3%) was >10 year figure (1) . This study has shown that incarceration is more frequent under 5year of age
76 and also that the incidence of incarceration decreases by increasing in age. In comparative to literature review
77 the incidence of pediatrics inguinal hernia is highest during the first year of life and then gradually decreases
78 thereafter. One-third of children undergoing surgery for hernia are less than 6 months of age . ??6) Most of the
79 patient came from rural area 31 cases (72,1%) while 12 cases (27,9%) from urban areas related to Gezira state
80 fig. (2), perhaps this may indicate the lower educational status in our rural areas and their lack of awareness
81 towards this problem. The fact that inguinal hernias are not self-limiting and that there is a risk of complications
82 dictate the need for expedient surgical repair. (7) Among our study we found that 28 cases (65,1%) of patients
83 underwent surgical exploration from the start because they present late with symptoms and signs of intestinal
84 obstruction. while only 15 cases (34,9%) were managed initially by manual reduction (MR). We observed that
85 those presented with no obstructive features came from areas near the hospital and presented rapidly when they
86 noticed the lump became irreducible.

87 A wait time for surgery of more than 14 days was associated with a doubling of the risk of incarceration.
88 The overall rate of incarceration is comparable to those reported by others. (8)(9) In a retrospective report of
89 institutional-based data, Stylianos and colleagues (10) reviewed all cases of children with an incarcerated hernia
90 who presented to an emergency department. They calculated a risk of hernia incarceration of 9% related to
91 a mean wait time of 8 days (range 0.5-28 days). we have long waiting time in comparable with above results
92 we have a minimum waiting for at least 3 months and this had been explained by shortage of resources and
93 overloaded hospital work. Our result showed that the waiting time was < 3months in 14 cases (32.6%) while it
94 was 10 cases (23.3%) waiting for 3-6 months and 19 cases (44.2%) in a period more than 6 months. We noticed
95 that most of our patients wait 6 months or more after booking for surgery this may contributed to a lot of factors
96 such as lack the parents information regarding this problem, traditional concepts, socioeconomic and financial
97 factors.

100 Among our patients we have 15 cases (34,9%) presented with no features of intestinal obstruction successful
101 attempt of (MR) was applied for 9 cases while failed in 6 cases so open surgery was carried out. For those
102 manually reduced admission for 24 hrs later for observation then operated within the next 24hrs, this going with
103 the study done by Gahukamble and Khamage whom concluded that all children with inguinal hernia should have
104 hernia repair within 5 days after reduction of incarceration as a precaution against recurrent incarceration. (11)
(D D D D) I ^{1 2}



Figure 1:

105

¹© 2013 Global Journals Inc. (US)

²() I Impact of Socioeconomic Status on Development of Incarcerated Inguinal Hernia in Children Treated in Gezira National Center for Paediatrics Surgery (GNCPS)




Figure 2: Figure 2

1

Valid	Frequency	Percentage
Yes	27	62.8%
No	7	16.3%
Total	34	79.1%

Figure 3: Table 1 :

-
- 106 [Borenstein et al. ()] 'Effect of subspecialty training and volume on outcome after pediatric inguinal hernia
107 repair'. S H Borenstein , Tt , A Wajja . *J Pediatr Surg* 2005. 40 p. .
- 108 [Ameh ()] 'Incarcerated and strangulated inguinal hernias in children in Zaria'. Ameh . *Nigeria. East Afr Med J*
109 1999. 76 (9) p. .
- 110 [Bronsther et al. ()] 'Inguinal hernia in children-a study of 1000 cases and a review of the literature'. B Bronsther
111 , Am , C Elboim . *J Am Ed Wom Assoc* 1972. 27 p. 524.
- 112 [Harper et al. ()] 'Inguinal hernia: a common problem of premature infants weighing 1000g or less at birth'. R
113 G Harper , Ga , C Sia . *Pediatrics* 1975. 56 p. 112.
- 114 [Takehara and Arakawa ()] 'Laparoscopic Strategy for Inguinal Ovarian Hernias in Children: When to Operate
115 for Irreducible Ovary'. H Takehara , Y Arakawa . *J Laparoendosc Adv Surg Tech* 2009. 19 p. . (Suppl)
- 116 [Lloyd and Hernia (ed.) ()] *Operative pediatric surgery*, D A Lloyd , Hernia . Ziegler M, Azizkhan R, Weber T
117 (ed.) 2003. New York: McGraw-Hill. p. .
- 118 [Neutra et al. ()] 'Risk of incarceration of inguinal hernia in Cell Colombia'. R Neutra , Va , R Ferrada , R Galan
119 . *J Chron Dis* 1981. 34 (11) p. .
- 120 [AP C (ed.)] *The anatomy and surgical treatment of abdominal hernia*, AP C (ed.) London: Longman & Co.
- 121 [Sj W ()] 'The incidence of external hernias in premature infants'. Sj W . *Acta Pediatr* 1962. 2003. McGraw-Hill.
122 51 p. . (ric surgery)
- 123 [Boley et al. ()] 'The irreducible ovary: a true emergency'. S J Boley , D Cahn , T Lauer . *J Pediatr Surg* 1991.
124 26 (9) p. .