

# Micronutrient Malnutrition, A Tragedy To Childhood Growth And Education

Dr. F. N. Uchendu<sup>1</sup>

<sup>1</sup> National Open University of Nigeria

Received: 29 March 2011 Accepted: 25 April 2011 Published: 6 May 2011

## Abstract

Micronutrient malnutrition is a serious childhood dietary problem in developing nations. Deficiencies in vitamins A and B, iron, folic acid and zinc, are preventable causes of poor childhood growth and school performance. Sustainable strategies exist to eradicate malnutrition. This paper discusses the negative effect of vitamin and mineral malnutrition on childhood growth and education, and effective strategies to eliminate them.

**Index terms**— Micronutrient malnutrition, Preschool-age children, Growth, Education, Strategies

## 1. I.

micronutrients are nutrients needed in minute specific quantities in the body. Most of them are not generated in the body but are derived from food intake. These micronutrients include vitamins A and B<sub>12</sub>, iron, folic acid, iodine, and zinc. Prolonged inadequate intake of foods rich in these micronutrients result in their deficiencies. Most developing countries are battling with hunger, poverty and high rate of unemployment. This gives rise to food insecurity in most of the households.

One-third of the world's population suffers from micronutrient deficiencies, due primarily to inadequate dietary intake (Fielder and Macdonald, 2009). Vitamin A is a fat-soluble vitamin, essential for vision in dim light, cellular, bone and tooth growth, formation and maintenance of healthy skin, hair, and mucous membranes, reproduction and immunity boosting. Vitamin A is so important in embryological development that without it, the fertilized egg cannot develop into a fetus (Brody, 2007). Its deficiency results in night blindness or impaired dark adaptation, lowered immunity to infections such as measles, diarrhoea, chickenpox, respiratory infections, anemia, poor growth, slowed bone development, blindness and death. All these have disastrous effect on the healthy growth and school performance of a child. Vitamin A deficiency (VAD) can be severe in children 6 years of age while blindness is more prevalent in children below 3 years. The average preschool child requires 400 µg of vitamin A daily for healthy vision, bone growth, reproduction, cell division, and cell differentiation. This must be derived from intake of foods rich in vitamin A such as fatty meat, School of Science and Technology National Open University of Nigeria, Lagos, Nigeria. +2348037065874; Email: Uchendu\_flo@yahoo.com egg, milk, butter, margarine, palm oil, fortified foods, dark green leafy vegetables and yellow fruits. For many parents in developing countries, apart from plant sources of vitamin A, animal sources are a luxury only enjoyed by the rich. The poor only depend on plant sources and it has been established that the efficiency of the conversion of plant sources of vitamin A (Provitamin A carotenoids) to vitamin A [bioefficacy] in a mixed diet is less than was previously thought (Wardlaw and Kessel, 2002). Retinol Equivalent overestimated the contribution of carotenoids to vitamin A needs until now (Thurnham and Northrop-Clewes, 1999). Then, I RE = I µg of all-trans Retinol = 6 µg of all-trans beta-carotene = 12 µg other carotenoids but the true contribution is I RAE = 12 µg of beta-carotene = 24 µg of other Carotenoids. Consequently, a child whose vitamin A source is solely dependent on plant sources of vitamin A will become vitamin A deficient over time. Worst still, the main component of the diet of children in developing countries is starchy foods. Another cause of vitamin A deficiency is the drifting from local foods rich in vitamins and minerals to fast foods which are highly refined as a result of the influence of urbanization and western culture. Prolonged shortage of vitamin A rich foods in the diet of the child, results

in low vitamin A Recommended Dietary Allowance (RDA) and eventual depletion of any available vitamin A stored in the liver.

World Health Organization (WHO) defines vitamin A deficiency as the tissue concentration of vitamin A low enough to cause adverse consequences even without clinical evidence of xerophthalmia ?? ??).

Iron deficiency occurs when iron requirements cannot be met by absorption from the diet, such as during periods of rapid growth (infancy, adolescence), in pregnancy, and as a result of menstrual or pathological blood loss (Hurrell, et al. 2010). Developing countries' diets are predominantly dominated by plantbased foods and so limit iron absorption due to their high phytate and polyphenol contents (Hurrell, 2002; ??immermann, et al. 2005;Hurrell, et al. 2010). Iron deficiency in infants and young children is associated with delayed mental and motor development (Lozoff, 2007). In summary, an iron-poor diet and rapid growth are primary causes of iron deficiency in infants and preschool children ??Li et al. 2009). Iron deficit children may experience emotional problems and fail to meet educational goals later in life leading to a negative impact on learning capacity in adulthood ??Hurrell et al. 2011).

An estimated 240,000 annual cases of folic acid-preventable spina bifida and anencephaly has been recorded (Oakley et al. 2004). Adequate consumption of folic acid before pregnancy and during the early weeks of gestation protects fetuses from developing neural tube defects (Folic Acid Working Group et al. 2010).

Interest in zinc was stimulated when zinc supplements given to short children and failure-to-thrive infants in the U.S. city of Denver improved growth . Trial studies concluded that zinc supplements are likely to improve the height gain of the most stunted children and to improve the weight gain of those with low plasma zinc concentrations . Intakes of absorbable zinc are often low in children and growthstunting occurs nearly universally during the first two years of life in underprivileged populations .

Because vitamin B-12 is found only in animal products, many poor populations, or those that avoid animal products for religious or other reasons, consume little or no vitamin B-12. Low serum B-12 concentration is associated with a higher risk of potentially irreversible harm to memory, cognitive function, and nerve conduction, as well as a higher risk of megaloblastic anaemia. Studies among low income people in Guatemala, Mexico, Nepal, Venezuela, and other countries show that 25 to 50 percent of individuals are deficient . Vitamin B-12 deficiency occurs in populations with low consumption of animal-source foods which are the only natural source of the vitamin. Vitamin B-12 deficiency is also prevalent in developed countries among the elderly due to their inability to release and absorb the vitamin from foods (Rosenberg, 2010).

## 2 II.

### 3 EFFECT OF MICRONUTRIENT MALNUTRITION ON THE GROWTH AND EDUCATION OF PRESCHOOL CHILDREN

Both chronic under nutrition and severe clinical malnutrition in childhood are related to scholastic that malnutrition in foetus and young children causes disturbances in the morphological and functional development of the central nervous system thus affecting the cognitive and emotional development of the child.

Micronutrient Malnutrition causes birth defects, mental retardation, learning difficulties, compromised immune systems, low work capacity, blindness and death. These consequences definitely have a negative effect on the healthy growth of the child via education in terms of intelligent quotient (I.Q) and school performance. There is evidence that a poor diet associated with high fat, sugar and processed food content in early childhood may be associated with small reductions in I.Q in later childhood (Northstone et al.

## 4 Global Journal of

## 5 Medical Research

Volume XI Issue I Version I May 2011 28 countries such as Zambia, U deficiency as a public health problem. The end-of-decade goal for elimination of VAD was widely promoted in the 1990s but, although progress was made, the goal was not met ?? Underwood, 2006). However, many countries now have success stories e.g. Vietnam, Venezuela, Bangladesh, Indonesia, Phillippine Egypt does not have Vitamin A and some parts of India. They have been able to reduce VAD and xerophthalmia to below WHO cut-off point constituting a public health problem ( West, 2002; Ramakrishnan and Darnton-Hill, 2002). The International Consultative Group (IVACG) recently adopted a cut off of more than 15% of pre-school population with serum retinol below 0.70 mol (or 20 g/dL) or displaying abnormal impression cytology as indicative of VAD (West, 2002;Wasantwisut, 2002 (Miguel and Kremer, 2004). Agreeably, studies have recognized that there is a relationship between school performance and child health nutritional status. Early malnutrition affects brain structure and learning ability (Liu et al. 2003). Malnourished children are inactive, inattentive and lack curiosity and explorative abilities and these affect their educational performance. Malnutrition also results in language retardation. A more serious effect of malnutrition is its permanent effect on children who were less than six months when they were malnourished which is a serious handicap on schooling and has a close impact on the ability to learn, read, and write (Amcoff, 1981). The consequences of these are school failure. Iron deficiency and anemia lead to compromised ability to

learn and poor physical growth, which can impair school performance ultimately resulting in retarded cognitive, motor and academic ability (REAP, 2010). Consequences of iron deficiency in children includes anaemia, poor growth, weak immune system, reduced cognition and development, poor attention span, concentration, memory, learning ability, poor muscle function and manual dexterity, behaviour, and social interaction. It has also been reported that even though VAD does not directly affect school performance, it may do so indirectly via its effect on infectious related morbidity, which in turn will affect school attendance. This was demonstrated in a study on school teachers' awareness about scholastic performance and nutritional status of Egyptian school children (van-Stuijvenberg, 2005). Many studies have shown associations between hunger, poor dietary intakes, stunting, underweight and poor school performance stating that children who were stunted, anaemic, or iodine deficient had poorer school achievement levels and attendance than other normal children. Figs. ?? and 2 are pictures showing some malnourished children in developing countries. Some already have blotted faces.

Folate deficiency results in learning disabilities. Recent evidence suggests that poor maternal folate status is also associated with a higher risk of abnormal pregnancy outcomes, including eclampsia, premature delivery, and birth defects such as club foot and cleft palate .

III.

## 6 NUTRITIONAL INTERVENTION STRATEGIES

The optimal growth, physical and intellectual development which will enable children to learn and reach their full potential in life must not be jeopardized. Strategies that have been identified to fight micronutrient deficiencies include exclusive breast-feeding, vitamin A supplementation (through use of capsules), nutrition education/communication, dietary diversification, food fortification, biofortification, home gardening, and disease control. Fortification of food with vitamin A, iron and folate results in smarter, stronger, healthier children. It increases the national Intelligent Quotient (I.Q.) by 5%, national GDP by 2% and prevents the 200,000 cases of severe disability in babies yearly (Moench-Pfanner, 2007).

Many local foods, fruits, and plants have been reported to be good sources of micronutrients. They are available in abundance and very cheap. Consumption of varieties of local foodstuffs will help the children have adequate nutrient stores especially during their season when the fruits, vegetables and foodstuffs are usually wasted due to poor storage facilities. Examples are palm oil, yellow maize (corn), orange-fleshed sweet potatoes, mango, banana, tomatoes etc. Red palm oil has proved effective in combating VAD in South Africa. Red palm oil has been used to fortify biscuits which provided beta-carotene at 50% of the Recommended Dietary Allowance (RDA) and red palm-oil based bread spreads for primary school feeding programs in South Africa (van Stuijvenberg, 2005). Similar fit has also been performed in Burkina Faso (Zagre, et al. 2004). The result from South Africa has a significant long-term positive implication on learning and school performance in children that are vitamin A deficient ??Zimmermann, et al. 2004). This technology could be transferred. There has been an increased promotion and utilization of orange-fleshed sweet potato to combat vitamin A deficiency in Burkina Faso, Uganda, and South Africa (Vebamba, 2004;Kapinga, et al. 2004;van-Stuijvenberg, 2005). Efforts to identify Nigerian traditional edible plants that are good sources of vitamin A have yielded positive results. Recent findings reported that Baobab leaf (*Adansonia Digitata* L) is a rich source of beta-carotene (156.5?g/g) and iron. It's use on rural Nigerian children improved their vitamin A and iron status by decreasing the number of children with serum retinol levels below 20?g/dl significantly from 21.25% to 10.0% and their serum beta-carotene rising from 6.8?g/dl to 14.1?g/dL while serum ferritin of children with low serum ferritin (HB<11.0g/dl) significantly increased to 19.0?g/dL (Nnam, 2004). Nnam and Onyeka (2004) reported that Sorrel (*Hibiscus sabdariffa* calyx is a good source of micronutrients to combat VAD. According to them,

7 )

Sorrel Calyx is a good source of retinol 285.29RE iron 833.00mg/100g), and ascorbate (53.00mg/100g). Rural Nigerian children fed with sorrel calyx based diet had higher hemoglobin (HB) and serum retinol (SR) levels and no VAD symptoms as against the control. Other lessons could be learnt from Malaysia where nutritious foods such as milk and multi-vitamins are distributed to families, primary school children, pregnant women and lactating mothers with twins. Milk is a good source of iron. In a study using milk fortified with 15mg iron (as iron sulphate/L) the incidence of anaemia was reduced from 36.4% (control) to 12.7% in the fortified group (Blum, 1997).

IV.

## 8 CONCLUSION

Studies have shown that maintaining high levels of micronutrients in the diet of children is important for optimal growth, development of their normal learning and cognitive functions. Adequate vitamins and mineral diet is needed for healthy and productive children. This should be a matter of right and not charity. Agriculture should be emphasized more and research encouraged to identify more local foods rich in micronutrients in various communities in developing nations. Nutrition education/communication should encourage increased consumption of animal sources of vitamins and minerals among preschool-aged children. Nutritional policies in developing countries should be encouraged in favour of children, pregnant and lactating mothers.




Figure 1: Fig. 1 .Fig, 2 .

1

Liberato

[Note: M INTRODUCTION deficiency (FORTAF, 2000) (Table]

Figure 2: Table 1 .

---

<sup>1</sup>Global Journals Inc. (US) Micronutrient Malnutrition, A Tragedy to Childhood Growth and Education ©  
<sup>2</sup>Global Journals Inc. (US) ©  
<sup>3</sup>. Li Y. O., Diosady L.L., and Wesley A. S. (2010). Iron in vitro bioavailability and iodine storage stability in double-fortified salt. Food and Nutr. Bull., vol. 30, no. 4. Pp. 327-335.  
<sup>4</sup>Global Journals Inc. (US) Micronutrient Malnutrition, A Tragedy to Childhood Growth and Education ©  
Références Referencias  
<sup>5</sup>Global Journals Inc. (US) Micronutrient Malnutrition, A Tragedy to Childhood Growth and Education ©

---

[ Accessed ()] , Accessed 1/2011. 17.

[ Accessed (2011)] , Accessed . 3/2/2011.

[Allen (2001)] *2020 Focus 5 (Health and Nutrition Emerging and Reemerging Issues in Developing Countries)*, L H Allen . <http://www.soilandhealth.org/01aglibrary/Arun/micronutrients.pdf>. Accessed 15/4/2011 2001. February 2001. (Brief 10 of 11)

[Fiedler and Macdonald ()] 'A strategic approach to the unfinished fortification agenda: Feasibility, costs, and costeffectiveness analysis of fortification programs in 48 countries'. J L Fiedler , B Macdonald . *Food and Nutr. Bull* 2009. 30 (4) p. .

[Northstone et al. ()] 'Are dietary patterns in childhood associated with IQ at 8 years of age? A population-based cohort study'. K Northstone , C Joinson , P Emmet , A Ness , T Paus . 10.1136/jech.2010.111955. *J Epidemiol CommunityHealth* 2010.

[West et al. (2008)] *Background document prepared for the second technical workshop on wheat flour fortification Atlanta*, K P West Jr , O Klemm , Q Dary , P Johnson , C Randel , Northrop -Cleves . 2008. March -3 April, 2008. Draft 31 March, 2008. (Vitamin A fortification group)

[Nnam ()] 'Effect of Baobab (*Adansonia L.*) diet on Vitamin A and Iron status Nigerian children'. N Nnam . *Report of the XXI 2004 International Vitamin A Consultative Group (IVACG) Meeting. Pp*, 2004. p. 66.

[West ()] 'Extent of Vitamin A Deficiency among Preschool Children and Women of Reproductive Age. Proceedings of the XX International Vitamin A Consultative Group Meeting'. K West Jr . *J. Nutr* 2002. 132 p. .

[Food Fortification in Africa: A strategy to eradicate Vitamin A and Mineral deficiencies] *Food Fortification in Africa: A strategy to eradicate Vitamin A and Mineral deficiencies*, <http://www.innovations-report.com/htm29/06/2006>

[Berry et al. ()] 'Fortification of flour with folic acid'. R J Berry , L Bailey , J Mulinare , Carol Bower . *Food and Nutr. Bull* 2010. 31 (1) p. .

[Liberato and Pinheiro-Sant'ana ()] 'Fortification of Industrialized Foods with vitamins'. S Liberato , H Pinheiro-Sant'ana . *Rev. Nutr* 2006. 2006. 19. 2 p. .

[Health, Nutrition, Care and Educational Performance REAP ()] 'Health, Nutrition, Care and Educational Performance'. [http://reap.stanford.edu/docs/nutrition\\_and\\_education/](http://reap.stanford.edu/docs/nutrition_and_education/) Accessed 10/2/2011 REAP 2010.

[Hurrell ()] 'How to ensure adequate iron absorption from iron-fortified food'. R Hurrell . *Nutr. Rev* 2002. 2002. 60 p. . (discussion S43)

[Vebamba et al. ()] 'Introduction of Orange-fleshed sweet potatoes in Gourma Province'. O Vebamba , M Bendeche , I Mariano , S Baker . *Report of the XXII 2004 International Vitamin A Consultative Group (IVACG) Meeting. Pp*, (Burkina Faso) 2004. p. 71.

[Lozoff ()] 'Iron deficiency and child development'. B Lozoff . *Food Nutr. Bull* 2007. 2007. 28 p. .

[Zimmerman et al. ()] 'Iron deficiency due to consumption of a habitual diet low in bioavailable iron: a longitudinal cohort study in Moroccan children'. M Zimmerman , N B; Chouki , R F Hurrell . *Am J. Clin. Nutr* 2005. 2005. 81 p. .

[Liu et al. ()] 'Malnutrition at Age 3 Years and Lower Cognitive Ability at Age 11 Years'. J Liu , A Raine , P H Venables , C Dalais , S Mednick . *Arch Pediatr. Adolesc* 2003. 2003. 157 p. .

[Black et al. ()] 'Maternal and Child undernutrition: global and regional exposures and health consequences'. R E Black , L H Allen , Z A Bhutta , F Caulfield , M De Onis , M Ezzati , C Mathers , J Rivera . *Lancet* 2008. 2008. 371 p. .

[Allen (2001)] 'Micronutrients.2020 Focus 5 (Health and Nutrition Emerging and Reemerging Issues in Developing Countries'. L H Allen . *Brief 10 of 11*, 2001. February 2001.

[Zagre et al. ()] *Moving with Red Palm Oil towards dietary diversification strategy for controlling Vitamin A deficiency in Burkina Faso*, N Zagre , H Delishe , Depeuch , P Traissac . 2004. (Report of the XXII 2004 International Vitamin A Consultative Group (IVACG) Meeting)

[Kraemer Zimmerman (ed.)] [http://sightandlife.org/images/stories/pageimages/content/publications/nutritional\\_anaemia\\_book.pdf](http://sightandlife.org/images/stories/pageimages/content/publications/nutritional_anaemia_book.pdf). Accessed 28/3/2011. 2011/01/31/tropej.fmq119.full#off.1. Accessed 7/2/2011 *Nutritional anaemias. Available at*, K Kraemer, M Zimmerman (ed.)

[David and Christine ()] *Optimal Nutrition: Vitamin A and the Carotenoids Proceedings of the Nutrition Society*, Thumham David , I , Northrop-Clewes Christine , A . 1999. 58 p. .

[Ramakrishan and Darnton-Hill ()] 'Proceedings of the XX International Vitamin A Consultative Group Meeting'. U Ramakrishan , I Darnton-Hill . *J. Nutr* 2002. 132 p. . (Assessment and Control of Vitamin A Deficiency Disorders)

- [Oakley et al. ()] *Recommendations for accelerating global action to prevent folic acid-preventable birth defects and other folate-deficiency diseases: meeting of experts on preventing folic acid-preventable neural tube defects*, G P Oakley , K N Bell , M B Weber . <http://www.ncbi.nlm.nih.gov/pubmed/15390317>. Accessed 15/4/2011 2004.
- [Wasantwisut ()] 'Recommendations for Monitoring and Evaluating Vitamin A Programs: Outcome Indicators'. E Wasantwisut . *Proceedings of the XX International Vitamin A Consultative Group Meeting*, (the XX International Vitamin A Consultative Group Meeting) 2002. 132 p. .
- [Hurrell et al. ()] 'Revised recommendations for iron fortification of wheat flour and an evaluation of the expected impact of current national wheat flour fortification programs'. R Hurrell , P Ranum , S De Pee , R Beibinger , L Hultheim , Q Johnson , S Lynch . *Food and Nutr. Bull* 2010. 31 (1) p. .
- [Van-Stuijvenberg ()] 'School Feeding system as a vehicle for Micronutrient Fortification: Experience from South Africa'. M Van-Stuijvenberg . *Food Nutr. Bull* 2005. 2005. (26) p. .
- [Nnam and Onyeka ()] 'Sorrel (*Hibiscus sabdariffa*) Calyx as a Promising Source of beta-carotene to control Vitamin A Deficiency'. N Nnam , N Onyeka . *Report of the XXII 2004 International Vitamin A Consultative Group (IVACG) Meeting. Pp*, 2004. p. 66.
- [Blum (ed.) ()] *Status Paper: Food fortification to end micronutrient malnutrition*, M Blum . Division, F. Hoffman-La Roche Limited (ed.) 1997. Basel, Switzerland. p. 97.
- [Summary Report. Second Technical Workshop on Wheat Flour Fortification: Practical Recommendations for National Application] 'Summary Report. Second Technical Workshop on Wheat Flour Fortification: Practical Recommendations for National Application'. *The Flour Fortification Initiative*, (Stone mountain, Georgia, USA) 2008. March 30 to April 3, 2008. p. .
- [The African Context (2000)] [http://www.fortaf.org/the\\_african\\_context.htm](http://www.fortaf.org/the_african_context.htm) *The African Context*, 2000. 2/2/2010.
- [Wardlaw and Kessel ()] 'The Fat Soluble Vitamins'. G Wardlaw , M Kessel . *Perspectives In Nutrition* 2002. McGraw Hill. 5 p. . (th ed))
- [Amcoff ()] *The Impact of Malnutrition on Learning Situation*, S Amcoff . <http://www.unesdoc.unesco.org/images/0014/001429/142929eo.pdf>. Accessed 10/2/2011 1981. p. .
- [Rosenberg ()] 'The opportunity of flour fortification: Building on the evidence to move forward'. I H Rosenberg . *Food and Nutr. Bull* 2010. 31 (1) p. .
- [Moench-Pfanner (2007)] *The Status Quo of Food Fortification. 3 rd International Muhlenchemie Symposium Hamburg*, R Moench-Pfanner . 14/15 June 2007. Hotel Hafewen Hamburg.
- [Underwood (2004)] 'Vitamin A Deficiency disorders: International efforts to control preventable "pox'. B A Underwood . *J. Nutrition* 2004. 2004 Jan: 134. (1) p. .
- [Brody] *Vitamin A deficiency Encyclopedia of medicine*, T Brody . [http://www.findarticles.com/p/article-mig2601/is0014/ai\\_2601001456](http://www.findarticles.com/p/article-mig2601/is0014/ai_2601001456)
- [Kapinga et al. ()] 'Vitamin A Partnership For Africa: A Food based Approach to combat Vitamin A Deficiency through increased utilization of Orange-fleshed Sweet potato'. R Kapinga , R Anderson , D Zhang , M Herman , F Opio . *of the XXII 2004 International Vitamin A Consultative Group (IVACG) Meeting. Pp*, 2004. p. 70. (Report)
- [Zimmerman et al. ()] 'Vitamin A Supplementation improves iodine efficacy in Goitrous vitamin A-deficient children receiving iodized salt'. M Zimmerman , R Wegmuller , F Rohner , C Zeder , N Chaouki , T Torresani . *of the XXII 2004 International Vitamin A Consultative Group (IVACG) Meeting. Pp*, 2004. p. 89. (Report)
- [Arcanjo et al. (FEB)] 'Weekly Iron Supplementation for the Prevention of Anemia in Pre-school Children: A Randomized, Double-blind, Placebo-controlled Trial'. F P Arcanjo , C C Arcanjo , Oms Amancio , Jap Braga , Ájm Leite . 10.1093/TROPEJ/FMQ119. [http://tropej.oxfordjournals.org/content/early/JTROP PEDIATR FEB. 57 \(1\) .](http://tropej.oxfordjournals.org/content/early/JTROP PEDIATR FEB. 57 (1) .)
- [McLean et al. (ed.) ()] *Worldwide prevalence of anaemia in preschool aged children, pregnant women and non-pregnant women of reproductive age*, E Mclean , I Egli , B De Benoist , D Wojdyla . Kraemer k., Zimmerman M.B. (ed.) 2007. 2007. Basel, Switzerland: Sight and Life Press. p. .
- [Miguel and Kremer ()] 'Worms: Identifying Impacts on education and Health in the Presence of Treatment Externalities'. E Miguel , M Kremer . *Econometrica* 2004. 2004. (1) p. .