

1 Pharmacological Activity and Chemical Constituents of Eclipta 2 Alba

3 Dharmender Jaglan¹, Amandeep Singh Brar² and Rupamjot Gill³

4 ¹ PTU, Jalandhar

5 Received: 14 December 2012 Accepted: 3 January 2013 Published: 15 January 2013

7 **Abstract**

8 Eclipta alba is a herb commonly found throughout India. This plant is known to have various
9 pharmacological activities and is traditionally used in treatment but it lacks adequate
10 scientific proof of this activity and constituents responsible for it. The present paper describes
11 the phytochemical and pharma-cological investigations of Eclipta alba.

13 **Index terms—**

14 Ethnopharmacology : Eclipta Alba (L.) has been used in various parts of tropical and sub-tropical regions
15 like south America, Asia, Africa. There are three kinds of Eclipta Alba-the white-flowering, the yellowflowering,
16 and the black-fruiting, but all three grow throughout India by marshes, rivers, and lakes or on the foothills of
17 the Himalayas. It is an active ingredient of many herbal formulations prescribed for liver ailments and shows
18 effect on liver cell generation. It is used as a tonic and diuretic in hepatic and spleen enlargement. It is also
19 used in catarrhal jaundice and for skin diseases (Scott Treadway1998). The alcoholic extract of the plant has
20 shown antiviral activity against Ranikhet disease virus (Sunita Dalal, et al., 2010). The plant is commonly used
21 in hair oil all over India for healthy black and long hair (Roy RK, et al., 2008). The fresh juice of leaves is used
22 for increasing appetite, improving digestion (Cheryl Lans, 2007) and as a mild bowel regulator. It is commonly
23 used in viral hepatitis to promote bile flow and protect the parenchyma and popularly used to enhance memory
24 and learning (Otilia Banji, et al., 2007).

25 Authors ? ?: Chandigarh College of Pharmacy, Landran (Mohali) M.pharmacy (Pharmacology Department).
26 e-mails: dharmenderjaglan@gmail.com, amandeepbrar678@gmail.com, Rjgill@gmail.com

27 The plant has a reputation as an antiageing agent in Ayurveda (Thakur VD, et al., 2005). It is used as a general
28 tonic for debility. Externally it is used for inflammation (Amritpal singh, et al., 2008; and Mahesh Sawant, et al.,
29 2004), minor cuts and burns and the fresh leaf-juice is considered very effective in stopping bleeding (Mukherjee
30 DR, et al., 1976). Leaf juice mixed with honey is also used for children with upper respiratory infections and
31 also used in eye and ear infections. It is a source of coumestans-type compounds used in phytopharmaceutical
32 formulations of medicines prescribed for treatment of cirrhosis of the liver and infectious hepatitis (Wagner H.
33 et al., 1986). (Tewtrakul S, et al., 2007). Vedic Guard, a polyherbal formulation is a synergistic combination of
34 16 medicinal plant extracts contains Eclipta Alba as a major ingredient (Rema Razdan, et al., 2008). Charaka
35 advises taking the juice of Eclipta Alba with honey to prevent the onset of senility, and its oil as the best
36 medicated massage oils for rejuvenation therapies.

37 Phytochemistry: Eclipta Alba (L.) contains wide range of active principles which includes coumestans, alka-
38 loids, flavonoids, glycosides, polyacetylenes, triterpenoids. The leaves contain stigmasterol, β -terthienylmethanol,
39 wedelolactone, demethylw-edelolactone and 1986). The roots give hentriacontanol and heptacosanol. The roots
40 contain polyacetylene substituted thiophenes. The aerial part is reported to contain a phytosterol, β -amyrin in
41 the n-hexane extract and luteolin-7-glucoside, β -glucoside of phytosterol, a glucoside of a triterpenic acid and
42 wedelolactone in polar solvent extract (Jadhav VM, et al., 2009). The polypeptides isolated from the plant yield
43 cystine, glutamic acid, phenyl alanine, tyrosine and methionine on hydrolysis. Nicotine and nicotinic acid are
44 reported to occur in this plant (Jadhav VM, et al., 2009).

45 Coumestan: Coumestan is an organic compound that is a derivative of coumarin. Coumestan forms the central
46 core of a variety of natural compounds known collectively as coumestans. Coumestans, including a coumestrol and

Pharmacological Activity and Chemical Constituents of *Eclipta Alba*

47 phytoestrogen are found in a variety of plants. Because of the estrogenic activity of some coumestans, a variety of
48 syntheses have been developed that allow the preparation of coumestans so that their pharmacological effects can
49 be explored. The major coumestan isolated from *Eclipta Alba* includes wedelolactone 0.5-0. Immunomodulator
50 activity : The protection of neuronal tissue may be possible due to the immunomodulatory action of *Eclipta*
51 *Alba*. Due to methanol extract of whole plant of *Eclipta Alba* (1.6% wedelolactone), the phagocytic index
52 antibody titer, phagocytic index and WBC count increased. Due to inhibition of non-specific humoral (lysozyme,
53 antiprotease and complement) and cellular (myelop-eroxidase content), the reactive oxygen and nitrogen species
54 are produced. The dietary intake of *Eclipta Alba* aqueous leaf extract also enhance the non-specific immune
55 response and disease resistance of *o.massambicus* against *A.Hydrphila*. *Eclipta Alba* serve as a potential memory
56 modulator also (Ghosh M et al., 1984 And Roitt I et al., 1998, Hudson L et al., 1991). Hair growth and alopecia
57 : *Eclipta Alba* is a well known Ayurvedic herb for hair growth. *Eclipta Alba* is used in hair oil preparation
58 since it promotes hair growth and maintains hair black. 10%w/v of *Eclipta Alba* used as a main ingredient in
59 the preparation of herbal formulation for hair growth. Petroleum ether and ethanolic extract were also used in
60 oleaginous cream and applied topically ??Roy R, et al., 2008).

61 Anticancer activity : The methanolic extract of *Eclipta Alba* has the inhibitory effect against colon cancer
62 due to inhibition of proliferation of cancer cells in a concentration dependent manner (Ruddon R. W, et al., 1995
63 and St. Luke, et al., 2007). Methanolic extract of *Eclipta Alba* was evaluated for its anticancer activity against
64 Ehrlich Ascites Carcinoma (EAC) in Swiss albino mice. On day 1, the extract of *Eclipta Alba* at a dose of 250 and
65 500 mg/kg body weight were administered orally and continued for 9 consecutive days. The anticancer activity
66 was examined by determining the tumor volume, tumor cell count, viable tumor cell count, nonviable tumor cell
67 count, mean survival time and increase in life span in experimental animal models. The extract increased the life
68 span of EAC treated mice and restored the hematological parameters as compared with the EAC bearing mice.
69 Thus, study revealed that the methanolic extract of *Eclipta Alba* showed anticancer activity in the tested animal
70 models (Malaya Gupta, et al., 2005). Coumestans are also known to act as phytoestrogens. These compounds
71 are present in soyabeans and clover. In many countries it is used as diet which act as chemopreventive agent
72 in breast and prostate cancer (Neerja Kaushik-Basu, et al., 2008). Dasyscyphin-C (saponins) a newer isolated
73 compound from *Eclipta prostrata* reported to have anticancer cytotoxic activity. It was tested under invitro
74 conditions in HeLa (Human cervical carcinoma) & vero cell lines. At the concentration of 50 μ g/ml it showed a
75 good anticancer-cytotoxic activity on HeLa cells (Khanna, 2008). A rat hepatic stellate cell line (HSCs) was used
76 as in-vitro assay system, the methanolic extract of aerial parts of *Eclipta prostrata* showed significant inhibitory
77 activity on HSCs proliferation (Mi Kyeong .

78 Hepatoprotective effect : *Eclipta Alba* is considered a powerful liver tonic. The hepatoprotective potential
79 of *Eclipta Alba* was studied by assessing the biochemical parameters like lipid peroxide (LPO), superoxide
80 dismutase (SOD), Catalase (CAT), glutathione peroxide (GPx), glutathione reductase (GR), ascorbic acid and
81 γ -tocopherol. Oral administration of the *Eclipta Alba* significantly decreased levels of LPO and elevated the
82 activity of antioxidant enzymes SOD, CAT, GPx, and GR as well as endogenous levels of ascorbic acid and γ -
83 tocopherol. *Eclipta Alba* has show protective effect on experimental liver damage in rats and mice and also used
84 for the treatment of liver cirrhosis and infective hepatitis by reducing centrilobular necrosis, hydropic degeneration
85 and fatty change of the hepatic parenchymal cells (Singh B et al., 1993). The coumestans constituents of the
86 *Eclipta Alba* plant, wedelolactone and demethylwedelolactone, are responsible for the potent antihepatotoxic
87 activities in CCL 4 -galactosamine and phalloidin induced liver damage in rats ??Chopra R et al., 1996). The
88 ethyl acetate fraction of *Eclipta Alba* improves the both enzymatic and non enzymatic antioxidant status in
89 rat liver (Singh B et al., 1993). Hepatoprotective activity also expressed by *Eclipta Alba* by regulating hepatic
90 lysosomal enzymes (Saxena A, et al., 1993).

91 Hepatoprotective activity of methanolic extract and sub fractions of leaves and the chloroform extract and sub
92 fractions of roots of *Eclipta alba* was carried out using carbon tetrachloride-induced liver damage and Lysosomal
93 enzymes level in wistar albino rats. The methanolic extract of leaves and the chloroform extract of roots of *Eclipta*
94 *Alba* showed significant activities and respectively causing 72.8% & 47.96% reduction of lysosomal enzyme. The
95 triterpenoid eclabasaponin fraction from methanolic extract of leaves produced significant (78.78%) and the
96 alkaloidal fraction (60.65%) reduction of carbon tetra chloride induced increase in lysosomal enzyme in blood.
97 Coumestan fraction and triterpenoidal saponin fraction from the chloroform extract of roots produced very
98 significant (75.6%) and (52.41%) respectively reduction of carbon tetra chloride induced increase in lysosomal
99 enzyme levels in blood (Lal V.K, et al., 2010 Antioxidant properties: The antioxidant effects of *Eclipta prostrata*
100 was reported when the level of serum hydroxyl radical (nmol/mg protein per minute) and serum lipid peroxide
101 (nmol/mg protein) levels reduced as compared to untreated group. 100mg/kg dose significantly reduced Carbonyl
102 content of oxidatively modified proteins8. Antioxidant activity of *Eclipta prostrata* was determined by FRAP,
103 radical scavenging activity, reducing activity, and DPPH assay. The antioxidant capacity was increased by
104 increasing the concentration of the extracts from 25 to 100mg/ml27. The antioxidant activity of the hexane,
105 ethyl acetate, ethanol and water extracts of *E. prostrata* was determined by ferric thiocyanate (FTC) (Karthikumar
106 S, et al., 2007).

107 Anthelmintic activity: The methanolic extract of whole plant of *Eclipta Alba* (L.) have the anthelmintic
108 activity.

109 Other pharmacological activities: It has been reported that the importance of free carboxylic acid at C-28

110 position in echinocystic acid derivatives from the methanolic extract *Eclipta prostrata* showed antifibrotic activity
111 (Mi Kyeong . Ethanolic and ethyl acetate fractions of *Eclipta prostrata* were tested for its antibacterial activities
112 against *Escherichia coli*, *Klebsiella pneumoniae*, *Shigella dysenteriae*, *Salmonella typhi*, *Pseudomonas aeruginosa*,
113 *Bacillus subtilis*, and *Staphylococcus aureus* (Karthikumar S, et al., 2007).

114 *Eclipta prostrata* is combined with a non-plant material which is used to bath children suffering from
115 malnutrition for 9 days and used as self medication by AIDS patients in southern Thailand (Sawangjaroen
116 N, et al., 2005 andCheryl Lans. 2007). 16 parts of *Eclipta prostrata* (bhringaraj), 1 part of Triphala formula
117 {*Embllica officinalis* (amalaki), *Terminalia chebula*, (haritaki), *Terminalia belerica* (bibhitaki)}, 1 part of *Caltropis*
118 *gigantean* (arka) and 1 part of *Smilax officinalis* (sariva) mixed with 80 parts of sesame oil and boiled to make a
119 medicated oil which is reported to be used in skin diseases (Bensky Dan, et al., 1986).

120 **1 Global Journal of Medical Research**

Volume XIII Issue VII Version I Year () B ¹



Figure 1:

121

¹© 2013 Global Journals Inc. (US) © 2013 Global Journals Inc. (US) Pharmacological Activity and Chemical Constituents of *Eclipta Alba*



Figure 2: Figure :

1

).

[Note: *Volatile oils*: The volatile components were isolated from the aerial parts of this plant by hydrodistillation and analysed by GC-MS. A total of 55 compounds, which were the major part (91.7%) of the volatiles, were identified by matching mass spectra with a mass spectrum library (NIST 05.L) (Xiong-HaoLin, et al., 2010).*Saponins*: From the whole plant of *Eclipta Alba*, a new triterpene saponin, named eclalbatin, together with alpha-amyrin, ursolic acid and oleanolic acid were isolated. *Dasyphytin C* was isolated from *Eclipta prostrata* which were studied on the HeLa cells for the anticancer activity(Khanna, et al., 2008).*Crude extract* : The crude extract has wound healing properties. The loss of hepatic lysosomal acid phosphatase and alkaline phosphatase by *CCl4* was significantly restored by *Eclipta Alba*. The previous studies shows that hepatoprotective activity of *Eclipta Alba* is by regulating the levels of hepatic microsomal drug metabolizing enzymes(Saxena AK, et al., 1993).The fresh plant is used as self medication by AIDS patients in]

Figure 3: Table 1 :

Sl.No	Chemical con- stituents	Pharmacological activites
1	Wedelolactone	Antihepatotoxic (Nazim Uddin, et al., 2010), Antibacterial (Karthikumar S, et al., 2007), Trypsin Inhibitor, Antivenom (Vianna-da-silva NM, et al., 2003)
2	Eclalbosapogenins	Antiinsrevitalizing (Rupali thorat, et al., 2009), Antiproleferative (Khanna, et al., 2008 and Neerja Kaushik-Basu, et al., 2008), Antigiardial (Sawangjaroen N, et al., 2005)
3	Demethylwaglerin	Antilepatotoxic (Wagner H. et al. 1986), Antihaemorrhage (Mukherjee DR, et al., 1976), Antivenom (Vianna-da-silva NM, et al., 2003), Dye (cosmetic) (Meena AK, et al., 2010)
4	Dasyphyllanthus	Antiviral, Anticancer (Khanna, et C al., 2008)
5	Eclalbatin	Antioxidant(Tewtrakul S, et al., 2007)
6	Ecliptalbin	Verazine Lipid lowering, Analgesic(Maged S. Abdel-Kader et al., 1998)
<p>Anti -Ulcer Activity : Eclipta Alba has the anti-ulcer activity. There are different type of herbal plants which are used to treat anti-ulcer such as The Polyherbal Formulation -RO7D consists of eleven medicinal plants namely Centella asiatica, Cassia auriculata, Cynodon dactylon, Rosa damascene, Myristica fragrans, Nelumbo nucifera, Hibiscus rosa-sinensis, Hemidesmus indicus, Glycyrrhiza glabra, Eclipta alba and Phyllanthus niruri. The Polyherbal Formulation -RO7D exhibited ($P < 0.001$) significant decrease in ulcer index in both the model and significant decrease in the gastric volume in pyloric ligation rat ulcer model. The study indicates that extract RO7D has anti-ulcer activity and its anti-ulcer potential may be due to anti -secretory and cyto-protective activity. (Srinivasan D et al., 2008)</p>		

Figure 4: Table 2 :

improvement in parameters like body weight and lipid profile by enhancing antioxidant defenses to protect against oxidative damage (Antihyperlipidemic properties : It has been reported that in the atherogenic diet induced hyperlipidemic model, the aqueous leaf extract of the *Eclipta prostrata* was given orally to the rats which significantly reduced total cholesterol, triglycerides, total protein. There was a significant elevation in the high density lipoprotein cholesterol levels (Dae-Ik Kima, et al., 2008).

Anti-inflammatory and Analgesic activity : The extract of *Eclipta Alba* was administered orally to investigate anti-inflammatory activity (Amritpal singh, et al., 2008). The anti-inflammatory activity which estimated by using carragenan induced paw oedema model. Inflammation occurs due to activation of platelet activation factors and release of pro-inflammatory mediators such as prostaglandins, kinins, tumor necrosis factors and nitric acid. The extract of *Eclipta Alba* has the potent inhibitor of the pro-inflammatory transcription factors and a promising agent for the treatment of the inflammatory cascade of cardiovascular diseases (Kaileh et al., 2007).

Antidiabetic Activity : Extracts of various plant materials capable of decreasing blood sugar (Bopanna K et al., 1982). The chloroform extract of *Eclipta Alba* exhibited significant

antidiabetic activity in alloxan

induced diabetic rats. This extract has showed

Figure 5:

122 [J.Venom.Anim. Toxins incl.Trop. Dis ()] , *J.Venom.Anim. Toxins incl.Trop. Dis* 2003. 9 (2) p. 381.

123 [Songklanakarin J. Sci. Technol ()] , *Songklanakarin J. Sci. Technol* 2006. 28 (4) p. .

124 [Sawant et al. ()] 'Analgesic studies on total alkaloids and alcohol extracts of *Eclipta alba* (Linn.) Hassk'. Mahesh
125 Sawant , C Jolly , Shridhar Isaac , Narayanan . *Phytotherapy research* 2004. 18 (2) p. .

126 [Tewtrakul et al.] *Anti-HIV-1 integrase activity of medicinal plants used as self medication by AIDS patients*,
127 Supinya Tewtrakul , Sanan Subhadhirasakul , Sopa Kum mee .

128 [Arunachalam et al. ()] 'Anti-inflammatory activity of methanolic extract of *Eclipta prostrata* L. (Asteraceae)'.
129 G Arunachalam , N Subramanian , G P Pazhani , V Ravichandran . *African Journal of Pharmacy and*
130 *Pharmacology* 2009. 3 (3) p. .

131 [Gupta et al. ()] 'Anticancer Activity of *Indigofera aspalathoides* and *Wedelia calendulacea* in Swiss Albino
132 Mice'. Malaya Gupta , K Upal , Palla K Anti Mazumder , Haldar , C Chandi , Laxmanan Kandar , G P
133 Manikanda , Senthil . *Iranian journal of pharmaceutical research* 2005. 6 (2) p. .

134 [Khanna ()] 'Anticancer-cytotoxic activity of saponins isolated from the leaves of *Gymnema sylvestre* and *Eclipta*
135 *alba* on HeLa cells'. Kannabiran Khanna . *International journal of green pharmacy* 2008. 1 p. .

136 [Ananthi et al. ()] 'Antihyperglycemic Activity of *Eclipta alba* Leaf on Alloxan-induced Diabetic Rats'. J Ananthi
137 , A Prakasam , K V Pugalendi . *Yale Journal of Biology and Medicine* 2003. 76 p. .

138 [Singh et al. ()] 'Antiinflammatory and analgesic agents from Indian medicinal plants'. Amritpal Singh , Ravi
139 Samir Malhothra , Subban . *International journal of integrative biology* 2008. 3 (1) p. .

140 [Vianna-Da-Silva et al.] *Antivenom effect of a new synthetic coumestan analog of wedelolactone*, N M Vianna-
141 Da-Silva , Moraes Da Silva , Ajm Costa , Prr Melo , PA .

142 [Jadhav et al. ()] 'Chemical composition, pharmacological activities of *Eclipta alba*'. V M Jadhav , R M Thorat
143 , V J Kadam , K P Salaskar . *Journal of Pharmacy Research* 2009. 2 (8) p. .

144 [Dan and Gamble ()] *Chinese Herbal Medicine: Materia Medica*, Bensky Dan , Andrew Gamble . 1986. Seattle:
145 Eastland Press.

146 [Sahu and Pankaj ()] 'Clinical Study of Pilex Combination Therapy Vs Conventional Ayurvedic Therapy in the
147 Management of Haemorrhoids'. M Sahu , Srivastava Pankaj . *The Indian Practitioner* 2001. 54 (11) p. .

148 [Meena et al. ()] 'Comparative evaluation of standardisation parameters between *Wedelia* genus species'. A K
149 Meena , M M Rao , Komalpreet Kaur , P Panda . *Int J. Ph Sci and Res* 2010. 1 (3) p. .

150 [Lans ()] 'Comparison of plants used for skin and stomach problems in Trinidad and Tobago with Asian
151 ethnomedicine'. Cheryl Lans . *Journal of Ethnobiology and Ethnomedicine* 2007. 3 (3) p. .

152 [Wagner ()] 'Coumestans as the main active principles of the liver drugs *Eclipta alba* and *Wedelia Calendulacea*'.
153 H Wagner . *Planta Med* 1986. 5 p. .

154 [Maged and Abdel-Kader ()] 'DNA damaging steroidal alkaloids from *Eclipta alba* from the suriname rainforest'.
155 S Maged , Abdel-Kader . *J. Nat. Prod* 1998. 61 (10) p. .

156 [Upadhyay et al. ()] 'Eclalbatin, a triterpene saponin from *Eclipta alba*'. R K Upadhyay , M B Pandey , R N
157 Jha , V B Pandey . *J Asian Nat Prod Res* 2001. 3 (3) p. .

158 [Singh et al. ()] *Eclipta Alba Linn. -Ancient remedy with therapeutic potential*, Amritpal Singh , Sanjiv Duggal ,
159 Asish Suttee , Jaswinder Singh , Shankar Katekhaye . 2010. 1 p. .

160 [Xiong-Hao Lin et al. ()] *Effects of Volatile Com-ponents and Ethanolic Extract from Eclipta prostrata on*
161 *Proliferation and Differentiation of Primary Osteoblasts*, Yan-Bin Xiong-Hao Lin , Shan Wu , Jian-Wei
162 Lin , Pei-Yuan Zeng , Jin-Zhong Zen , Wu . 2010. 15 p. .

163 [Abdul Viqar Khan ()] 'Ethnomedicinal uses of *Eclipta prostrata*. Indian journal of traditional knowledge'. Abdul
164 Viqar Khan . *Pharmacologyonline* 2008. 2011. 7 (2) p. .

165 [Rupali Thorat et al. ()] *Evaluation of a herbal hair oil in reducing hair fall in human volunteers*, Rupali Thorat
166 , Vilasrao Varsha Jadhav , Kadam . 2009. 6 p. . (Ninad sathe, Ashwini save, Vikas ghorpade)

167 [Bhaskar Rao et al. ()] *Evaluation of antioxidant potential of clitoria ternate L. and Eclipta prostrata L. Indian*
168 *journal of biochemistry and biophysics*, D Bhaskar Rao , Ch Ravi Kiran , Y Madhavi , P Koteshwara Rao ,
169 Raghava Rao , T . 2009. 46 p. .

170 [Kima et al. ()] 'Gun-Soon Lee. The butanol fraction of *Eclipta prostrata* (Linn) effectively reduces serum lipid
171 levels and improves antioxidant activities in CD rats'. Dae-Ik Kima , Sung-Hyen Lee , Jin-Ho Choia , Hyun
172 Soon Lillehoj , Mi-Hee Yu . *Nutrition Research* 2008. 28 p. .

173 [Roy et al. ()] 'Hair growth promoting activity of *Eclipta alba* in male albino rats'. R K Roy , Mayank Thakur ,
174 V K Dixit . *Arch Dermatol Res* 2008. 300 p. .

175 [Bruce Milliman et al. ()] 'Hepatitis C: A Retrospective Study, Literature Review, and Naturopathic Protocol'.
176 W Bruce Milliman , Davis W Lamson , MS , Matthew Brignall , S . *Alternative Medicine Review* 2000. 5 (4)
177 p. .

178 [Samudram et al. ()] 'Hepatoprotective activity of Biherbal ethanolic extract on CCl₄ induced hepatic damage
179 in rats'. P Samudram , Rajeshwari Hari , R Vasuki , Geetha , Sathiya . *Clinical nutrition insights* 2008. 1998.
180 2 (2) p. . (AJBR)

181 [Saxena et al. ()] 'Hepatoprotective effects of Eclipta alba on subcellular levels in rats'. A K Saxena , B Singh ,
182 K K Anand . *Journal of ethnopharmacology* 1993. 40 (3) p. .

183 [Tewtrakul et al. ()] 'HIV-1 protease and HIV-1 integrase inhibitory substances from Eclipta prostrata'. S
184 Tewtrakul , S Subhadhirasakul , S Cheenpracha , C Karalai . *Phytother Res* 2007. 21 (11) p. .

185 [Dhandapani ()] *Hypolipidemic activity of Eclipta prostrata (L.) L. leaf extract in atherogenic diet induced*
186 *hyperlipidemic rats*, R Dhandapani . 2007. 45 p. .

187 [Kaushik-Basu et al. ()] *Identification and characterization of coumestans as novel HCV NS5B polymerase*
188 *inhibitors*, Neerja Kaushik-Basu , Alain Bopda Waffo , Tanaji Talele , T , Amartya Basu , Paulo Costa
189 R R , Alcides Da Silva , JM , Stefan Sarafianos , G , Francois Noel . 2008. 36 p. .

190 [Banji et al. ()] 'Investigation on the effect of Eclipta alba on animal models of learning and memory'. Otilia
191 Banji , David Banji , Annamalai Ar , R Manavalan . *Indian J physiol pharmacol* 2007. 51 (3) p. .

192 [Sagar et al. ()] *Invitro Production of Anti-hepatotoxic Compounds in Cultures of Eclipta alba Linn. and Silybum*
193 *mariannum gaertn*, B P Sagar , A Sangwan , A Panwar , A Sangwan . 2006. 6 p. 88.

194 [Lal et al. ()] 'Kuldeep Singh Yadav. Screening of Leaves and Roots of Eclipta alba for Hepatoprotective Activity'.
195 V K Lal , Amit Kumar , Prashant Kumar . *Arch. Appl. Sci. Res* 2010. 2 (1) p. .

196 [Uddin et al. ()] 'Masudul Azad Chowdhury AM. Antioxidant, cytotoxic and antimicrobial properties of Eclipta
197 alba ethanolic extract'. Nazim Uddin , Atiar Rahman , Sohel Nazim Ahmed , Rasheda Rana , Akter .
198 *Int J Biol Med Res* 2010. 1 (4) p. .

199 [Thakur and Mengi ()] 'Neuropharmacological profile of Eclipta alba (Linn.) Hassk'. V D Thakur , S A Mengi .
200 *Journal of Ethnopharmacology* 2005. 102 p. .

201 [Shoji Yahara et al. ()] 'Oleanane glycosides from Eclipta alba'. Shoji Yahara , Toshihiro Ning Ding , Nohara .
202 *Chem.pharm.bull* 1994. 42 (6) p. .

203 [Mohammad S Rahman et al. ()] *Oleanane Glycosides from Eclipta prostrata. The Dhaka university journal of*
204 *pharmaceutical sciences*, Rasheduzzaman Mohammad S Rahman , Chowdhury , M Chowdhury , Hasan , A
205 Mohammad , Rashid . 2005. 4.

206 [Christyapita et al. ()] *Oral administration of Eclipta alba leaf aqueous extract enhances the non-specific immune*
207 *responses and disease resistance of Oreochromis mossambicus. Fish and shellfish immunology*, D Christyapita
208 , M Divyagnaneswari , Dinakaran Michael , R . 2007. 23 p. .

209 [Dalal et al. ()] 'Phytochemical Screening of Methanolic Extract and Antibacterial Activity of Active Principles
210 of Hepatoprotective Herb'. Sunita Dalal , Sudhir Kataria , K Sastry , K Rana , Svs . *Eclipta alba.*
211 *Ethnobotanical Leaflets* 2010. 14 p. .

212 [Mukherjee and Poddar ()] *Pilex therapy in Piles -a Preliminary Report. The antiseptic*, D R Mukherjee , H
213 Poddar . 1976. 10 p. 541.

214 [Mitesh Phale et al. ()] 'Precise and sensitive HPTLC method for quantitative estimation of wedelolactone in
215 Eclipta alba Hassk'. D Mitesh Phale , Purnima Hamrapurkar , D , Manasi Chachad , A , Priti Patil , S ,
216 Sandeep Pawar , B . *Pharmacophore* 2010. 1 (2) p. .

217 [Jayathirthaa and Mishraa ()] 'Preliminary immunomodulatory activities of methanol extracts of Eclipta alba
218 and Centella asiatica'. M G Jayathirthaa , S H Mishraa . *Phytomedicine* 2004. 11 (4) p. .

219 [Razdan et al. ()] 'Preventive and curative effects of Vedic Guard against antitubercular drugs induced hepatic
220 damage in rats'. Rema Razdan , Amar Imranulla , M J Dev . *Phcog mag* 2008. 4 (15) p. .

221 [Lee et al. ()] *Sang Hyun Sung, Gun Hee Kim, Young Choong Kim. Antiproliferative activity of triterpenoids from*
222 *Eclipta prostrata on hepatic stellate cells*, Mi Kyeong Lee , Na Ry Ha , Hyekyung Yang . 2008. 15 p. .

223 [Karthikumar et al. ()] 'Screening of antibacterial and antioxidant activities of leaves of Eclipta prostrata (L)'.
224 S Karthikumar , K Vigneswari , K Jegatheesan . *Sci. Res. Essays* 2007. 2 (4) p. .

225 [Rolf Teschke and Bahre ()] *Severe hepatotoxicity by Indian herbal products: A structured causality assessment*,
226 Ruediger Rolf Teschke , Bahre . 2009. 8 p. .

227 [Hemalatha et al. ()] *Shrivijaya kirubha T. Evaluation of antidiabetic and diuretic activity of polyherbal formu-*
228 *lation. Indian Journal of Traditional Knowledge*, S Hemalatha , T Ayyappan , S Shanmugan , D Nagavalli .
229 2006. 5 p. .

230 [Baig and Bhagwat ()] *Study the efficacy of Galactin Vet Bolus on milk yield in dairy cows. Veterinary world*,
231 M I Baig , V G Bhagwat . 2009. 2 p. .

232 [Sawangjaroen et al. ()] 'The in vitro anti-giardial activity of extracts from plants that are used for self-
233 medication by AIDS patients in southern Thailand'. N Sawangjaroen , S Subhadhirasakul , S Phongpaichit ,
234 C Siripanth , K Jamjaroen , K Sawangjaroen . *Parasitol Res* 2005. 95 (1) p. .

235 [Haveliwala ()] *The role of 'Sapera Forte Tablet' in the management of hypertension*, M D Haveliwala . 1963. 9

236 p. .

237 [Shoji Yahara and Ding ()] 'Toshihiro nohara, Kazuo mazuda, Hiroyuki agenta. Taraxastane glycosides from

238 'Eclipta alba'. Ning Shoji Yahara , Ding . *Phytochemistry* 1997. 40 (1) p. .