

1 Early Suspicion of Vasa Previa with Velamentous Umbilical Cord 2 Insertion and Low Laying Placenta and its Management

3 Natasha Gupta MD¹ and Natasha Gupta MD²

4 ¹ Mount Sinai Hospital

5 *Received: 11 December 2012 Accepted: 1 January 2013 Published: 15 January 2013*

7 Abstract

8 We present a case of a 34 years old female who was suspected to have vasa previa due to
9 sonographic evidence of low lying placenta and velamentous cord insertion, early in the
10 pregnancy. Vasa previa can lead to significant perinatal morbidity by causing fetal
11 exsanguination, if diagnosis is delayed. This case demonstrates the need for early suspicion of
12 vasa previa in presence of persistent high risk features on repeated ultrasounds, importance of
13 patient preparation with antenatal steroids as well as comprehensive education of patient
14 regarding the warning signs and symptoms.

16 *Index terms*—

17 Early Suspicion of Vasa Previa with Velamentous Umbilical Cord Insertion and aying Placenta and its
18 Management Natasha Gupta Md ? , Mina Tirabassi Rdms ? , Jeff B Chapa Md ? , Ori Kushnir Md ? &
19 Josef Blankstein Md ¥ Abstract -We present a case of a 34 years old female who was suspected to have vasa
20 previa due to sonographic evidence of low lying placenta and velamentous cord insertion, early in the pregnancy.
21 Vasa previa can lead to significant perinatal morbidity by causing fetal exsanguination, if diagnosis is delayed.
22 This case demonstrates the need for early suspicion of vasa previa in presence of persistent high risk features
23 on repeated ultrasounds, importance of patient preparation with antenatal steroids as well as comprehensive
24 education of patient regarding the warning signs and symptoms.

25 1 I.

26 2 Case Presentation

27 34-years old female, gravida 2, para 1, presented at 32.5 weeks gestation with complaints of contractions every
28 2 minutes. She denied any vaginal bleeding or leakage of fluid. She was being closely followed for low lying
29 placenta and velamentous insertion of umbilical cord, with serial ultrasounds. Her sonogram at 20.2 weeks
30 showed an anterior, low lying placenta. A follow up ultrasound at 24.2 weeks noted anterior, marginal placenta.
31 A transvaginal sonogram at 29.5 weeks revealed anterior marginal placenta with suspicion for funic presentation
32 (FIGURE A). Color flow mapping was suggestive of velamentous cord insertion with suspicion for vasa previa
33 (FIGURE B).

34 Patient was administered a course of steroids in preparation for preterm delivery for suspected vasa previa.
35 She also received instructions to return to hospital in case of leakage of fluid or vaginal bleeding. An ultrasound at
36 31.2 weeks again noted velamentous cord insertion, with vessels seen coursing the membranes in close proximity
37 of internal cervical os, thus providing strong suspicion for previa. Patient returned to labor and delivery with
38 frequent contractions and was delivered with Cesarean section to prevent fetal head compression of vessels or
39 their rupture from uterine contractions. Placental inspection confirmed the diagnosis of velamentous insertion
40 and vasa previa. Infant with APGAR scores of 7 and 7 at 1 and 5 minutes respectively was born. Baby was
41 admitted to neonatal intensive care unit (NICU) for further care, did not require transfusion and was discharged
42 after 15 days stay in NICU. Mother and baby continued to do well on their follow up.

3 II.

4 Discussion

45 Vasa Previa is an uncommon obstetric complication where fetal blood vessels travel the membranes covering the
46 internal cervical os, in front of the fetal presenting part [1]. Type 1 vasa previa is associated with velamentous
47 umbilical cord insertion, where cord is inserted marginally into the placenta and lacks the protective covering
48 of Wharton's jelly around it, such that umbilical vessels are covered only by the membranes [1]. These vessels
49 are prone to rupture and compression with onset of labor and with rupture of membranes. Type 2 vasa previa
50 is seen in association with bilobed placenta or placenta with a succenturiate lobe, where fetal vessels travel the
51 membranes connecting 2 lobes of placenta or those connecting placenta with its accessory or succenturiate lobe.
52 The incidence of vasa previa is 1 in 2500 in United States.

53 Risk factors for vasa previa include placental and cord abnormalities like velamentous insertion of umbilical
54 cord [2], bilobed placenta, placenta with succenturiate or accessory lobe, low-lying placenta or placenta previa
55 as well as multifetal pregnancies and pregnancies following in vitro fertilization [3]. It is essential to identify
56 vasa previa in a timely manner since onset of labor or premature rupture of membranes results in rapid fetal
57 exsanguination causing fetal anemia, hypotension and demise within doppler can be reliably used for umbilical
58 cord and placental pathologies and has a very low false positive rate. Also, transvaginal approach is considered
59 superior to the transabdominal approach, which may not be practical in obese patients, those with abdominal
60 scars or difficult fetal presentations [3]. Rarely, transvaginal ultrasound may pose challenges in diagnosis due to
61 motion artifacts or when a funic presentation is misinterpreted as vasa previa [3]. Second trimester is considered
62 the best time for antenatal screening for vasa previa [7].

63 Several studies have compared the neonatal outcomes in vasa previa recognized prenatally with those recognized
64 intrapartum and they described significantly better survival rates, improved APGAR scores, shorter NICU stay
65 and lower transfusion rates in those diagnosed prenatally. Perinatal mortality up to 56% is noted if vasa previa
66 remains undiagnosed prenatally, compared to 3% in those diagnosed prenatally. Similarly, neonatal blood
67 transfusion rates of 58.5% is reported in those that remain undiagnosed prenatally versus 3.4% in those
68 diagnosed prenatally [8][9][10].

69 Optimal management of vasa previa consists of patient education regarding signs of labor, early administration
70 of a course of steroids, hospitalization at 31-32 weeks for fetal monitoring and cesarean delivery at 34 weeks or
71 before, if labor ensues or if fetal wellbeing is compromised by membrane rupture or cord compression. ??obinson
72 () E insertion site of umbilical cord using color doppler. This revealed velamentous insertion and a close follow
73 up with serial scans suggested vasa previa. Cipriano et al studied the cost effectiveness of a screening ultrasound
74 for vasa previa in all twin gestations and in-vitro fertilization pregnancies and concluded that these screening
75 ultrasounds are cost effective [6]. Catanzarite et al followed 11 cases of vasa previa diagnosed by color doppler
76 sonography, of which 10 were confirmed to have vasa previa upon delivery by the delivering physician, thus
77 noting a 91% specificity of the sonographic diagnosis of vasa previa [1]. Thus, color minutes [3]. Due to lethal
78 nature of this rare condition, several authors recommend early prenatal diagnosis of vasa previa by using color flow
79 mapping in second trimester, especially in patients with risk factors [4]. Color flow imaging is popularly employed
80 for the purpose of identifying umbilical cord and placental pathologies as compared to Gray scale imaging alone.
81 Nomiya et al reported 100% sensitivity of color Doppler imaging in identifying velamentous cord insertion,
82 when performed routinely between 18-20 weeks gestation [5]. They proceeded to identify vasa previa in the
83 patients that were noted to have velamentous cord insertion on color doppler imaging and they concluded that
84 color doppler had 100% sensitivity, 99.8% specificity, 83% positive predictive value and 100% negative predictive
85 value in identification of velamentous cord insertion. We applied same investigational technique in our patient
86 when we noted a low lying placenta on a second trimester scan, following which careful attention was paid to
87 placental outcomes [11]. Thus, most authors recommend a scheduled delivery at 34 or 35 weeks in the patients
88 that do not go into labor or rupture spontaneously [11]. Chmait et al described a patient with type 2 vasa previa
89 that they treated with fetoscopic laser ablation, who delivered at 33.3 weeks with a good perinatal outcome
90 [12,13].

91 Our patient was administered a course of steroids in anticipation of preterm delivery due to suspected vasa
92 previa. She was also educated about this condition and about warning signs and symptoms suggestive of onset
93 of labor. She underwent an uncomplicated cesarean section which confirmed our diagnosis of vasa previa and
94 there were no neonatal sequelae due to vasa previa or preterm delivery.

95 5 III.

96 6 Conclusion

97 ? Vasa Previa is a condition where fetal vessels travel unprotected in front of the cervical os and are at risk of
98 rupture with the rupture of membranes or at risk of compression from the fetal head when the labor ensues.

99 ? It is commonly associated with velamentous insertion of umbilical cord or low lying placenta and other
100 placental abnormalities. ? A screening ultrasound in second trimester can detect these risk factors. Patients
101 noted to have these risk factors on ultrasound should be followed with color doppler sonography to diagnose vasa
102 previa.

103 ? If vasa previa is suspected, patient should be educated about this condition and counseled to return to hospital in case of rupture of membranes or onset of labor. ^{1 2}



Figure 1:

104

¹© 2013 Global Journals Inc. (US)

²© 2013 Global Journals Inc. (US) © 2013 Global Journals Inc. (US)

6 CONCLUSION




Figure 2: Figure A :




Figure 3:

013
2
Year

Figure 4:

Figure 5:

-
- 105 [Derbala et al. ()] , Y Derbala , F Grochal , P Jeanty , Vasa Previa . *J Prenat Med* 2007. 1 (1) p. .
- 106 [Hasegawa ()] ‘Analysis of the ultrasonographic findings predictive of vasa previa’. J Hasegawa . *Prenat Diagn*
107 2010. 30 p. .
- 108 [Nomiya et al. ()] ‘Antenatal diagnosis of velamentous umbilical cord insertion and vasa previa with color
109 Doppler imaging’. M Nomiya , Y Toyota , H Kawano . *Ultrasound Obstet Gynecol* 1998. 12 (6) p. .
- 110 [Robinson and Grobman ()] ‘Effectiveness of timing strategies for delivery of individuals with vasa previa’. B K
111 Robinson , W A Grobman . *Obstet Gynecol* 2011. 117 (3) p. .
- 112 [Gagnon ()] ‘Guidelines for the management of vasa previa’. R Gagnon . *J Obstet Gynaecol Can* 2009. 31 (8) p. .
- 113 [Quintero ()] ‘in utero laser treatment of type II vasa previa’. R A Quintero . *J Matern Fetal Neonatal Med* 2007.
114 20 (12) p. .
- 115 [Smorgick ()] ‘Is neonatal risk from vasa previa preventable? The 20-year experience from a single medical
116 center’. N Smorgick . *J Clin Ultrasound* 2010. 38 (3) p. .
- 117 [Kanda ()] ‘Prenatal diagnosis and management of vasa previa: a 6-year review’. E Kanda . *J Obstet Gynaecol*
118 *Res* 2011. 37 (10) p. .
- 119 [Catanzarite ()] ‘Prenatal sonographic diagnosis of vasa previa: ultrasound findings and obstetric outcome in ten
120 cases’. V Catanzarite . *Ultrasound Obstet Gynecol* 2001. 18 (2) p. .
- 121 [Cipriano et al. ()] ‘The cost-effectiveness of targeted or universal screening for vasa praevia at 18-20 weeks of
122 gestation in Ontario’. L E Cipriano , W H Barth , Jr , G S Zaric . *BJOG* 2010. 117 (9) p. .
- 123 [Chmait ()] ‘Third trimester fetoscopic laser ablation of type II vasa previa’. R H Chmait . *J Matern Fetal*
124 *Neonatal Med* 2010. 23 (5) p. .
- 125 [Lijoi and Brady ()] ‘Vasa previa diagnosis and management’. A F Lijoi , J Brady . *J Am Board Fam Pract* 2003.
126 16 (6) p. .
- 127 [Lee ()] ‘Vasa previa: prenatal diagnosis, natural evolution, and clinical outcome’. W Lee . *Obstet Gynecol* 2000.
128 95 (4) p. .