

Breast Conserving Surgery and Whole Breast Radiation therapy Followed by High Dose Rate Brachytherapy Boost Versus Electron Beam Boost in the Treatment of Early Breast Cancer in Young Indian Women: Which is Cosmetically Better?

Dr. Aseem Rai Bhatnagar¹, Dr. Aseem Rai Bhatnagar² and Dr. Rameshwaram
Sharma³

¹ S.M.S Medical college, Jaipur, India.

Received: 11 April 2013 Accepted: 1 May 2013 Published: 15 May 2013

Abstract

Breast conserving treatment (BCT) is the treatment of choice in early breast cancer. Despite years of observation it is still regarded as controversial not in regard to the very idea; the controversy pertains rather to the way it is being performed by radiation oncologists and surgeons. There are no uniform indications as far as the optimal surgery range is concerned (lumpectomy alone, lumpectomy with the macroscopic margin of 1cm, excision of the breast tissue block of a segment or a quadrant). BCT has produced survival equivalent to mastectomy in the treatment of patients with early-stage invasive breast carcinoma in several randomized Phase III clinical trials. Breast irradiation is an essential element of the conservative approach. Local recurrence risk after surgery alone reaches 35

Index terms— Breast conserving treatment (BCT).

Background reast conserving treatment (BCT) is the treatment of choice in early breast cancer. Despite years of observation it is still regarded as controversial not in regard to the very idea; the controversy pertains rather to the way it is being performed by radiation oncologists and surgeons. There are no uniform indications as far as the optimal surgery range is concerned (lumpectomy alone, lumpectomy with the macroscopic margin of 1cm, excision of the breast tissue block of a segment or a quadrant). BCT has produced survival equivalent to mastectomy in the treatment of patients with early-stage invasive breast carcinoma in several randomized Phase III clinical trials.

Breast irradiation is an essential element of the conservative approach. Local recurrence risk after surgery alone reaches 35%, compared to 10% in patients undergoing adjuvant radiotherapy [1]. First, the whole breast is irradiated using external beam technique, usually with a dose of 50Gy. Subsequently it is necessary to increase the dose delivered to the tumour bed using a so called "boost".

Primary boost dose methods include teleradiotherapy (TRT) with external photon or electron beam (usually) and high dose rate (HDR) or low dose rate (LDR) brachytherapy (BT) [2]. The cost and time required (for both the patient and physician) for these two boosting techniques differ greatly. Whole-breast EBRT involves a 6-week course of fractionated treatments.

In contrast, BT can be completed in a 4-to 5day treatment course. In addition, BT adds the risk of an invasive procedure with an outcome that is highly dependent upon the expertise of the physician/physicist team. Biomathematical models are often used to estimate equivalent high-dose-rate regimens. For example, linear quadratic modelling has suggested that a high-dose-rate regimen of 5 fractions of 310 cGy per fraction should approximate the early and late effects of a 20-Gy low dose rate delivered at 0.5 Gy/h. Although biomathematical models can be used to estimate the appropriate dose, there is no standardized high-dose-rate fractionation schedule that can be recommended [3,4,5].

3 A) REVIEW OF LITERATURE

In two studies the efficacy of HDR BT and TRT as a boost in non-advanced breast cancer patients with breast conserving treatment was compared. First, whole breast irradiation was performed using an external photon beam (50Gy in classical fractionation). Subsequently Hammer et al. delivered a boost to the tumour bed using either an electron beam (TRT-11Gy in 5 fractions) or HDR BT (single 10Gy boost). Local recurrence rates were 8.2% and 4.3% ($p < 0.04$), respectively. Excellent or good cosmetic results were achieved in 70% and 88%, respectively ($p < 0.0001$) [6].

Polgar et al., in a randomized clinical Phase III trial, after the first stage of the study randomized the patients into 2 groups. In the first group the patients received external electron beam therapy of 16Gy in 8 fractions. In the second group the same total dose was delivered in the form of HDR BT. Local recurrence rates were 6% and 8.5%, respectively. Excellent or good cosmetic results were achieved in 83% and 88%, respectively [7]. The differences between rates in the two groups were not statistically significant.

Kulik from the Oncology Centre in Warsaw presented the results of a HDR BT boost study in 93 patients undergoing conservative treatment. During the 3-year follow-up one case of local recurrence was observed; excellent or good cosmetic results were achieved in 85% of patients. In a ProbRough rule induction analysis including all clinical and therapeutic variables it was shown that patients with a mammography diameter of tumour not exceeding 11mm have the best chance of excellent or good cosmetic results [8].

We undertook this study, the second such study in our Department, to evaluate the effect of HDR BT boost versus electron beam boost on local tumor control, side effects and cosmesis after breast conserving surgery in early breast cancer II.

Methods 40 patients with invasive early-stage breast cancer (Stage I-II as defined by the AJCC 7th edition guidelines) were treated prospectively with breast conservation surgery. All patients signed informed consent forms prior to treatment. All the patients underwent tumorectomy i.e. macroscopic total resection of the primary tumor. Re-excision to achieve negative surgical margins was performed as needed to obtain margins of 2 mm, if initially the surgical margin was positive. All patients underwent full axillary lymph node dissection (all III levels of the axillary fossa). The median number of lymph nodes excised was 16.

In all patients, adjuvant EBRT to the whole breast was used. Patients were positioned supinely on a breast board with both arms raised overhead. 3D CT planning of the breast was used. Patients were treated with two tangential fields with either gamma-rays from a cobalt unit or with 4-6 MV photon X-rays. Whole breast radiotherapy was delivered as 50 Gy in 25 fractions over 5 weeks.

Boost to the tumor bed was given to an equivalent dose of 15-16 Gy with either HDR BT using Iridium-192 interstitial temporary implants or electron beam using a linear accelerator. Electron beam boost was given in continuation with EBRT to maintain the continuity. There was one week gap between completion of EBRT and HDR BT boost to reduce chances of infection. In the HDR BT boost group, implants were designed to irradiate the lumpectomy cavity with at least a 1-2 cm margin. The dose rate was 350 cGy twice a day for two days. In the electron group the boost was 250 cGy once daily with 9-12 MeV electron over 6 days.

The toxicities and cosmesis were assessed at a specific time point: at 1.5 years of follow-up. The toxicity parameters examined included the following: breast edema, erythema, fibrosis, hyperpigmentation, hypopigmentation, breast pain, breast infection, telangiectasia, and fat necrosis. Toxicities were graded by using the Radiation Therapy Oncology Group (RTOG) / European Organization for Research & Training of Cancer (EO-RTC) late radiation morbidity scoring scheme and Common Terminology Criteria for Adverse Events (CTC-AE) for skin, subcutaneous tissues, pain and dermatitis. Breast edema, erythema, pigmentary changes, and telangiectasia fell under the domain of radiation dermatitis and skin; breast fibrosis and breast pain were under the domains of subcutaneous tissues and pain due to radiation, respectively. Breast infections and fat necrosis were either present or not and were noted accordingly.

In accordance with the guidelines of Common Toxicity Criteria, version 4.0, toxicities were graded by using the acute/chronic radiation morbidity scale: Grade 0 - no observable radiation effects; Grade 1 : mild radiation effects; Grade 2 : moderate radiation effects; Grade 3 : severe radiation effects.

Cosmetic evaluation was based on the standards set forth by the Harvard criteria as shown below in Table ?? . The treating physician at a scheduled follow-up visit scored the cosmetic result. No patient-reported scoring of cosmetic outcome was done. Likewise, the treating radiation oncologist did all toxicity scoring for each patient. ()I

The statistical method employed for the incidence/severity of toxicities and cosmetic outcome with various parameters was Pearson chi-square analysis stratified for no toxicity versus any toxicity.

1 III.

2 Results and Discussion

3 a) Review of Literature

Oedema of the breast, hyperpigmentation, hypopigmentation/ depigmentation of the nipple and papillae, telangiectases and fibrosis are all consequences of radiation therapy [10]. Generally breast pain, edema, erythema, and hyperpigmentation all diminish in frequency over time. Edema of the breast is observed mainly

during and directly the end of radiot-herapy. In 10-20% of patients, it can appear as a late reaction after 18-36 months after radiotherapy; in such cases it is moderate and reversible [10].

Sequelae that increases until the 2-year mark and later stabilizes includes breast fibrosis and hypopigmentation. Fat necrosis and telangiectasia increase with the passage of time. A study from Peter Y. Chen et al showed fat necrosis increased from 1% at 6 months to 9% at 2 years and 11% at 5 years. The median time to occurrence of fat necrosis was 5.5 years after completion of radiation therapy and HDR BT.

Telangiectases are observed mainly in areas of high doses of radiotherapy given by electrons or HDR BT or in areas of skin folds. They can be observed in 30% of patients and time to their appearance is the longest out of all side effects of radiotherapy. Contrary to other side effects, the probability and intensity of telangiectases increases in the course of follow-up. The most important late effect of radiation is breast fibrosis. Contrary to other factors, which are reversible (oedema) or limited to a small area of the breast (telangiectases), fibrosis encompasses the whole breast and is the most important factor of breast's retraction [10]. Fibrosis appears after 6-18 months and the highest intensity is observed after 3 years. Longer observations of patients did not reveal progression of the retraction of the treated breast. It is advised to perform cosmetic evaluation 3 years after primary treatment because at this point most late effects already appear. Late effects, those that appear years after, don't affect final cosmesis. In our study we evaluated the cosmetic outcomes and adverse events at 1.5 years after the completion of whole breast EBRT and boost. So we did not evaluate the changing trends for these events and as such our study follow up was short. For the implant group, nearly all pigmentary changes, whether hyperpigmentation or hypopig-ment-ation, were pinpoint rather than diffuse, corresp-onding to the sites where the HDR catheters were been placed. E-xcellent or good cosmetic results were achi-eved in 10-0% and 90% patients of electron boost and HDR BT Boost respectively ($p=0.0009$), as shown below in table ?? . In assessing toxicities and cosmesis, interpretations of changes like fibrosis and cosmetic outcomes are not entirely objective. In our study, grading of fibrosis was based on the degree of induration palpated at the time of each follow-up visit. Because induration dimesions were not always recorded, the grading by the examining clinician became the basis on which the degree of fibrosis was assessed. Fibrotic changes can be difficult to differentiate between sequelae from postsurgical changes and sequelae from radiation effects. Reexcision, such second surgical procedure, also contributed to breast fibrosis/induration. Thus, fibrosis is a conti-num and a morbidity of both surgical excision and a late radiation effect. It would be difficult to determine the proportional contribution of surgery versus the contribution of radiation that leads to fibrosis. However, we conservatively assigned any degree of induration under subcutaneous tissue-late RT morbidity scoring (fibrosis) solely related to a late radiation sequelae.

4 Volume XIII Issue V Version I

There was no significant difference in local tumor control between patients treated with electron bosst or HDR BT boost over a period of one and a half year in our study. The rate of local recurrence was same between the 2 patient groups: The HDR BT group demonstrated a local recurrence rate of 5% compared with patients who received electron beam boost, who had a similar 5% risk of local failure ($p=1.00$).

5 IV.

6 Conclusions

Breast conservation therapy nowadays is an effective treatment for early breast cancer with more and more patients preferring this option due to better psychosexual quality of life. Breast conserving therapy in patients with early breast cancer allows us to achieve an excellent and good (satisfactory) cosmetic effect in a majority of cases (95% in our study). The results of the quailtative cosmetic evaluation vary between the patients and the physicians. We have done two such Volume XIII Issue V Version I Year 013 2 () studies to address cosmesis in BCT in our Department. In one of our studies, patients with early breast cancer after undergoing breast conserving surgery and whole breast irradiation have better cosmetic results and reduced chances of fibrosis at one and a half years of follow-up, when they are given electron boost as compared to HDR BT boost. Local tumor control rates were similar between the two groups. For local tumor control assessment long term follow up studies are needed.

Reaching an unequivocal opinion on which of the two boost techniques, TRT boost with electrons or HDR BT, is more efficient is not an easy task. Hammer et al. showed significantly lower local recurrence rates with significantly higher rate of excellent and good cosmetic results for the HDR BT, while the group from the National Oncology Institute in Budapest did not confirm these results in the settings of a randomized study [6,7].

It has been stated that publications showing inferior cosmetic outcomes after brachytherapy boost have lacked the necessary attention to technical details such as dose homogeneity. But this was not seen in our study. Recent experiences have demonstrated equivalent or superior results for HDR BT as compared to electron-beam boosting- despite the higher doses. Irrespective of the dose rate (HDR or LDR) better cosmetic results by BT boost can be explained by the lower dose delivered to the skin. This results from the fact that the distance between the most "superficial" interstitial guide needle and the skin should reach 5mm. Thus the danger of teleangiectasias and fibrosis, which significantly influences cosmetic outcomes, is reduced. Due to the beam geometry this cannot be achieved using electron beam TRT [9]. So the debate as to which is the optimal boosting technique goes on.

V.

163 7 Bibliography



Figure 1:

[Note: I.]

Figure 2:

164 1

Figure 3:

2

	HDR BT Boost (% of patients)			Electron Beam Boost (% of patients)			p-value
	Grade 1	Grade 2	Grade 3	Grade 1	Grade 2	Grade 3	
Breast Pain	15		5 0	15	0	0	0.211
Breast edema	15		0 0	10	0	0	0.161
Erythema	15		0 0	15	0	0	1.000
Hyperpigmentation	40		5 0	30	0	0	0.002
Hypopigmentation	35		0 0	20	0	0	0.001
Fibrosis	50		5 5	30	0	0	<0.001
Telangiectasia	25		5 0	20			0.029
Fat necrosis	10			5			0.095
Breast Infection	5			0			0.021

[Note: *Fat necrosis and infection are not graded.]

Figure 4: Table 2 :

-
- 165 [Int J Radiat Oncol Biol Phys ()] , *Int J Radiat Oncol Biol Phys* 1993. 26 p. .
- 166 [Nag and Gupta ()] 'A simple method of obtaining equivalent doses for use in HDR brachytherapy'. S Nag , N
167 Gupta . *Int J Rdiat Oncol Biol Phys* 2000. 46 p. .
- 168 [Hammer et al. ()] 'Breast cancer: primary treatment with external beam radi-ation therapy and high-dose-rate
169 iridium impla-ntat-ion'. J Hammer , D H Seewald , C E Trach . *Radiology* 1994. 193 p. .
- 170 [Breast Conserving Surgery and Whole Breast Radiation therapy Followed by High Dose Rate Brachytherapy Boost Versus Elect
171 *Breast Conserving Surgery and Whole Breast Radiation therapy Followed by High Dose Rate Brachytherapy*
172 *Boost Versus Electron Beam Boost in the Treatment of Early Breast Cancer in Young Indian Women: Which*
173 *is Cosmetically Better?*,
- 174 [Hannoun-Levi and Marsiglia ()] *Complement d'irradiation par curietherapie dans les cancers du sein: que savons*
175 *nous? Ou allons nous? Cancer/ Radiotherapie*, J M Hannoun-Levi , H Marsiglia . 2004. 8 p. .
- 176 [Borger et al. ()] 'Dose and volume effects on fi brosis after breast conservation therapy'. J H Borger , K H
177 Kemperman , H S Smith . *Int J Radiat Oncol Biol Phys* 1994. 30 p. .
- 178 [Kurtz ()] 'Impact of radiotherapy on breast cosmesis'. J M Kurtz . *Breast* 1995. 4 p. .
- 179 [Deore et al.] *Influence of dose rate and dose per fraction on clinicaloutcome of breast cancer treated by external*
180 *beam irradiation plus irridium-192 implants: Analysis of 289 cases*, S M Deore , R Sarin , K A Dinshaw .
- 181 [Serkies et al. ()] 'Metody wyznaczania i podawania dodatkowej dawki (boost) w oszcz-z?dzaj?cym leczeniu raka
182 piersi'. K Serkies , E W?grzynowicz , J Jassem . *Nowotwory* 2004. 54 p. .
- 183 [Kulik ()] 'Ocena efektu kosmetycznego u chorych na niezaawansowanego raka piersi leczonych w sposob
184 oszcz?dzaj?cy z wykorzystaniem brachy-terapii o wysokiej mocy dawki'. A Kulik . *Praca Doktorska, Ce-ntrum*
185 *Onkologii-Institut im. M. Sk?odowskiej -Curie w Warszawie* 2004. p. 40.
- 186 [Orton et al. (ed.) ()] Cg: Orton , Radiobiology . *Principles and Practice of Brachytherapy*, S Nag (ed.) (Armonk,
187 NY) 1997. Futura Publishing Co. p. .
- 188 [Polgar et al. ()] C Polgar , J Fodor , Z Orosz . *Electron and highdose-rate brachytherapy boost in the conservative*
189 *treatment of stage I-II breast cancer: fi rst results of the randomized Budapest boost trial*, 2002. 178 p. .