

# Clinico Haematological Study of Acute Lymphoblastic leukemia

Dr. Preethi C R<sup>1</sup>

<sup>1</sup> JJM MEDICAL COLLEGE DAVANGERE

*Received: 12 December 2013 Accepted: 31 December 2013 Published: 15 January 2014*

## Abstract

Context: Acute Lymphoblastic leukemia encompasses a group of neoplasms composed of immature, precursor B (pre-B) or T (pre-T) lymphocytes referred to as lymphoblast. Aims: 1. To know the relative incidence of Acute Lymphoblastic leukemia among the patients referred for complete haemogram at the department of pathology, JJMMC, Davangere. 2. To study the clinical manifestations and their correlation with various types of acute Lymphoblastic leukemia. 3. To study the haematological profiles in acute Lymphoblastic leukemia. Settings and design: The study was a hospital based study conducted at haematology unit, Department of Pathology, JJM Medical college, Davangere. Methods and material: The present study was done during the period of June 2006 to May 2008 at haematology unit department of Pathology, JJM Medical college, Davangere. Cases from chigateri general hospital, Bapuji hospital and other private hospitals situated in and around Davangere were included for the study. The case selection was based on clinical features and supported by laboratory evidence. Bone marrow aspiration was subsequently carried out after obtaining written consent from the patient or the guardian. Statistics: The results were expressed in percentage. Results and conclusion: A total of 1039 patients who were referred to the department of haematology out of which 13 patients were diagnosed as Acute Lymphoblastic Leukemia. The present study is to highlight that light microscopic features of peripheral smear and bone marrow will still remain mainstay in the diagnosis of acute Lymphoblastic leukemias. Aims: 1. To know the relative incidence of Acute Lymphoblastic leukemia among the patients referred for complete haemogram at the department of pathology, JJMMC, Davangere. 2. To study the clinical manifestations and their correlation with various types of acute Lymphoblastic leukemia. 3. To study the haematological profiles in acute Lymphoblastic leukemia. Settings and design: The study was a hospital based study conducted at haematology unit,

**Index terms**— leukemias, all, hospital-based study.

## 1 Clinico Haematological Study of Acute Lymphoblastic leukemia

Dr. Preethi C R Abstract-Context: Acute Lymphoblastic leukemia encompasses a group of neoplasms composed of immature, precursor B (pre-B) or T (pre-T) lymphocytes referred to as lymphoblast.

Aims:

1. To know the relative incidence of Acute Lymphoblastic leukemia among the patients referred for complete haemogram at the department of pathology, JJMMC, Davangere. 2. To study the clinical manifestations and their correlation with various types of acute Lymphoblastic leukemia. 3. To study the haematological profiles in acute Lymphoblastic leukemia.

Settings and design: The study was a hospital based study conducted at haematology unit, Department of Pathology, JJM Medical college, Davangere.

Methods and material: The present study was done during the period of June 2006 to May 2008 at haematology unit department of Pathology, JJM Medical college, Davangere.

Cases from chigateri general hospital, Bapuji hospital and other private hospitals situated in and around Davangere were included for the study. The case selection was based on clinical features and supported by laboratory evidence. Bone marrow aspiration was subsequently carried out after obtaining written consent from the patient or the guardian.

## 2 I. Introduction

acute lymphoblastic leukemia encompasses a group of neoplasms composed of immature, precursor(B-pre-B) or T (pre-T) lymphocytes referred to as lymphoblast 1 .

Although ALL affects all age groups, ALL has its highest incidence in children between ages, 1-5 years with a peak at 3-4 years 2 . In 1976, Bennet JM et al classified ALL into three sub types (L1, L2 and L3). According to a) the occurrence of individual cytological features b) The degree of heterogeneity in the distribution among the leukemic population of some or all of these features. The features considered are cell size, nuclear chromatin, nuclear shape, nucleoli, amount and basophilia of the cytoplasm ?? . In the year 1985, the first MIC (morphologic, immunologic, cytogenetic cooperative study group) proposed a classification of ALL 4 . The children cancer study group (CCSG) has presented their own classification of ALL which borrows from FAB nomenclature 5 . The latest WHO classification of the acute leukemias differs from the FAB classification in that greater than or equal to 20% blasts are used for the diagnosis of acute leukemias 6 .

## 3 II. Objectives

1. To know the relative incidence of Acute lymphoblastic leukemia among the patients referred for complete haemogram at the department of pathology, JJMMC, Davangere. 2. To study the clinical manifestations and their correlation with various types of acute lymphoblastic leukemia. 3. To study the haematological profiles in acute lymphoblastic leukemia.

## 4 III. Materials and Methods

The present study on "Clinico-Haematological study of Acute LymphoblasticLeukemias" was undertaken during the period of June 2006 to may 2008 at haematology unit, Department of Pathology, J J M Medical College, Davangere.

The cases from chigateri hospital, Bapuji hospital and other private hospitals situated in and around Davangere formed the material of the study. Case selection was based on clinical features and supported by laboratory evidences. Bone marrow aspiration was subsequently carried out after obtaining written consent from the patient or the guardian.

## 5 Inclusion criteria -New cases of ALL Exclusion criteria - Treated cases of ALL

The following investigations were done: 1. Complete haemogram was performed and peripheral smear was stained by Leishman stain for all cases and examined in detail.

2. Bone marrow aspiration and study was done in all cases and leishman stained smears were examined. In all cases, the following cytochemical stains were employed for diagnosis and subtyping of leukemias.

? MPO -Myelo-peroxidase stain ? SBB -Sudan Black B ? PAS-Periodic Acid Schiff stain ? NSE-Non specific esterase stain Acute lymphoblastic leukemias were classified based on FAB Criteria.

## 6 IV. Results

13 patients in this study were diagnosed as ALL. ALL patients were seen in the age range of 5months to 16 years with a mean age of 8.3 years. The mean ages for males and females were 9.2 years and 6.0 years respectively. Out of the 13 patients, 9 were males and 4 were females, with a male to female ratio of 2.2:1.

The main presenting symptoms were fever in 6 patients (46.1%), generalized weakness in 4 patients (30.8%) and backache in 3 patients (23%).

Physical examination showed pallor of varying degrees in all patients. Lymphadenopathy was present in 8 out of 13 patients constituting (61.5%). All the patients had localized lymphadenopathy among which cervical lymphadenopathy was common.

Mild to moderate Hepatomegaly was seen in 7 patients (53.8%).

Mild to moderate Splenomegaly was seen in 10 patients (76.9%).

Anemia of variable degree was seen in all patients of ALL. The Hb level ranged from 5.9-7.2gm/dl. The mean Hb level being 6.7gm/dl. TLC ranged from  $16.9 \times 10^9 /l$  to  $210 \times 10^9 /l$ . the mean TLC being  $75.8 \times 10^9 /l$ . 5 patients (38.4%) had count between  $11-49 \times 10^9 /l$ , 4 patients (30.7%) had counts between  $50-100 \times 10^9 /l$ , 4 (30.7%) patients had counts  $>100 \times 10^9 /l$ .

All the 13 patients had thrombocytopenia at the time of diagnosis, 5 patients had counts from  $11-49 \times 10^9 /l$ , 4 patients had counts from  $50-100 \times 10^9 /l$ , 4 patients had counts  $>100 \times 10^9 /l$ , with a mean platelet count of  $82.3 \times 10^9 /l$ . 10 out of 13 patients had an ESR of  $>50mm$  at the end of 1 st hour and 3 patients between 20-50mm at the end of 1 st hour.

Bone marrow aspiration was performed in all 13 patients. Lymphoblast was the predominant cell with an average of 85% blasts on differential count. All the patients had decreased megakaryocytes. All were subtyped on morphological basis using FAB criteria into 3 subtypes-L1, L2, and L3. The identification of lymphoblasts was mainly on morphological grounds as stated in the FAB proposals. MPO/SBB and PAS stains were performed. Most of the ALL patients were MPO/SBB negative, but characteristic block positivity on PAS was seen in 38.5% of the patients. All the 13 patients who were diagnosed as ALL were ALL-L2 in this study. The age range was 5 months to 16 years with a mean age of 8.3 years. The mean ages for males and females were 9.2 years and 6 years respectively. Out of the 13 patients, 9 were males and 4 were females, with a male to female ratio of 2.2:1.

The main presenting symptoms were fever in 6 patients (46.1%), generalized weakness in 4 patients (30.8%) and backache in 3patient (23%).

Physical examination showed pallor of varying degrees in all patients. Lymphadenopathy was present in 8 of the 13 patients constituting (61.5%). All the 8 patients had localized lymphadenopathy among which cervical lymphadenopathy was common.

Mild to moderate hepatomegaly was seen in 7 patients (53.8%).

Mild to moderate Splenomegaly was seen in 10 patients (76.9%).

Anemia of variable degree was seen an all patients of ALL. The Hb level ranged from 5.9-7.2gm/dl. The mean Hb level being 6.7gm/dl. TLC ranged from  $16.9 \times 10^9 /l$  to  $210 \times 10^9 /l$ . the mean TLC being  $75.8 \times 10^9 /l$ . 5 patients (38.4%) had count between  $11-49 \times 10^9 /l$ , 4 patients (30.7%) had counts between  $50-100 \times 10^9 /l$ , 4 (30.7%) patients had counts  $>100 \times 10^9 /l$ .

All the 13 patients had thrombocytopenia at the time of diagnosis, 5 patients had counts from  $11-49 \times 10^9 /l$ , 4 patients had counts from  $50-100 \times 10^9 /l$ , 4 patients had counts  $>100 \times 10^9 /l$ , with a mean platelet count of  $82.3 \times 10^9 /l$ . 10 out of 13 patients had an ESR of  $>50mm$  at the end of 1st hour and 3 patients between 20-50mm.

Bone marrow aspiration was done in all 13 patients. Marrow was hypercellular. Aspiration showed reduced erythropoiesis and megakaryopoiesis. Leucopoiesis showed a predominance of lymphoblasts which comprised of heterogenous population of both large and small lymphoblasts. The average blast count was 85%.

Cytochemical staining for MPO/SBB was negative and PAS positivity was seen in 38.5% of the patients.

## 7 c) Acute Lymphoblastic Leukemia (L3)

No case of L3 was encountered in this study.

## 8 V. Discussion

The mean age incidence in the present study was 8.3%. In studies conducted by Shome et al (1985)<sup>7</sup> and Mathur (1993)<sup>8</sup> et al, the mean age incidence was 15.6 and 29.7 respectively. This was more when compared to the present study.

The male female ratio in our study was 2.2:1. In studies done by Shome et al (1985) and Mathur (1993) et al, the male to female ratio was 3.4:1 and 2.4:1 respectively. This is more when compared to the present study.

The average figures for age incidences in the present study are less than the figures quoted in other Indian studies.

The main presenting symptoms were fever and generalized weakness in our study. The same was noted by Shome et al (1985) and Mathur (1993) et al. Bleeding manifestation as a presenting symptom was not noted in this study. Higher incidences of bleeding manifestations were noted by Shome et al (55%) and Mathur et al (47%). A high incidence of lymphadenopathy was seen consistently in this study and also in other studies. Hepatosplenomegaly was also a presenting symptom in this study. Similar observation was noted by Shome et al (1985) and Mathur (1993) et al, but with a more frequency among patients. Backache was seen in 23% of the patients in this study. Backache was not seen in study done by Shome et al (1985) and Mathur (1993) et al.

Pallor was present in our study and it correlates well with a study done by Mathur (1993) Anemia was seen in all the cases of ALL in the present study. Similar finding was observed in study conducted by Mathur Thrombocytopenia was present in all the patients in the present study. The mean platelet count was  $82.3 \times 10^9 /l$  in the present study. This was high when compared to the study conducted by Mathur et al, where it was  $55.2 \times 10^9 /l$ . Bone marrow examination was performed in all 13 cases.

The mean blast percentage was 85%. This was more when compared to Mathur et al where it was 57%. In the present study, all the patients, that were diagnosed as ALL were ALL-L2 (100%). In the study done by Mahendrakumar (1998)<sup>9</sup> L2 was a predominant subtype. Shome et al (1985) reported an almost equal incidence of L1 and L2 (45.8% and 42.4%).

The mean age for L2 was 8.3 years in the present study. Shome et al (1985) reported a mean age of 17.7, whereas Mahendrakumar (1998) reported a mean age 13.6 for L2. In this study, the sex ratio showed a male

predominance in L2 subtype (2.2:1). The Shome et al (1985) showed a sex ratio of 2.6:1 in L2 type, whereas the study of Mahendrakumar (1998) showed a sex ratio of 1.8:1 in L2 type.

The general pattern of clinical features varies with the findings of Shome et al (1985). Fever and generalized weakness were a common initial clinical presentation in the present study whereas in study done by Shome et al (1985) a higher percentage of fever and generalized weakness were noted. Lymphadenopathy was common in ALL-L2 in the present study (61.5%) and is less when compared to Shome et al (73%). Splenomegaly correlates well with study by Shome et al (1985). Hepatomegaly was seen in less frequency when compared to Shome et al (1985). Backache was seen in 23% of the patients in this study and was not seen in study conducted by Shome et al (1985).

The mean Hb levels in ALL-L2 subtype correlates with Shome et al study (1985). TLC also correlates with Shome et al series.

Thrombocytopenia was seen in all patients in this study. The mean platelet count was little less when compared to study done by Shome et al (1985).

## 9 VI. Conclusion

ALL was diagnosed in 13 patients (20.63%). All the patients that were diagnosed with ALL were ALL-L2. Lymphadenopathy was the most consistent feature with fever, generalised weakness, backache and hepatomegaly. Anemia and thrombocytopenia were pre -sent in all the patients. TLC count ranged from  $16.9$  to  $210 \times 10^9 / l$  with a mean of  $75.8 \times 10^9 / l$ . The mean blast percentage was 85%.

The present study is to highlight that light microscopic features of peripheral smear and bone marrow still remain mainstay in the diagnosis of acute leukemias, whereas immunotyping and cytogenetics are complimentary procedures at specialized centres.

However, with newer modalities of therapy and rewarding curative results in haematological malignancies, the use of cytochemistry, immunotyping and cytogenetics have become gold standards for arriving at a specific diagnosis.



Figure 1:

1

| Haematological parameters                    | ALL (n=13) | L1                            | L2       | L3 |
|----------------------------------------------|------------|-------------------------------|----------|----|
| Hb (gm/dl)                                   |            |                               |          |    |
| Range                                        | 5.9-7.2    | -                             | 5.9-7.2  | -  |
| Mean                                         | 6.7        | -                             | 6.7      | -  |
|                                              |            | TLC<br>( $\times 10^9$ /L)    |          |    |
| Range                                        | 16.8-210   | -                             | 16.8-210 | -  |
| Mean                                         | 75.8       | -                             | 75.8     | -  |
|                                              |            | Platelets ( $\times 10^9$ /L) |          |    |
| Range                                        | 0.46-145   | -                             | 0.46-145 | -  |
| Mean                                         | 82.3       | -                             | 82.3     | -  |
| Blast range (%)                              | 60-90      | -                             | 60-90    | -  |
| Bone marrow decreased megakaryocytes         | 13         | -                             | 13       | -  |
| Blast mean                                   | 85%        | -                             | 85%      | -  |
| a) Acute Lymphoblastic Leukemia (L1)         |            |                               |          |    |
| No case of L1 was encountered in this study. |            |                               |          |    |

[Note: b) Acute Lymphoblastic Leukemia (L2)]

Figure 2: Table 1 :

2

et al.

Figure 3: Table 2 :

3

| Haematological parameters     | Mathur et al (1993) | Present study (2008) |
|-------------------------------|---------------------|----------------------|
| Hb gm/dl                      | 2-9.5               | 5.9-7.2              |
| Range                         | 5.2                 | 6.7                  |
| Mean                          |                     |                      |
| TLC ( $\times 10^9$ /l)       | 8-90                | 16.8-210             |
| Range                         | 35.8                | 75.8                 |
| Mean                          |                     |                      |
| Platelets ( $\times 10^9$ /l) | 20-150              | 0.46-145             |
| Range                         | 55.2                | 82.3                 |
| Mean                          |                     |                      |
| Blasts (%)                    | 20-90               | 60-90                |
| Range                         | 57                  | 85                   |
| Mean                          |                     |                      |

Figure 4: Table 3 :

4

| Haematological parameters                   | L1<br>PGI | PS | L2<br>PGI PS |      | L3<br>PGI | PS |
|---------------------------------------------|-----------|----|--------------|------|-----------|----|
| Hb <sub>gm</sub> /dl)                       | 6.6       | -  | 6.8          | 6.7  | 8.4       | -  |
| Mean                                        |           |    |              |      |           |    |
| TLC ( $\times 10^9$ /l)                     | 39.10     | -  | 77.6         | 75.8 | 29        | -  |
| Mean                                        |           |    |              |      |           |    |
| Platelets ( $\times 10^9$ /l)               | 81        | -  | 85           | 69.3 | 49.3      | -  |
| Mean                                        |           |    |              |      |           |    |
| Bone marrow decreased megakaryocytes (in %) | 96        | -  | 90           | 100  | 100       | -  |

Figure 5: Table 4 :

---

181 [Schumacher ()] , Harold Schumacher . *Acute leukemia* 2002. 22 p. .

182 [Fayazmukarab ()] ‘Clinico-haematological study of childhood acute leukemia (unpublished doctoral dissertation’.

183 Khan Fayazmukarab . *Rajiv Gandhi University of Health Sciences* 2003. p. 146.

184 [Abbas and Fausto] *Diseases of white blood cells, lymph nodes, spleen and thymus: Robbins and cotran’s*

185 *Pathologic basis of disease*, Abdul Abbas , Nelson Fausto . Elsevier. 14 p. . (seventh edition)

186 [Kumar ()] Mahendra Kumar , H . *Clinico-haematological study of leukemias*, (Shimoga) 1998. p. 150. Kuvempu

187 University (unpublished doctoral dissertation)

188 [Mathur ()] S K Mathur . *Clinical profile of acute leukemias: A study of 50 cases: Indian Practitioner*, 1993. 46

189 p. .

190 [Behm and Frederick ()] *Morphologic and cytochemical characteristics of childhood lymphoblastic leukemia:*

191 *HaematolOncolClin North Am*, G Behm , Frederick . 1990. 4 p. .

192 [Bennet ()] ‘Morphologic, immunologic and cytogenetic (MIC) working classification of the acute myeloid

193 leukemias second MIC co-operative study group’. J M Bennet . *Br J Haematol* 1988. 68 p. .

194 [Bennet ()] ‘Proposals for classification of the acute leukemias’. J M Bennet . *Br J Haematol* 1976. 33 p. .

195 [Shome ()] ‘The leukemias at presentation: Clinical, Demographic and cytologic variables’. D K Shome . *Ind J*

196 *Cancer* 1985. p. .

197 [Whitlock et al. ()] A Whitlock , Gaynon S James , Paul . *Acute lymphoblastic leukemia in children: Chapter 80*

198 *in Wintrobe’s clinical haematology: 11 th ed: Lippincot Williams and Wilkins company*, 2004. 2 p. .