

Genetic Risk Factors, Biology, Psychosocial Aspects, and Comparison between Male Breast Cancer and Female Breast Cancer

Mojgan Haji Seyed Ebrahim Darkeh¹ and Edward Azavedo²

¹ Karolinska Institute

Received: 7 December 2013 Accepted: 4 January 2014 Published: 15 January 2014

Abstract

To review genetic risk factors, biological and psychosocial aspects of male breast cancer, and comparison between male and female breast cancer. Method: A systematic review of the literature between 1990 and 2013 was conducted to identify studies relevant to the objective. Searches were carried out on the database PubMed, by using the title term "Male Breast Cancer" (MBC). Results: Genetic risk factors for MBC are poorly understood. Family history is a definite risk factor. BRCA2 mutations are more frequent than BRCA1. Men with Klinefelter's syndrome have a high risk of being affected by MBC. The majority of male breast cancers are hormone positive. Informative and psychological support for male breast cancer patients is poor. Diagnosis and treatment of MBC is similar to that of female patients. It is believed that prognosis of male patients is equal to that of age- and stage-matched women, but there is no consensus about this.

Index terms— men at higher risk to reduce symptom duration before diagnosis.

1 Introduction

Male breast cancer is a relatively rare disease, which accounts for less than 1% of all instances of cancer in men and about 1% of all breast cancer cases (1)(2)(3)(4)(5)(6)(7). It accounts for less than 0.2% of all cancer related deaths among men (8)(9)(10)(11). Because of the rarity of the disease, most information about male breast cancer has been obtained from small, monocentric, retrospective studies or through extrapolation from randomized prospective studies or from clinical experience of breast cancer in women (12). But this enormous volume of data on female breast cancer may not be completely relevant to men, particularly with regard to differences concerning the hormonal environment for men and women, and also in terms of gender differences that may affect the cancer patient's condition, medical and/or psychosocial side effects from treatments, and survival priorities.

2 II.

3 Method

This study is a systematic review of the literature. The literature review was conducted in order to describe the current state of knowledge and to compile the scientific literature within the field of breast cancer in men. The study processes the scientific papers in a systematic manner, which consisted of both empirical studies of quantitative and qualitative design, and theoretical or meta-analytic and overview studies. All of them had a clear link to breast cancer in men. The search for scientific literature was conducted in the PubMed database by searching for the key words "male breast cancer" and some articles were also selected from bibliographies from other publications.

Articles that were included in this study meet these criteria ? Articles in English language published between 1990-01-01 and 2013-09-30.

? Articles were about primary breast cancer in men.

? Articles touched heredity and genetic aspects, clinical features, clinical histopathology, diagnosis and diagnostic methods (mammography, ultrasound, fine needle aspiration biopsy / core needle biopsy and sentinel lymph node biopsy), treatment (surgery, radiotherapy, hormone therapy and chemotherapy), prognosis (prognostic factors and survival), and psychosocial aspects.

4 ? Articles made a clear comparison of breast cancer in men and breast cancer in women

Articles were excluded if one or more of the following criteria were matched ? Articles that were case studies or studies with less than 10 patients (with the exception of case studies of unknown / rare genetic factors to MBC or articles with qualitative approach and in-depth interviews). ? Articles that were studies of a certain group of people e.g. breast cancer in transsexual men or in HIV-infected men.

? Articles that were about MBC metastasis.

? Of the total of 812 articles, 187 were included in this review study that deals with genetic aspect, histopathology, and psychosocial aspects of MBC and also comparison between MBC and FBC.

III.

5 Results and Discussion

6 a) Heritability and genetic aspects

The interaction between genetic and environmental factors generally, is probably of major importance for the occurrence of MBC (13). Most known risk factors related to genetic predisposition include positive family history, BRCA gene mutations and Klinefelter's syndrome (14).

Several studies indicate that a family history of breast cancer is associated with greater risk of MBC (15-22). Approximately 15% to 20% of male patients with breast cancer have a positive family history (12)(13). Quite a large percentage of MBC patients have a history of breast cancer in first-degree relatives (13,19,20,(23)(24)(25)(26)(27). A positive family history of either male or female breast cancer among first-degree relatives leads to 2-3 times higher risk of the emergence of MBC (9,21,23,25,28). This risk increases with increasing numbers of affected first-degree relatives and the early onset of breast cancer in the affected relatives (28). There is a strong correlation between heritable mutations in BRCA and the risk of MBC, but BRCA2 mutations are far more frequent than BRCA1 mutations in MBC cases (13,(27)(28)??29)(30)(31)(32). BRCA2 mutations in MBC patients have been assessed in several studies and in various countries (13,(26)(27)(33)(34)(35)(36)(37)(38)(39)(40)(41)(42)(43)(44)(45)(46)(47) and the results vary between 3% and 40% (Figure 1). These studies suggest that the frequency of BRCA2 mutations may reflect the possible genetic differences between different populations but caution should be exercised in interpreting these estimates, because of the small sample populations in the studies (9) or possible selection bias (18). The estimated lifetime risk of breast cancer among male BRCA2 mutation carriers is 5-10%, compared with a 0.1% risk of MBC in the general male population, i.e. 50-100 times higher (48). Moreover, the cumulative risk of MBC in BRCA2 mutation carriers is always higher than in non-carriers, and in all age-groups, but it is highest among those in their thirties (1,500 times higher) and in their forties (630 times higher) and lowest for those in their eighties (69 times higher) (28, ??9). A recent multi-centric study from Italy by Ottini et al. (2012) (30) has shown that BRCA2 is correlated with aggressive tumour behaviour and with higher tumour grade.

Klinefelter's syndrome is also strongly associated with breast cancer in men (9,33). The syndrome is characterized by a rare chromosomal abnormality, 47 XXY, with breast growth, small testes, infertility and increased excretion of follicle-stimulating hormone (FSH) (33), which occurs in less than one man per thousand (9). The mean age of breast cancer patients with Klinefelter's syndrome is 58, which is slightly lower than the average age for other male breast cancer patients (9,33,50). Up to 7% of men with breast cancer may have Klinefelter's syndrome (33,(51)(52)(53). Compared with the frequency of the disease in the general population, breast cancer can be at least 20 times more common in these men (3,9,54). Other less known genetic mutations that have been reported in men with breast cancer include Reifenshtein syndrome or androgen receptor (AR) mutations (55), CYP17 polymorphism (56), Li-Fraumeni syndrome or p53 or CHEK2 mutation ??57-59), Cowden syndrome or PTEN mutation (60) and Lynch syndrome or HNPCC (61), but the correlation between these mutations and increased risk of MBC has not yet been adequately researched (32,62). Almost all histopathological types of breast cancer have been identified in men (32,63). The most common type is invasive ductal carcinoma, which is at least 80% in several studies (10,14,(63)(64)(65)(66)(67)(68)(69)??70)(71) and ductal carcinoma in-situ (DCIS) is far less common, less than 10% (1,63,69,(72)(73)(74). A study by Lanitis et al. (2008) (75) showed that in-situ cancer in men is not as rare as reported in earlier studies, which indicates earlier detection of breast cancer in men (72). Rare tumour types include invasive papillary and medullary lesions (3) and Paget's disease (3,62) and lobular breast tumours (3,69,(76)(77)(78). Male breast tumours are usually sensitive to the hormones oestrogen and progesterone (10,76,79,80,(81)(82)(83)(84)??85)(86)(87), which has been reported at between 55-92% for

oestrogen and 39-89% for progesterone for MBC cases in large, retrospective series (63, 69, 88-89) (Figure 2). Lymph node involvement in the armpit is very frequent, from 41% to 57% in large retrospective series (10, 90-96) and 11-20% of male breast tumours grade I, 55-61% grade II and 22-33% grade III have been reported in large retrospective series (95-94, 81). Generally, 5-15% of MBC patients have metastases at diagnosis (6, 64, 76, 81, 97-98) but the numbers are higher in African and Asian series, i.e. up to 30% (71,99)(100)(101). Details of the growth factor HER2 from various studies are highly variable and up to 56% of male breast tumours display an over-expression of HER2 (12, 43, 57, 102-107).

7 c) Psycho-social aspects

The general perception of breast cancer as a female disease causes surprise in many men when they are diagnosed with breast cancer (77). France et al. (2000) (708) described in their study that men with breast cancer have been shocked to get a breast cancer diagnosis. These patients have not been aware that men can also be affected by such a disease, which is associated with femininity and they have found it difficult to understand that the disease can develop in a male body. Patients have also had difficulty revealing their breast cancer diagnosis to those around them. Iredale et al. (2005) (709) have described in their study that men with breast cancer are afraid of others' reactions, which reduce or question their masculinity. These patients were also uncertain about discussing this sensitive topic with those around them (109). However, when these patients try to talk about it with their friends and colleagues, the reaction becomes distrust and often a subject for fun in a lamentable way (77).

A sense of frustration also occurs due to a lack of relevant information about breast cancer in men (108)(109)(110)(111)(112)(113). Men with breast cancer are more vulnerable in social contacts in comparison with female breast cancer patients (114), which leads to high levels of disease-specific stress (111). The cancer impact and cancer-related stress are worse in young MBC patients compared to those who are older (111, 715). Concerns about masculinity, fear of stigma and experience of isolation are associated with the general lack of knowledge and information about the disease (111). Different experiences of breast cancer care have also been reported by men and women in a study by Sime (2012) (116). MBC and FBC differ mostly with regard to incidence figures, age at diagnosis, frequency of histological tumour types and frequency of expression of hormone receptors (5). The incidence of male breast cancer varies. Some studies report an increase in MBC in a few countries (4,69,77,(117)(118)(119). However, the incidence of female breast cancer (FBC) has been found to be increasing in most countries (50,102,120). The incidence of breast cancer in men has been stable in Europe for several decades (121) and a new international population-based study by Miao et al. (2011) (120) also shows that male breast cancer incidence rates have remained at a stable low level during for the past four decades. Epidemiologically, MBC occurs continuously with a certain average frequency in the general population and is little affected by environmental factors, but conversely, FBC has a tendency to continuously increase, which may be due to the triggering effect of one or more environmental factors (122). Male breast cancer incidence is generally less than 1 per 100,000 population, in contrast to the much higher incidence of female breast cancer of 122 per 100,000 population (102), i.e. the incidence ratio between MBC and FBC is 0.008, but this ratio is higher among African Americans (123) and is fivefold in Africa, 0.042 (124). While the incidence of MBC generally exhibits a uni-modal distribution, with peak incidence at the age of 71 (50,125), the incidence of FBC tends to have a bimodal distribution (126,80) with early-onset and late-onset incidence at 52 and 71 years of age (50). Age-specific incidence for men is steadily increasing either constantly (50) or exponentially (62), but increased age-specific incidence for women is rapid up to 50 years of age, but then at a slower pace after menopause (50). Differences in age-specific incidence between men and women reflect differences in underlying risk factors for the disease (102).

But on the other hand, the international correlation between male and female breast cancer incidence rates is quite strong ($r = 0.69$), meaning that both sexes have several common risk factors for breast cancer (102). Age-specific incidence patterns among men also display a biological similarity between male breast cancer and late-onset female breast cancer (127). This similarity shows that hormonal mechanisms are important (127) but the differences between them may reflect unique mechanisms that may be associated with androgens (18).

Several studies indicate that male breast cancer patients are, on average, 5 to 10 years older than female breast cancer patients at the time of diagnosis (69, 82, 89, 120, 122, 128 -132), but the age gap between men and women is likely to be less in the Middle East and South Asia (5,(133)(134)(135)(136). The differences between men and women in the age presentation may also reflect gender differences in underlying risk factors, pathogens, and/or over-diagnosis (102).

In both sexes, a family history of breast cancer can increase the risk of developing this disease (15,(18)(19)(20)(21)(22). Genetically, MBC is distinct from FBC (43,137). BRCA1 and BRCA2 mutation genes give an increased risk of breast cancer in both sexes (14, 52, 138) but mutations in these genes do not increase the risk of developing male breast cancer at the same rate in women (139). While mutations in BRCA1 in women can give up to 80% lifetime risk of breast cancer, they do not cause as high a level of risk of breast cancer in men (128). In women, between 30% and 86% of hereditary breast cancer has been estimated to be aetiologically linked to BRCA1/2 gene mutations, but estimates of these mutations in MBC are significantly smaller (9). Several studies on men with hereditary breast cancer have shown that BRCA1 mutations are significantly less common (26,27,37,42,140) but many studies show that BRCA2 mutations play a particularly prominent role in

the development of breast cancer in men (13,26,27,39,40,43,46,48,(139)(140)(141)(142)(143) ??144) ??145)(146) BRCA2 mutation also seems to have a stronger role in MBC development than in FBC development in younger people (13,36). It is also suggested that male breast cancer has a higher genetic than female breast cancer (147).

The relationship between MBC and CBC (contra-lateral breast cancer) is much stronger than the relationship between FBC and CBC i.e. 30 times increased risk of CBC in men, compared to 2-4 times increased risk of CBC in women (28).

Histologically, it is impossible to distinguish between MBC and FBC (14). Almost all the histological breast cancer types described in relation to FBC have also been reported in relation to MBC (12). In both sexes, invasive ductal carcinoma (IDC) is a very common form of breast cancer ??8, 65 149-152). But IDC is more common in men than in women (5).

When we look at percentages, non-invasive, insitu breast cancer in men is not higher than is seen in women before the introduction of mammography screening and it may depend on the size of the male breast, which simplifies the detection of small breast lumps in men using clinical breast examination (63). Papillary breast cancer is relatively more common in men (8,89) i.e. 2% to 4% in men compared with 1% in women (49), but lobular carcinoma is less common in men because of the absence of the mammary glands in the normal male breast (8,69,76,78,81,(89)(90) Even rarer subtypes of breast cancer, such as medullary, tubular and mucinous types, have been reported in men, although the male equivalent may be somewhat more unusual than the female (6,63,64,81). Inflammatory breast cancer and Paget's disease have been seen with similar levels of frequency in men and women (10,63). Male breast tumours have a significantly higher frequency of hormone sensitivity, with regard to oestrogen and progesterone, than their female counterparts (85, 103,107,127,135,[153][154][155][156][157][158][159][160] ??161] which implies a different pathogenesis in the development in of this disease (87). Such differences may play key roles in the therapeutic treatment, which should be grounds for different treatment strategies in comparison with female breast cancer (87).

Older studies show that in contrast to women, the frequency of hormone sensitivity in men does not increase with age (63). But a new study by Giordano et al. (2004) (69) has shown that there is a strong link between this hormone sensitivity in male breast tumours and their age, in the same way that has been observed in women. Oestrogen receptor positivity in males may be a result of the low level of circulating oestrogen in the male body, which is similar to that observed in women after menopause (62). In contrast to this interpretation, several researchers suggest that sexually reducing endocrine condition and/or increased oestrogen production may be the underlying hormonal pattern in MBC cases (162). The amount of testosterone that is converted to oestrogen by aromatase is much greater in men than in women, regardless of a woman's menstrual status, which may explain the differences in approach and response with regard to hormonal treatment of MBC and FBC (163).

HER2-positive breast cancer in men is less common than in women (33,(164)(165) which is more correlated with higher cancer stage and with higher histological grade (33). There are several similarities in clinical signs and symptoms of breast cancer in both sexes (73), such as a hard and fixed lump in the breast, with skin or nipple retraction, and nipple discharge and enlarged lymph nodes in the armpit (166). But in MBC, it is more common for the nipple to be affected (76,131,167) because male breast tumours develop just below the nipple, where rudimentary milk ducts are located, and not in the upper lateral quadrant of the breast, which is characteristic in women (33). Selfdetection is the primary form of detection among men, both with cancer in-situ and with invasive breast cancer. However, ductal carcinoma in situ (DCIS) in women is usually detected through screening (168).

Compared to FBC lesions, the edges on MBC lesions are often more defined and calcifications are less frequent and coarser (33,169), Micro-calcifications occur mainly in ductal carcinoma in-situ (DCIS), which is rarer in men (33,73,170) Cancer metastases in the lymphatic tissue of the skin are far more common in male breast cancer compared with female breast cancer (136).

Men undergo mastectomy more often than females (171,172). Men who have undergone mastectomy are more likely than women to receive radiotherapy (95, ??68,173) because of more advanced disease stage and/or more nipple and skin involvement (62). Men are also less likely than women to receive chemotherapy after surgical treatment (173).

FBC prognosis is correlated with patients' age at diagnosis, but conversely, there is no association between age at diagnosis and MBC prognosis. Relative cancer survival in women increases from 35 years of age to age 45-49, and then decreases to the age of 50-59 and then increases again after the age of 65. This means that relative cancer survival in women is a function of age at diagnosis, but relative survival in men has no significant link with age at diagnosis (174).

Compared to women, cancer survival in men is lower, especially in regions where women are routinely examined with mammography (49). This is because of a more advanced stage of disease at presentation, with higher incidence of lymph node involvement (7,67,131) and the low standard of loco-regional treatment for MBC has a significant role for the poorer results (120).

Several studies have shown that survival in MBC patients is almost equal to the survival of FBC patients if the age and stage of disease at diagnosis are matched (50,69,83, ??61,(175)(176)(177), and after adjustment for age at diagnosis, the stage of the disease and the treatment methods, men have actually had significantly better survival from the disease than comparable women (120). A matched analysis of male and female breast cancer patients in a German study by Foerster et al. (2011) (178) has also shown that the 5year disease-free survival

rate was 53.4% (95% CI, range from 54.1 to 66.3%) in men and 62.6%. (95% CI, range from 63.5 to 75.3%) in women, which was not a significant difference ($p > 0.05$), and the 5-year overall survival rate was 71.4% (95% CI, range from 62.1 to 72.7%) in men and 70.3% (95% CI, range from 32.6 to 49.6%) in women, which was also not a significant difference ($p > 0.05$). Xia et al. (2010) (179) have shown in their study that the 5-year and 10-year overall survival rate between Chinese men and women in general are not equal, and that Chinese men have poorer survival rates compared to Chinese women, but when the male group was compared to post-menopausal women, the difference disappeared. In their population-based cohort study, Thalib & Hall (2009) (180) have shown that gender has no significant effect on the prognosis, which was confirmed in a large retrospective study by Hill et al (2005) (181) when 2,923 male breast cancer cases were compared with 442,500 female breast cancer cases. With respect to the variables tumour size, lymph node status, age at diagnosis, histological grading and receptor status, no significant difference has been demonstrated in survival rates for male breast cancer patients compared with female patients in a multivariate analysis in a study by Borgen et al. (1997) (156).

Nahleh et al. (2007) (89) showed in a multivariate analysis that not only tumour size and lymph node status are independent prognostic factors for survival in men, but that gender also serves as an independent prognostic factor. Median survival age between men and women had significant differences when patients have breast cancer at stages I and II, but this difference disappeared at stages III and IV. It was also shown that MBC patients with negative lymph node status had shorter median survival age than FBC patients with the same lymph node status, but this difference also disappeared when both genders had positive lymph node status.

A new study by Ioka et al. (2006) (136) also showed that the 5-year survival rate only decreases with increasing age in men, and that male breast cancer patients have significantly poorer 5-year survival rates compared with women at a corresponding stage of the disease, which is also confirmed in several recent studies (171,(182)(183)(184) Deaths due to primary breast cancer in men is higher than in women, which is also reported in a study of Gnerlich et al. (2011) (185) and this mortality rate has not changed, unlike female breast cancer (117). It should be emphasised here, the importance of adjuvant systemic therapy, mammography screening, and reduced use of hormone replacement therapy for decreasing mortality among women with breast cancer (127).

The gender difference for prognosis may be a result of anatomical differences between the male and female breast, i.e. undeveloped breast tissue in men facilitates the spread of tumours to the lymphatic tissue in the skin and early regional and distant metastases, both on the overlying skin and on the underlying chest muscle (76) and possibly depends on the biological differences between male and female breast tumours (182,184) or on a result of the lack of adjuvant systemic therapy (chemotherapy and/or hormonal therapy) (183) or on the effect of co-morbidity and other primary tumours that act as confounding factors (179). With overall survival as a benchmark for comparison, there is no difference between MBC and FBC prognosis in several studies. However, with disease-specific survival as the benchmark for comparison, the same study shows a significant difference between the two groups (176,179).

It is still unclear whether the MBC prognosis is worse than the FBC prognosis, so there is a need for multicentric prospective studies in this area (177). One should focus on identifying prognostic factors and on defining optimal therapy for MBC patients (173). Psychological differences between male and female patients with breast cancer are also grounds to introduce a different treatment strategy, especially with regard to hormone replacement therapy (117). An early diagnosis with the absence of lymph node involvement has a significant role in improving the outcome of MBC treatment (188).

IV.

8 Conclusion

A significant advance in understanding MBC can improve MBC diagnosis and prognosis. The treatment of MBC has been extrapolated from knowledge available about FBC, although there are many differences in pathogenesis, in biology and in genetics for these two diseases, especially with regard to differences in the role of hormone oestrogen in MBC compared to FBC.

An increased understanding of the potential differences between male and female breast cancer is important, because this can provide new opportunities for therapeutic intervention and probably improved outcomes for MBC treatment. Increased awareness chances of early detection and result in improved prognosis. Clearer treatment guidelines are also necessary in order to improve MBC prognosis. Increased input is needed for informative and psychological support for MBC patients and public education should be oriented toward men at higher risk to reduce symptom duration before diagnosis.



Figure 1: Figure 1 :



Figure 2: Figure 2 :



Figure 3:



Figure 4:

[Falchetti et al.] , M Falchetti , R Lupi , P Rizzolo , K Ceccarelli , I Zanna , V Calò , S Tommasi , G Masala , A Paradiso , A Gulino , G Giannini , A Russo , D Palli , L Ottini .

[Syndrome ()] , Reifenshtein Syndrome . *Nat Genet* 1992. 2 (2) p. .

[Jonasson et al. ()] , J G Jonasson , B A Agnarsson , S Thorlacius . *Male breast cancer in Iceland Int J Cancer* 1996. 65 (4) p. .

[Int J Cancer ()] , *Int J Cancer* 2002. 98 (6) p. .

[Mokbel ()] , Joseph A & Mokbel , K . *Male breast cancer Int J Fertil Womens Med* 2004. 49 (5) p. .

[Eur J Med Genet ()] , *Eur J Med Genet* 2008.

[Macdonald et al. ()] 'A comparative use and patient outcome in males and females with breast cancer'. G Macdonald , C Paltiel , I A Olivotto , S Tyldesley . *Ann Oncol* 2005. 16 p. .

[Willsher et al. ()] 'A comparison of outcome of male breast cancer with female breast cancer'. P C Willsher , I H Leach , I O Ellis , J B Bourke , R W Blamey , J F Robertson . *Am J Cancer* 1997a. 173 p. .

[Satram-Hoang et al. ()] 'A pilot study of male breast cancer in the Veterans Affairs healthcare system'. S Satram-Hoang , E M Moran , Anton - Culver , H Burras , R W Heimann , T M Boggio , I Dykstra-Long , G R Wood , P A Zulka , R Hufnagel , G Bahan , KK . *J Environ Pathol Toxicol Oncol* 2010. 29 (3) p. .

[Young et al. ()] 'A polymorphism in the CYP17 gene is associated with male breast cancer'. I E Young , K M Kurian , C Annink , I H Kunkler , V A Anderson , B B Cohen , M L Hooper , A H Wyllie , C M Steel . *Br J Cancer* 1999. 81 (1) p. .

[Ribeiro et al. ()] *A review of the management of the male breast carcinoma based on an analysis of 420 treated cases*, G G Ribeiro , R Swindell , M Harris , S Baerjee , A Cramer . 1996. p. .

[Thorlacius et al. ()] 'A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes'. S Thorlacius , G Olafsdottir , L Tryggvadottir , S Neuhausen , J G Jonasson , S V Tavtigian , H Tulinius , H M Ogmundsdottir , J E Eyfjörd . *Nat Genet* 1996. 13 p. .

[Giordano ()] 'Adjuvant systemic therapy for male breast carcinoma'. Giordano . *Cancer* 2005a. 104 (11) p. .

[Scott-Conner et al.] *An analysis of male and female breast cancer*, C E Scott-Conner , P R Jochimsen , H R Menck .

[Wang et al. ()] 'Are males with early breast cancer treated differently from females with early breast cancer in Australia and New Zealand?'. J Wang , J Kollias , C Marsh , G Maddern . *Breast* 2009. 18 (6) p. .

[Müller et al. ()] 'Are there biologic differences between male and female breast cancer explaining inferior outcome of men despite equal stage and treatment?'. A C Müller , C Gani , H M Rehm , F Eckert , M Bamberg , T Hehr , M Weinmann . *Strahlenther Onkol* 2012. 188 (9) p. .

[Palli et al. ()] 'Association between the BRCA2 N372H variant and male breast cancer risk: a population-based case-control study in'. D Palli , M Falchetti , G Masala , R Lupi , F Sera , C Saieva , D 'amico , C Ceroti , M Rizzolo , P Caligo , M A Zanna , I Ottini , L . *BMC Cancer* 2007. 7 p. 170.

[Barh ()] 'Biomarkers, critical disease pathways, drug targets, and alternative medicine in male breast cancer'. D Barh . *Curr Drug Targets* 2009. 10 (1) p. .

[Anelli et al. ()] 'Borgen PI Mutations of the p53 gene in male breast cancer'. A Anelli , T F Anelli , B Youngson , P P Rosen . *Cancer* 1995. 75 (9) p. .

[Ottini et al. ()] 'BRCA1 and BRCA2 mutation status and tumor characteristics in male breast cancer: a population-based study in Italy'. L Ottini , G Masala , D 'amico , C . *Cancer Res* 2003. 63 (2) p. .

[Basham et al. ()] 'BRCA1 and BRCA2 mutations in a population-based study of male breast cancer'. V M Basham , J M Lipscombe , J M Ward , S A Gayther , B A Ponder , D F Easton , P D Pharoah . *Breast Cancer Res* 2002. 4 (1) p. R2.

[Evans et al. ()] 'BRCA1/2 mutation analysis in male breast cancer families from North West England'. D G Evans , M Bulman , K Young , E Howard , S Bayliss , A Wallace , F Laloo . *Fam Cancer* 2008. 7 (2) p. .

[Ottini et al. ()] 'BRCA1/BRCA2 mutation status and clinicalpathologic features of 108 male breast cancer cases from Tuscany: a populationbased study in central Italy'. L Ottini , P Rizzolo , I Zanna , M Falchetti , G Masala , K Ceccarelli , V Vezzosi , A Gulino , G Giannini , S Bianchi , F Sera , D Palli . *Breast Cancer Res Treat* 2009. 116 (3) p. .

[BRCA1/BRCA2 rearrangements and CHEK2 common mutations are infrequent in Italian male breast cancer cases Breast Cancer Res Treat 2008. 110 (1) p. .

[Besic et al. ()] 'BRCA2 gene mutations in Slovenian male breast cancer patients'. N Besic , B Cernivc , J De Grève , K Lokar , M Krajc , S Novakovic , J Zgajnar , E Teugels . *Genet Test* 2008. 12 (2) p. .

- [Haraldsson et al. ()] 'BRCA2 germ-line mutations are frequent in male breast cancer patients without a family history of the disease'. K Haraldsson , N Loman , Q X Zhang . *Cancer Res* 1998. 58 (7) p. .
- [Díez et al. ()] 'BRCA2 germ-line mutations in Spanish male breast cancer patients'. O Díez , J Cortés , M Domènech , C Pericay , J Brunet , C Alonso , M Baiget . *Ann Oncol* 2000. 11 (1) p. .
- [Couch et al. ()] 'BRCA2 germline mutations in male breast cancer cases and breast cancer families'. F J Couch , L M Farid , M L Deshano , S V Tavtigian , K Calzone . *Nat Genet* 1996. 13 p. .
- [Kwiatkowska et al. ()] 'BRCA2 germline mutations in male breast cancer patients in the Polish population'. E Kwiatkowska , M Teresiak , K M Lamperska . *Hum Mutat* 2001. 17 (1) p. 73.
- [Anan et al. ()] 'Breast cancer in Japanese men: does sex affect prognosis?'. K Anan , S Mitsuyama , K Nishihara , Y Abe , T Iwashita , T Ihara , K Tamae , M Ono , S Toyoshima . *Breast Cancer* 2004. 11 p. .
- [Rai et al. ()] 'Breast cancer in males: a PGIMER experience'. B Rai , S Ghoshal , S C Sharma . *J Cancer Res Ther* 2005. 1 (1) p. .
- [Donegan and Redlich ()] 'Breast cancer in men'. & Donegan , Redlich . *Surg Clin North Am* 1996. 76 (2) p. .
- [Giordano et al. ()] 'Breast cancer in men'. S H Giordano , A U Buzdar , G N Hortobagyi . *Ann Intern Med* 2002. 137 (8) p. .
- [Harlan et al. ()] 'Breast cancer in men in the United States: a population-based study of diagnosis, treatment, and survival'. L C Harlan , J A Zujewski , M T Goodman , J L Stevens . *Cancer* 2010. 116 (15) p. .
- [Perkins (2003)] *Breast cancer in men Treatment is based on results extrapolated from trials for women with breast cancer BMJ*, Perkins . August 2003. 327 p. .
- [Robinson et al. ()] 'Breast cancer in men: A need for psychological intervention'. J D Robinson , K P Metoyer , N Bhayani . *Journal of Clinical Psychology in Medical Settings* 2008. 15 (2) p. .
- [Rosenblatt et al. ()] 'Breast cancer in men: aspects of familial aggregation'. K A Rosenblatt , D B Thomas , A Mctiernan , M A Austin , H Stalsberg , A Stemhagen . *J Natl Cancer Inst* 1991. 83 p. .
- [Thomas et al. ()] 'Breast cancer in men: risk factors with hormonal implications'. D B Thomas , L M Jimenez , A Mctiernan , K Rosenblatt , H Stalsberg , A Stemhagen , W D Thompson , M G Curnen , W Satariano , D F Austin . *Am J Epidemiol* 1992. 135 p. .
- [Anderson and Badzioch ()] 'Breast cancer risks in relatives of male breast cancer patients'. D E Anderson , M D Badzioch . *J Natl Cancer Inst* 1992. 84 p. .
- [Levi et al. ()] 'Breast cancer survival in relation to sex and age'. F Levi , L Randimbison , La Vecchia , C . *Oncology* 1992. 49 (6) p. .
- [Giordano ()] 'Breast carcinoma in men: a population-based study'. Giordano . *Cancer* 2004. 101 (1) p. .
- [Gill et al. ()] 'Breast diseases in males—a morphological review of 150 cases'. M S Gill , N Kayani , M N Khan , S H Hasan . *J Pak Med Assoc* 2000. 50 (6) p. .
- [Morimoto et al. ()] 'Cancer of the male breast'. T Morimoto , K Komaki , T Yamakawa , T Tanaka , Y Oomine , Y Konishi , T Mori , Y Monden . *J Surg Oncol* 1990. 44 (3) p. .
- [Sandler et al. ()] 'Cancer of the male breast'. B Sandler , C Carman , R R Perry . *Am Surg* 1994. 60 (11) p. .
- [Donegan et al. ()] 'Carcinoma of the breast in males: a multiinstitutional survey'. W L Donegan , P N Redlich , P J Lang , M T Gall . *Cancer* 1998. 83 (3) p. .
- [Amir and Hirji ()] 'Carcinoma of the male breast in Tanzania'. H Amir , K F Hirji . *J Natl Med Assoc* 1992. 84 (4) p. .
- [Vinod and Pendlebury ()] 'Carcinoma of the male breast: a review of adjuvant therapy'. & Vinod , Pendlebury . *Australas Radiol* 1999. 43 (1) p. .
- [Jamal et al. ()] 'Carcinoma of the male breast: a study of 141 cases from northern Pakistan'. S Jamal , N Mamoon , S Mushtaq . *Asian Pac J Cancer Prev* 2006. 7 (1) p. .
- [Borgen et al. ()] 'Carcinoma of the male breast: analysis of prognosis compared with matched female patients'. P I Borgen , R T Senie , W M Mckinnon . *Ann Surg Oncol* 1997. 4 (5) p. .
- [Schaub et al. ()] *Changes in Male Breast Cancer Over a 30-Year Period The American Surgeon*, N P Schaub , N Maloney , H Schneider , E Feliberti , R Perry . 2008. p. .
- [Xia et al. ()] 'Chinese female breast cancer patients show a better overall survival than their male counterparts'. L P Xia , F F Zhou , G F Guo , F Wang , X Wang , Z Y Yuan , B Zhang . *Chin Med J (Engl)* 2010. 123 (17) p. .
- [Ottini et al. ()] 'Clinical and pathologic characteristics of BRCA-positive and BRCA-negative male breast cancer patients: results from a collaborative multicenter study in Italy'. L Ottini , V Silvestri , P Rizzolo , M Falchetti , I Zanna , C Saieva , G Masala , S Bianchi , S Manoukian , M Barile , P Peterlongo , L Varesco , S Tommasi , A Russo , G Giannini , L Cortesi , A Viel , M Montagna , P Radice , D Palli . *Breast Cancer Res Treat* 2012. 134 (1) p. .

387 [Frank et al. ()] 'Clinical characteristics of individuals with germline mutations in BRCA1 and BRCA2: analysis
388 of 10,000 individuals'. T S Frank , A M Deffenbaugh , J E Reid , M Hulick , B E Ward , B Lingenfelter , K
389 L Gumpfer , T Scholl , S V Tavtigian , D R Pruss , G C Critchfield . *Clin Oncol* 2002. 20 (6) p. .

390 [Meijer-Van Gelder et al. ()] *Clinical relevance of biologic factors in male breast cancer*, Meijer-Van Gelder , M
391 P Look , J Bolt-De Vries , H A Peters , J G Klijn , J A Foekens . 2001. p. .

392 [Tawil et al. ()] *Clinicopathologic and immunohistochemical characteristics of male breast cancer: a single center
393 experience*, A N Tawil , F I Boulos , Z I Chakhachiro , Z K Otrrock , L Kandaharian , El Saghir , NS , Abi
394 Saad , Gs . 2012. 18 p. .

395 [Liu et al. ()] 'Clinicopathologic characteristics and survival of male breast cancer'. D Liu , G Xie , M Chen . *Int
396 J Clin Oncol* 2014. 19 (2) p. .

397 [Park et al. ()] 'Clinicopathological characteristics of male breast cancer'. S Park , J H Kim , Koo J Park , B W
398 Lee , KS . *Yonsei Med J* 2008. 49 p. .

399 [Shah et al. ()] 'Clinicopathological study of male breast carcinoma: 24 years of experience'. P Shah , I Robbani
400 , O Shah . *Ann Saudi Med* 2009. 29 (4) p. .

401 [Hill et al. ()] 'Comparison of male and female breast cancer incidence trends, tumor characteristics, and survival'.
402 T D Hill , H J Khamis , J E Tyczynski , H J Berkel . *Ann Epidemiol* 2005. 15 (10) p. .

403 [Sperlongano and Pisaniello ()] 'Current management of male breast cancer'. P Sperlongano , D Pisaniello . *Ann
404 Ital Chir* 2000. 71 (2) p. .

405 [Borgen et al. ()] 'Current management of male breast cancer. A review of 104 cases'. P I Borgen , G Y Wong ,
406 V Vlamis , C Potter , B Hoffmann , D W Kinne , M P Osborne , W M Mckinnon . *Ann Surg* 1992. 215 (5)
407 p. 451.

408 [Vecchia et al. ()] 'Descriptive epidemiology of male breast cancer in Europe'. La Vecchia , C Levi , F Lucchini
409 , F . *Int J Cancer* 1992. 51 p. .

410 [Lanitis ()] 'Diagnosis and management of male breast cancer'. Lanitis . *World J Surg* 2008. 32 (11) p. .

411 [Rudan et al. ()] 'Differences between male and female breast cancer. I. Epidemiological features'. I Rudan , N
412 Rudan , M Strnad . *Acta Med Croatica* 1995. 49 (3) p. .

413 [Hittmair et al. ()] 'Ductal carcinoma in situ (DCIS) in the male breast: a morphologic study of 84 cases of pure
414 DCIS and 30 cases of DCIS associated with invasive carcinoma—a preliminary report'. A P Hittmair , R A
415 Lininger , F A Tavassoli . *Cancer* 1998. 83 (10) p. .

416 [Weiss et al. ()] 'Epidemiology of male breast cancer'. J R Weiss , K B Moysich , H Swede . *Cancer Epidemiol
417 Biomarkers Prev* 2005. 14 (1) p. .

418 [Temnim et al. ()] 'Evaluation of prognostic factors in male breast cancer'. L Temnim , Y A Luqmani , M
419 Jarallah , I Juma , M Mathew . *Breast* 2001. 10 (2) p. .

420 [Pituskin et al. ()] 'Experiences of men with breast cancer: a qualitative study'. E Pituskin , B Williams , H-J
421 Au , K Martin-Mcdonald . *The Journal of Men's Health & Gender* 2007. 4 (1) p. .

422 [Williams et al. ()] 'Experiences of men with breast cancer: an exploratory focus group study'. B G Williams ,
423 R Iredale , K Brain , E France , P Barrett-Lee , J Gray . *Br J Cancer* 2003. 89 (10) p. .

424 [Chung et al. ()] 'Expression of prognostic factors (EGFR, ER) by immunohistochemical staining method in
425 male breast cancer'. H C Chung , D L Kim , E H Koh . *Yonsei Med J* 1991. 32 (2) p. .

426 [Hemminki and Vaittinen ()] 'Familial cancers in a nationwide family cancer database: age distribution and
427 prevalence'. K Hemminki , P Vaittinen . *Eur J Cancer* 1999. 35 (7) p. .

428 [Struewing and Coriaty ()] 'Founder BRCA1/2 mutations among male patients with breast cancer in Israel'. J
429 P Struewing , Z M Coriaty , RonE . *Am J Hum Genet* 1999. 65 (6) p. .

430 [Callari et al. ()] 'Gene expression analysis reveals a different transcriptomic landscape in female and male breast
431 cancer'. M Callari , V Cappelletti , De Cecco , L Musella , V Miodini , P Veneroni , S Gariboldi , M Pierotti
432 , M A Daidone , MG . *Breast Cancer Res Treat* 2011. 127 (3) p. .

433 [De Jong et al. ()] 'Genes other than BRCA1 and BRCA2 involved in breast cancer susceptibility'. M M De Jong
434 , I M Nolte , G J Te Meerman , W T Van Der Graaf , J C Oosterwijk , J H Kleibeuker . *J Med Genet* 2002.
435 39 (4) p. .

436 [Sverdlov et al. ()] 'Genetic analyses of male breast cancer in Israel'. R S Sverdlov , I Barshack , Bar Sade , R B
437 Baruch , R G Hirsh-Yehezkel , G Dagan , E Feinmesser , M Figer , A Friedman , E . *Genet Test* 2000. 4 (3)
438 p. .

439 [Deb et al. ()] 'Genotypic and phenotypic analysis of familial male breast cancer shows under representation of
440 the HER2 and basal subtypes in BRCA-associated carcinomas'. S Deb , N; Jene , Kconf Ab Investigators , S
441 B Fox . *BMC Cancer* 2012. 12 p. 510.

- [Mavraki et al. ()] 'Germline BRCA2 mutations in men with breast cancer'. E Mavraki , I C Gray , D T Bishop , N K Spurr . *Br J Cancer* 1997. 76 p. .
- [Giordano A review of the diagnosis and management of male breast cancer Oncologist ()] 'Giordano A review of the diagnosis and management of male breast cancer'. *Oncologist* 2005b. 10 (7) p. .
- [Giordano A review of the diagnosis and management of male breast cancer Oncologist ()] 'Giordano A review of the diagnosis and management of male breast cancer'. *Oncologist* 2005b. 10 (7) p. .
- [Rudlowski et al. ()] 'Her-2/neu gene amplification and protein expression in primary male breast cancer'. C Rudlowski , N Friedrichs , A Faridi , L Füzesi , R Moll , G Bastert , W Rath , R Büttner . *Breast Cancer Res Treat* 2004. 84 (3) p. .
- [Evans et al. ()] 'High detection rate for BRCA2 mutations in male breast cancer families from North West England'. D G Evans , M Bulman , K Young . *Fam Cancer* 2001. 1 (3-4) p. .
- [Csokay et al. ()] 'High frequency of germ-line BRCA2 mutations among Hungarian male breast cancer patients without family history'. B Csokay , N Udvarhelyi , Z Sulyok , I Besznyak , S Ramus , B Ponder , E Olah . *Cancer Res* 1999. 59 p. .
- [Stalsberg et al. ()] 'Histologic types and hormone receptors in breast cancer in men: a population-based study in 282 United States men'. H Stalsberg , D B Thomas , K A Rosenblatt , L M Jimenez , A Mctiernan , A Stemhagen , W D Thompson , M G Curnen , W Satariano , D F Austin . *Cancer Causes Control* 1993. 4 (2) p. .
- [Sharif and Mamoon ()] 'Histological and immuno-histochemical study of male breast carcinoma in Northern Pakistan'. M A Sharif , N Mamoon , Arif . *J Pak Med Assoc* 2009. 59 (2) p. .
- [Czene et al. ()] 'How to treat male breast cancer'. K Czene , J Bergqvist , P Hall , J Bergh . *Breast* 2007. 16 (2) p. . (Suppl)
- [Ndom et al. ()] 'Huo D A meta-analysis of male breast cancer in Africa'. P Ndom , G Um , E M Bell , A Eloundou , N M Hossain . *Breast* 2012. 21 (3) p. .
- [Chen et al. ()] 'Imaging characteristics of malignant lesions of the male breast'. L Chen , P K Chandra , L H Larsen , P Barton , M Rohitopakarn , E Q Zhu , L W Bassett . *Radiographics* 2006. 26 (4) p. .
- [Dawson et al. ()] 'Immunocytochemical characterization of male breast cancer'. P J Dawson , T M Paine , S R Wolman . *Mod Pathol* 1992. 5 (6) p. .
- [Anderson and Devesa ()] 'In situ male breast carcinoma in the Surveillance, Epidemiology, and End Results database of the National Cancer Institute'. W F Anderson , S S Devesa . *Cancer* 2005. 104 (8) p. .
- [Miao ()] 'Incidence and outcome of male breast cancer: an international population-based study'. Miao . *J Clin Oncol* 2011. 20 (33) p. .
- [Burga et al. ()] 'Invasive carcinomas of the male breast: a morphologic study of the distribution of histologic subtypes and metastatic patterns in 778 cases'. A M Burga , O Fadare , R A Lininger . *Virchows Arch* 2006. 449 (5) p. .
- [Anderson et al. ()] 'Is male breast cancer similar or different than female breast cancer?'. W F Anderson , M D Althuis , L A Brinton , S S Devesa . *Breast Cancer Res Treat* 2004. 83 p. .
- [Koc Polat Epidemiology and aetiological factors of male breast cancer: a ten years retrospective study in eastern Turkey Eur J Cancer] 'Koc & Polat Epidemiology and aetiological factors of male breast cancer: a ten years retrospective study in eastern Turkey'. *Eur J Cancer Prev* 2001. 10 (6) p. .
- [Hill et al. ()] 'Localized male breast carcinoma and family history. An analysis of 142 patients'. A Hill , Y Yagmoury , K N Tran , Bolton Js , M Robson , P I Borgen . *Cancer* 1999. 86 p. .
- [Wick ()] 'Low-stage carcinoma of the male breast. A histologic, immunohistochemical, and flow cytometric comparison with localized female breast carcinoma'. Wick . *Am J Clin Pathol* 1999. 111 (1) p. .
- [Crichlow and Galt ()] 'Male breast cancer'. & Crichlow , Galt . *Surg Clin North Am* 1990. 70 (5) p. .
- [Kayani et al. ()] 'Male breast cancer'. N Kayani , M N Khan , Y Bhurgri , S Gill , M I Nasir , T Siddiqui . *J Pak Med Assoc* 2003. 53 (3) p. .
- [Fentiman et al. ()] 'Male breast cancer'. I S Fentiman , A Fourquet , G N Hortobagyi . *Lancet* 2006. 367 (9510) p. .
- [Ottini et al. ()] 'Male breast cancer'. L Ottini , D Palli , S Rizzo , M Federico , V Bazan , A Russo . *Crit Rev Oncol Hematol* 2010. 73 (2) p. .
- [Gómez-Raposo et al. ()] 'Male breast cancer'. C Gómez-Raposo , Zambrana Tévar , F , Sereno Moyano , M , López Gómez , M Casado , E . *Cancer Treat Rev* 2010. 36 (6) p. .
- [Reis et al. ()] 'Male breast cancer'. L O Reis , F G Dias , M A Castro , U Ferreira . *Aging Male* 2011. 14 (2) p. .
- [Rudlowski ()] *Male Breast Cancer Breast Care (Basel)*, Rudlowski . 2008. 3 p. .

497 [Fackenthal et al. ()] 'Male breast cancer in Cowden syndrome patients with germline PTEN mutations'. J D
498 Fackenthal , D J Marsh , A L Richardson , S A Cummings , C Eng , B G Robinson , O I Olopade . *J Med*
499 *Genet* 2001. 38 p. .

500 [Chikaraddi et al. ()] 'Male breast cancer in Indian patients: is it the same?'. S B Chikaraddi , R Krishnappa ,
501 V Deshmane . *Indian J Cancer* 2012. 49 (3) p. .

502 [Boyd et al. ()] 'Male breast cancer in the hereditary nonpolyposis colorectal cancer syndrome'. J Boyd , E Rhei
503 , M G Federici , P I Borgen , P Watson , B Franklin . *Breast Cancer Res Treat* 1999. 53 (1) p. .

504 [Nahleh et al. ()] 'Male breast cancer in the Veterans Affairs population: a comparative analysis'. Z A Nahleh ,
505 R Srikantiah , M Safa . *Cancer* 2007. 109 (8) p. .

506 [El-Habbash and Alwindi ()] 'Male breast cancer in Tripoli, Libya'. M M El-Habbash , A A Alwindi . *Saudi Med*
507 *J* 2009. 30 (8) p. .

508 [André et al. ()] 'Male breast cancer—a reappraisal of clinical and biologic indicators of prognosis'. S André , I
509 Fonseca , A E Pinto , P Cardoso , T Pereira , J Soares . *Acta Oncol* 2001. 40 (4) p. .

510 [Krause ()] 'Male breast cancer—an andrological disease: risk factors and diagnosis'. Krause . *Andrologia* 2004.
511 36 (6) p. .

512 [Herman ()] 'Male breast cancer. Does the prognosis differ compared to female?'. Herman . *Neoplasma* 2000. 47
513 p. .

514 [Cutuli et al. ()] 'Male breast cancer. Evolution of treatment and prognostic factors. Analysis of 489 cases'. B
515 Cutuli , C C Le-Nir , D Serin , Y Kirova , Z Gaci , C Lemanski , B De Lafontan , M Zoubir , P Maingon ,
516 H Mignotte , C T De Lara , Edeline J Penault-Llorca , F Romestaing , P Delva , C Comet , B Belkacemi ,
517 Y . *Crit Rev Oncol Hematol* 2010. 73 (3) p. .

518 [Nahleh and Girnius ()] 'Male breast cancer: a gender issue'. Z Nahleh , S Girnius . *Nat Clin Pract Oncol* 2006.
519 3 p. .

520 [Anderson et al. ()] 'Male breast cancer: a population-based comparison with female breast cancer'. W F
521 Anderson , I Jatoi , J Tse , P S Rosenberg . *J Clin Oncol* 2010. 28 (2) p. .

522 [Bourhahfour et al. ()] 'Male breast cancer: a report of 127 cases at a Moroccan institution'. M Bourhahfour , R
523 Belbaraka , A Souadka , M 'rabeti , H Tijami , F Errihani , H . *BMC Res Notes* 2011. 4 p. 219.

524 [Yoney et al. ()] 'Male breast cancer: A retrospective anal-ysis'. A Yoney , A Kucuk , M Unsal . *Cancer Radiother*
525 2009. 13 p. .

526 [Baojiang et al. ()] 'Male breast cancer: A retrospective study comparing survival with female breast cancer'. L
527 Baojiang , L Tingting , L Gang , Z Li . *Oncol Lett* 2012. 4 (4) p. .

528 [Shukla et al. ()] 'Male breast cancer: a retrospective study from a regional cancer center in northern India'. N
529 K Shukla , V Seenu , A K Goel , V Raina , G K Rath , R Singh , A K Kriplani , S V Deo , M C Misra . *J*
530 *Surg Oncol* 1996. 61 (2) p. .

531 [Tan and Sng ()] 'Male breast cancer: a retrospective study with immunohistochemical analysis of hormone
532 receptor expression'. P H Tan , I T Sng . *Pathology* 1997. 29 (1) p. .

533 [Agrawal et al. ()] *Male breast cancer: a review of clinical management Breast Cancer Res Treat*, A Agrawal , A
534 A Ayantunde , R Rampaul , J F Robertson . 2007. 103 p. .

535 [Hayes ()] 'Male breast cancer: a review of the literature'. Ravandi-Kashani & Hayes . *Eur J Cancer* 1998. 34
536 (9) p. .

537 [Gennari et al. ()] 'Male breast cancer: a special therapeutic problem. Anything new? (Review)'. R Gennari , G
538 Curigliano , B A Jereczek-Fossa , S Zurrida , G Renne , M Intra , V Galimberti , A Luini , R Orecchia , G
539 Viale , A Goldhrisch , U Veronesi . *Int J Oncol* 2004. 24 (3) p. .

540 [Liukkonen et al. ()] 'Male breast cancer: a survey at the Helsinki University Central Hospital during 1981-2006'.
541 S Liukkonen , T Saarto , H Maenpaa . *Acta Oncol* 2010. 49 (3) p. .

542 [Mclachlan et al. ()] 'Male breast cancer: an 11 year review of 66 patients'. S A Mclachlan , C Erlichman , F F
543 Liu . *Breast Cancer Res Treat* 1996. 40 (3) p. .

544 [Onami et al. ()] 'Male breast cancer: an update in diagnosis, treatment and molecular profiling'. S1 Onami , M
545 Ozaki , J E Mortimer , S K Pal . *Maturitas* 2010. 65 (4) p. .

546 [Stierer et al. ()] 'Male breast cancer: Austrian experience'. M Stierer , H Rosen , W Weitensfelder , H
547 Hausmaninger , B Teleky , R Jakesz , H Fruhwirth , M Dünser , S Beller , A Haid . *World J Surg* 1995. 19
548 (5) p. .

549 [Hodgson et al. ()] 'Male breast cancer: is the incidence increasing?'. N C Hodgson , J H Button , D Franceschi
550 , F L Moffat , A S Livingstone . *Ann Surg Oncol* 2004. 11 (8) p. .

551 [Contractor et al. ()] 'Male breast cancer: is the scenario changing'. K B Contractor , K Kaur , G S Rodrigues ,
552 D M Kulkarni , H Singhal . *World J Surg Oncol* 2008. 16 p. 58.

- [Cutuli ()] 'Male breast cancer: results of the treatments and prognostic factors in 397 cases'. Cutuli . *Eur J Cancer* 1995. 1995. 31 (12) p. .
- [Akbulut et al. ()] 'Male breast cancer: thirteen years' experience of a single center'. S Akbulut , I Arer , A Kocbiyik , M C Ya?murdur , H Karakayal? , M Haberal . *Int Semin Surg Oncol* 2009. 6 p. 4.
- [Tahmasebi et al. ()] 'Male breast cancer; analysis of 58 cases in Shiraz, South of Iran'. S Tahmasebi , M Akrami , S Omidvari . *Breast Dis* 2010. 31 (1) p. .
- [Meguerditchian et al. ()] 'Male breast carcinoma'. A N Meguerditchian , M Falardeau , G Martin . *Can J Surg* 2002. 45 (4) p. .
- [Goss ()] 'Male breast carcinoma: a review of 229 patients who presented to the Princess Margaret Hospital during 40 years: 1955-1996'. Goss . *Cancer* 1999. 85 (3) p. .
- [Joshi ()] 'Male breast carcinoma: an evaluation of prognostic factors contributing to a poorer outcome'. Joshi . *Cancer* 1996. 77 (3) p. .
- [Davis et al. ()] 'Male breast carcinoma: clinical experience in a suburban community'. S L Davis , F Barse , F G Meoli . *J Am Osteopath Assoc* 1992. 1013. 92 (8) p. .
- [Ciatto et al. ()] 'Male breast carcinoma: review of a multicenter series of 150 cases. Coordinating Center and Writing Committee of FONCAM (National Task Force for Breast Cancer)'. S Ciatto , A Iossa , R Bonardi , P Pacini . *Italy. Tumori* 1990. 76 (6) p. .
- [France et al. ()] 'Male cancer: a qualitative study of male breast cancer'. L France , S Michie , P Barrett-Lee , K Brain , P Harper , J Gray . *Breast* 2000. 9 (6) p. .
- [Muir et al. ()] 'Male versus female breast cancers. A population-based comparative immunohistochemical analysis'. D Muir , R Kanthan , S C Kanthan . *Arch Pathol Lab Med* 2003. 127 (1) p. .
- [Cooper et al. ()] 'Mammography in men'. R A Cooper , B A Gunter , L Ramamurthy . *Radiology* 1994.
- [Ahmed ()] 'Management and outcomes of male breast cancer in zaria, Nigeria'. Ahmed . *Int J Breast Cancer* 2012. 2012 p. 845143.
- [Foerster et al. ()] 'Matched-pair analysis of patients with female and male breast cancer: a comparative analysis'. R Foerster , F G Foerster , V Wulff , B Schubotz , D Baaske , M Wolfgarten , W C Kuhn , C Rudlowski . *BMC Cancer* 2011. 11 p. 335.
- [Marchal et al. ()] 'Men with breast cancer have same diseasespecific and even-free survival as woman'. F Marchal , M Salou , C Marchal , A Lesur , E Desandes . *Ann Surg Oncol* 2009. 16 p. .
- [Sime ()] *Men's experiences of having breast cancer: a comparison with women's experiences*, C A Sime . 2012. University of Glasgow (PhD thesis)
- [Rayson et al. ()] 'Molecular markers in male breast carcinoma'. D Rayson , C Erlichman , V J Suman , P C Roche , L E Wold , J N Ingle , J H Donohue . *Cancer* 1998. 83 (9) p. .
- [André and Soares ()] 'Morphology of male breast carcinoma in the evaluation of prognosis'. S André , J Soares . *Pathol Res Pract* 1990. 186 (6) p. .
- [Korde et al. ()] 'Multidisciplinary meeting on male breast cancer: summary and research recommendations'. L A Korde , J A Zujewski , L Kamin , S Giordano , S Domchek , W F Anderson , J M Bartlett , K Gelmon , Z Nahleh , J Bergh , B Cutuli , G Pruneri , W Mccaskill-Stevens , J Gralow , G Hortobagyi , F Cardoso . *J Clin Oncol* 2010. 28 (12) p. .
- [Friedman et al. ()] 'Mutation analysis of BRCA1 and BRCA2 in a male breast cancer population'. L S Friedman , S A Gayther , T Kurosaki , D Gordon , B Noble , G Casey , B A Ponder . *Am J Hum Genet* 1997. 60 p. .
- [Fox et al. ()] 'Oestrogen receptor and epidermal growth factor receptor expression in male breast carcinoma'. S B Fox , S Rogers , C A Day , J C Underwood . *J Pathol* 1992. 166 (1) p. .
- [Idelevich et al. ()] 'Oncogenes in male breast cancer'. E Idelevich , M Mozes , N Ben-Baruch , M Huszar , A Kruglikova , R Katsnelson , A Shani . *Am J Clin Oncol* 2003. 26 (3) p. .
- [Patterson ()] 'Outcome of men presenting with clinical breast problems: the role of mammography and ultrasound'. Patterson . *Breast J* 2006. 2006. 12 (5) p. .
- [Arslan et al. ()] 'Outcome of non-metastatic male breast cancer: 118 patients'. U Y Arslan , B Oksüzo?lu , N Ozdemir , S Aksoy , N Alk?? , A Gök , M A Kaplan , M Gümü? , V Berk , D Uncu , M Baykara , D Colak , U Uyetürk , I Türker , A I?kdo?an . *Med Oncol* 2012. 29 (2) p. .
- [Chen et al. ()] 'Poorer Survival of Male Breast Cancer Compared with Female Breast Cancer Patients May Be Due to Biological Differences'. X Chen , X Liu , L Zhang , S Li , Y Shi , Z Tong . *Jpn J Clin Oncol* 2013. 43 (10) p. .
- [Gnerlich et al. ()] 'Poorer survival outcomes for male breast cancer compared with female breast cancer may be attributable to in-stage migration'. J L Gnerlich , A D Deshpande , D B Jeffe , S Seelam , E Kimbuende , J A Margenthaler . *Ann Surg Oncol* 2011. 18 (7) p. .

- [Olsson et al. ()] 'Population-based cohort investigations of the risk for malignant tumors in first degree relatives and wives of men with breast cancer'. H Olsson , H Andersson , O Johansson , T R Moller , U Kristoffersson , E Wenngren . *Cancer* 1993. 71 p. .
- [Wolpert et al. ()] 'Prevalence of BRCA1 and BRCA2 mutations in male breast cancer patients in Canada'. N Wolpert , E Warner , M F Seminsky , A Futreal , S A Narod . *Clin Breast Cancer* 2000. 1 (1) p. .
- [Hultborn et al. ()] 'Prevalence of Klinefelter's syndrome in male breast cancer patients'. R Hultborn , C Hanson , I Kopf , I Verbiene , E Warnhammar , A Weimarck . *Anticancer Res* 1997. 17 p. .
- [Mathew et al. ()] 'Primary breast cancer in men: clinical, imaging, and pathologic findings in 57 patients'. J Mathew , G H Perkins , T Stephens , L P Middleton , W T Yang . *AJR Am J Roentgenol* 2008. 191 (6) p. .
- [Salvadori et al. ()] 'Prognosis of breast cancer in males: an analysis of 170 cases'. B Salvadori , R Saccozzi , A Manzari , S Andreola , R A Conti , F Cusumano , M Grassi . *Eur J Cancer* 1994. 30 (7) p. 930.
- [Avisar ()] 'Prognostic factors in node-negative male breast cancer'. Avisar . *Clin Breast Cancer* 2006. 7 (4) p. .
- [Moore et al. ()] 'Prognostic indicators in male breast carcinoma'. J Moore , M I Friedman , T Gansler . *Breast J* 1998. 4 (4) p. .
- [Brinton et al. ()] 'Prospective evaluation of risk factors for male breast cancer'. L A Brinton , D A Richesson , G L Gierach . *J Natl Cancer Inst* 2008. 100 (20) p. .
- [Brain et al. ()] 'Psychological distress in men with breast cancer'. K Brain , B Williams , R Iredale , L France , J Gray . *Journal of Clinical Oncology* 2006. 24 (1) p. .
- [Izquierdo ()] 'Report of a series of 50 cases'. Izquierdo . *Acta Oncol* 1994. 33 (7) p. 767. (Male breast cancer)
- [Sasco et al. ()] 'Review article: epidemiology of male breast cancer. A metaanalysis of published case-control studies and discussion of selected aetiological factors'. A J Sasco , A B Lowenfels , -De Pasker , P Jong . *Int J Cancer* 1993. 20 (4) p. .
- [Vecchia ()] 'Risk factors for male breast cancer'. D'avanzo B & La Vecchia , C . *Br J Cancer* 1995. 71 (6) p. .
- [Johnson et al. ()] 'Risk factors for male breast cancer in Canada, 1994-1998'. K C Johnson , S Pan , Y Mao . *Eur J Cancer Prev* 2002. 11 (3) p. .
- [Ewertz et al. ()] 'Risk factors for male breast cancer—a case-control study from Scandinavia'. M Ewertz , L Holmberg , S Tretli , B V Pedersen , A Kristensen . *Acta Oncol* 2001. 40 (4) p. .
- [Lenfant-Pejovic et al. ()] 'Risk factors for male breast cancer: a Franco-Swiss case-control study'. M H Lenfant-Pejovic , N Mlika-Cabanne , C Bouchardy , A Auquier . *Int J Cancer* 1990. 45 (4) p. .
- [Bevier et al. ()] 'Risk of breast cancer in families of multiple affected women and men'. M Bevier , K Sundquist , K Hemminki . *Breast Cancer Res Treat* 2012. 132 (2) p. .
- [Tournier et al. ()] 'Significant contribution of germline BRCA2 rearrangements in male breast cancer families'. I Tournier , B B Paillerets , H Sobol , D Stoppa-Lyonnet , R Lidereau , M Barrois , S Mazoyer , F Coulet , A Hardouin , A Chompret , A Lortholary , P Chappuis , V Bourdon , V Bonadona , C Maugard , B Gilbert , C Nogues , T Frébourg , M Tosi . *Cancer Res* 2004. 15 (22) p. .
- [Chodick et al.] *Similar prevalence of founder BRCA1 and BRCA2 mutations among Ashkenazi and non-Ashkenazi men with breast cancer: evidence from 261 cases in Israel*, G Chodick , J P Struewing , Ron E Rutter , J L Iscovich , J . p. .
- [Nilsson et al. ()] 'Similarities and differences in the characteristics and primary treatment of breast cancer in men and women -a population based study (Sweden)'. C Nilsson , M Holmqvist , L Bergkvist , I Hedenfalk , M Lambe , M L Fjällskog . *Acta Oncol* 2011. 50 (7) p. .
- [Tirkkonen et al. ()] 'Somatic genetic alterations in BRCA2-associated and sporadic male breast cancer'. M Tirkkonen , T Kainu , N Loman , O T Jóhannsson , H Olsson , R B Barkardóttir , O P Kallioniemi , A Borg . *Genes Chromosomes Cancer* 1999. 24 (1) p. .
- [Dimitrov et al. ()] 'Some aspects of the endocrine profile and management of hormonedependent male breast cancer'. N V Dimitrov , P Colucci , S Nagpal . *Oncologist* 2007. 12 (7) p. .
- [Vetto et al.] *Stages at presentation, prognostic factors, and*, J Vetto , S Y Jun , D Paduch , H Eppich , R Shih .
- [Bloom et al. ()] 'Status of HER-2 in male and female breast carcinomas'. K J Bloom , H Govil , P Gattuso , V Reddy , D Francescatti . *Am J Surg* 2001. 82 p. .
- [Cutuli ()] 'Strategies in treating male breast cancer'. B Cutuli . *Expert Opin Pharmacother* 2007. 8 (2) p. .
- [Salehi et al. ()] 'Survival of male breast cancer in fars, South of Iran'. A Salehi , H Zeraati , K Mohammad , M Mahmoudi , A R Talei , A Ghaderi , M H Imanieh , A Fotouhi . *Iran Red Crescent Med J* 2011. 13 (2) p. .
- [Ioka et al. ()] 'Survival of male breast cancer patients: a population-based study in Osaka, Japan'. A Ioka , H Tsukuma , W Ajiki , A Oshima . *Jpn J Clin Oncol* 2006. 36 p. .

- 663 [Thalib and Hall ()] ‘Survival of male breast cancer patients: population-based cohort study’. & Thalib , Hall .
664 *Cancer Sci* 2009. 100 p. .
- 665 [Iredale et al. ()] ‘The experiences of men with breast cancer in the United Kingdom’. R Iredale , K Brain , B
666 Williams , E France , J Gray . *European Journal of Cancer* 2005. 42 p. .
- 667 [De Perrot et al. ()] ‘Thirty-year experience of surgery for breast carcinoma in men’. M De Perrot , J Deléaval ,
668 J Robert , A Spiliopoulos . *Eur J Surg* 2000. 166 (12) p. .
- 669 [Pant and Dutta ()] ‘Understanding and management of male breast cancer: a critical review’. & Pant , Dutta .
670 *Med Oncol* 2008. 25 (3) p. .
- 671 [Volpe et al. ()] ‘unilateral male breast masses: cancer risk and their evaluation and management’. C M Volpe ,
672 J D Raffetto , D W Collure , E L Hoover , R J Doerr . *Am Surg* 1999. 65 (3) p. .
- 673 [Donovan and Flynn ()] ‘What Makes a Man a Man? The lived experience of male breast cancer’. C Donovan ,
674 M Flynn . *Cancer Nursing* 2007. 30 (6) p. .
- 675 [Hecht ()] ‘Winchester DJ Male breast cancer’. J R Hecht . *Am J Clin Pathol* 1994. 102 (4) p. . (Suppl 1)