

# Effects of Adding Human Follicular Fluid and Pentoxifylline on IUI Outcome of Asthenozoospermic Patients

Saad S. Al-Dujaily<sup>1</sup>

<sup>1</sup> University of Baghdad

Received: 7 December 2013 Accepted: 3 January 2014 Published: 15 January 2014

## Abstract

Background: Although there are many motility stimulants have been found to play an important role in improving the active sperm motility and morphologically normal sperm (MNS) percentages when added in vitro with the culture medium for semen, the results of pregnancy rate following intra-uterine insemination (IUI) still not exceed 20

## Index terms—

## 1 Effects of Adding Human

Follicular

Abstract-Background: Although there are many motility stimulants have been found to play an important role in improving the active sperm motility and morphologically normal sperm (MNS) percentages when added in vitro with the culture medium for semen, the results of pregnancy rate following intra-uterine insemination (IUI) still not exceed 20% .

Objective: This study is a trial to improve the results of pregnancy rate following IUI by adding the follicular fluid (FF), and pentoxifylline (PX) to the culture medium used for preparation of asthenozoospermic semen samples.

Subjects, Material and methods: Ninety five infertile couples were involved in the current study (asthenozoospermic male partners with intact spouses) divided into three groups according to the sperm stimulants which were used for in vitro sperm activation followed by intrauterine insemination technique. Group one was FF-treated semen samples and consists of 40 couples, group two was FF+PX-treated semen samples and consists of 25 couples and lastly, group three was Ham's F-12 culture medium as a control group and consists of 30 couples. All the couples were followed up for determining outcome of pregnancy after IUI was done.

Results: Out of the all 95 couples, 22 pregnancies (

## 2 Introduction

Infertility is the inability of a sexually active noncontraceptive couple to achieve pregnancy through one year (1). It is a relatively common condition affecting approximately one in ten of the population. Infertility is either primary, when no pregnancy has ever occurred, or secondary, where there has been a pregnancy, regardless of the outcome. About 67-71% and 29-33% of patients have primary and secondary infertility, respectively (2). In half of these cases, a male factor is involved, making defective sperm function the largest single, defined cause of human infertility (3). Asthenozoospermia is one of the major causes of infertility or reduced fertility in men (4).

In vitro studies in human showed that spermatozoal accumulation into follicular fluid (FF) is significantly higher than into simple medium and that chemoattractant effect of fluid from an individual follicle correlates with the fertilizability of the egg from the same follicle (5,6).

Follicular fluid can also alter the physiology and behavior of spermatozoa by accelerating capacitation of the cell to induce sperm motility, increasing AR and signaling interaction between sperm and oocyte during fertilization.

This is due to FF components which are a low molecular hydrophobic compounds, platelet activating factor and progesterone (7,8) .

On the other hand, certain metabolic stimulants induce sperm capacitation and AR via changes in the values of cyclic adenosine monophosphate (cAMP) and intracellular calcium ( $[Ca^{2+}]_i$ ). Examples of these stimulators are pentoxifylline (PX) and L-carnitine (LC) (9) .

Pentoxifylline is a methylxanthine derivative which inhibits phosphodiesterase, thereby elevating the concentrations of intracellular cAMP and/or cyclic guanosine monophosphate (cGMP). It is used in ART to enhance the function of inherently poor quality spermatozoa from oligoasthenozoospermic patients (10) . Other study does not suggest any increase in teratogenicity or evidence of congenital malformations in pregnancies following the using of PX in the media of IVF cycles (9) .

The Artificial Insemination (AI) is the first option treatment for infertile couples with cervical factor subfertility, mild to moderate male subfertility and unexplained infertility (11) .IUI, with or without controlled ovarian hyperstimulation, is one of the treatment modalities offered most often to subfertile couples because it is less stressful, invasive and expensive than other interventions such as in vitro fertilization and gamete intrafallopian transfer (12)(13) . The goal of the present proposal is to examine the effects of adding human FF alone or in combination with PX on the

### 3 Materials and Methods

This study was conducted at the High Institute for Infertility Diagnosis and Assisted Reproductive Technologies, Al-Nahrain University through the period from September 2012 to June 2013.

Clinical examination was performed by a Consultant Urologist in charge of Male Infertility Unit in the Institute including presence or absence of varicocele, cryptorchidism, hydrocele, hernia and others. Alcohol consumption and smoking habits were also reported. Those patients were classified into three subgroups according to WHO criteria of seminal fluid analysis (1) .In this study ninety five infertile couples (asthenozoospermic male partners with intact spouses) divided into three groups according to the sperm stimulants which were used for in vitro sperm activation followed by intrauterine insemination techniques: 1. Group I: FF-treated semen samples. Include 40 couples. 2. Group II: FF and PX-treated semen samples.

Include 25 couples. 3. Group III: Ham's F-12 culture medium as control group. Include 30 couples.

### 4 Preparation of follicular fluid for in vitro sperm activation

Patients underwent oocyte retrieval 36 hr after hCG administration by transvaginal ultrasound guided follicular aspiration. The first aspirated follicle was usually the one with the largest diameter, and the first follicle allowed the collection of fluid without contamination of flushing medium. Specimens that were contaminated with blood were discarded .After removal of potential oocytes for treatment , the fluid was centrifuged ( $500 \times g$ ) to eliminate granulosa cells and to monitor the contamination of red blood cells and the clear supernatant was divided into aliquots and frozen at  $-20^{\circ}C$  (14) .The frozen samples were thawed and analyzed (15) . Solution of 20% hFF was prepared by the addition of 0.2 ml of hFF to 0.8 ml of Ham ' s F-12 medium.

### 5 b) Investigations to assess the female reproductive status

Physical examination was done following the case history. The investigations were performed to assess the normal menstruation and ovulation. Hormonal assay was done to the female namely; early follicular estradiol ( $E_2$ ), FSH and LH level with serial vaginal ultra-sonographic examination for monitoring of follicular development and endometrial thickness. Their tubal patency were checked from the previous cycles by hysterosalpingography. c) Intra-Uterine Insemination Technique IUI was performed when 0.5 -0.75 ml of prepared semen was aspirated into 1 ml syringe attached to endo-cervical IUI catheter (Gynetics, Belgium). In IUI room, the spouse was prepared for insemination in lithotomic position. Non lubricated Cusco's speculum was placed in the vagina to visualize the uterine cervix (16,17) . The specimen was slowly ejected from the syringe with gentle grasping of the cervical lips to prevent escape of the solution out of the cervix then after few minutes the catheter was removed slowly. The patient remains on supine position for 30 minutes then she allowed leaving the theater. Following the day of insemination the patient instructed to take progesterone tablets (Duphaston ® 10 mg/day) for two weeks as a luteal phase support. No use of antibiotic as it was detected that incidence of pelvic infection following IUI has been estimated to be less than 0.5% (17) . The diagnosis of pregnancy was done either by biochemical analysis (the detection of  $\beta$ hCG in blood after two weeks of insemination) or by sonographic examination to assess the pregnancy status

### 6 d) Statistical analysis

The data of this study was analyzed by using SPSS (Statistical package for social sciences) versions 16 and Microsoft excel 2013. Numeric variables were expressed as mean  $\pm$ SEM whereas nominal variables were expressed as numbers and percentage. The analysis was done by using one way analysis of variance (ANOVA) to compare among different groups of in vitro sperm activation techniques. Also, descriptive analysis was done by using Chi-square depending on the nature of the data. Differences between values of means were considered statistically significant at ( $P < 0.05$ ) (18) .

---

## 7 III. Results

### 8 a) Comparison of certain sperm function parameters after activation among IUI groups

In Table 1, there was no significant ( $P>0.05$ ) difference in the mean of sperm concentration, active sperm motility (grade A, B and A+B) and the percentage of MNS between semen samples activated in vitro by Ham's F12 medium only (control) and other treated media. 2 and 3 reveal the effect of three different sperms stimulants (FF, FF+PX and Ham's F-12) on pregnancy rate and percentage (%) following IUI. Out of the all 95 couples, 22 pregnancies (23.15%) were resulted following IUI. The best results were obtained by group 1(FF treated group) in which out of 40 cases enrolled in this group, 12 pregnancies (30%) were resulted following IUI. Regarding group 2 (FF+PX treated group),out of 25 cases enrolled in this group, 5 women (20%) were conceived following IUI. Regarding group 3 (Ham's F-12 treated group), out of 30 cases enrolled in this group, 5 women (16.67%) were conceived following IUI.

## 9 Volume XIV Issue

There was a significant ( $P<0.05$ ) difference in the rate of pregnancy between group 1 and group 2 (table 4-21), while there was no statistically significant ( $P>0.05$ ) difference in the rate of pregnancy between group 1 and 3. Year 2014

## 10 E

IV.

## 11 Discussion

The current study demonstrates that IUI by a sperm washed in vitro using a medium supplemented with FF results in a higher pregnancy rates per cycle (30%) in asthenozoospermic patients than by using Ham's F-12 medium alone (16.6%) even the results of certain sperm function parameters after in vitro preparation did not reveal any significant between them. This finding can be explained by the fact that the extent of hyperactivated motility produced by FF is positively correlated with the extent of zona binding, the acrosome reaction, zona-free oocyte penetration and fertilizing capacity in vitro (19). Additionally, human FF is an efficient capacitating agent and only the capacitated sperm are able to migrate towards the ovum site, under thermotaxis and chemotaxis stimuli (20,21).

It has been reported that FF may modulate spermatozoa function to be indicative of pregnancy outcome (22), this fact can verify and confirms the original results.

The contents of FF also play a crucial role in the pregnancy outcome. A lower activity of the conversion enzymes, ovarian 11 beta hydroxy steroid dehydrogenase (HSD11?) which is found in FF interconverts 'active' cortisol to 'inert' cortisone, have been associated with positive pregnancy outcomes for patients undergoing ART (23).

In this study, there was a significant ( $P<0.05$ ) difference in the rate of pregnancy between IUI-FF treated group (pregnancy rate was 30%) and IUI-FF+PX treated group (pregnancy rate was 20%).This finding suggests that IUI following in vitro sperm activation by FF containing culture media is associated with a higher pregnancy outcome than FF+PX containing media. Moreover, the proper time of IUI that performed enclosed with ovulation time can lead to increase in pregnancy rates, and is best seen in patients with pure andrological indication. At the same time, careful selection of patients might play part in achieving success with IUI. Furthermore, the use of fertility drugs like gonadotropins in the controlled ovarian stimulation followed by IUI provide better pregnancy rate (24) than the using of other ovarian stimulants.

The current study found that pregnancy rate of IUI following FF+PX (20%) was approximately similar to the results of Al-Dujaily et al (20.8%) using PX alone (25).

Both results had no significant different in pregnancy outcome than the pregnancy outcome following in vitro sperm activation with FF alone. This suggest that there is no further advantage to add PX to FF-containing culture media for the enhancement of active sperm motility, morphologically normal sperm percentages or pregnancy rate following IUI.

Thus the present study concluded that adding the FF to medium used for IUI can enhance the results of pregnancy rate of infertile couples. <sup>1</sup>



Figure 1:

1

Year

Figure 2: Table 1 :

2

Status of pregnancy		Group 1(FF treated)		Pregnancy Rate Group 2 (FF+PX treated)	
		No.	%	No.	%
Pregnant	Yes	12	30	5	20
	No	28	70	20	80
	Total	40	100	25	100
X2 = 9.474					
DF = 1		P < 0.05			

Figure 3: Table 2 :

Status of Pregnancy		Pregnancy rate			
		Group 1 (FF treated)		Group 3(Ham's F-12 treated)	
		No.	%	No.	%
Pregnancy	Yes	12	30	5	16.67
	No	28	70	25	83.33
	Total	40	100	30	100
X2 = 1.656					
DF = 1					
		P			
		>0.05			

Figure 4: Table 3 :



- 
- [Casadei et al.] , L Casadei , V Zamaro , M Calcagni .
- [Steiber et al. ()] ‘Carnitine: a nutritional, biosynthetic and functional perspective’. A Steiber , J Kemer , C Hoppelm . *Mol. Aspects Med* 2004. 25 (5-6) p. .
- [Bjorndahl et al. ()] ‘Contamination by seminal plasma factors during sperm selection’. L Bjorndahl , M Mohammadi , M Pourian . *J Androl* 2005. 2 p. .
- [Bedaiwy ()] ‘Effect of follicular fluid oxidative stress parameters on intracytoplasmic sperm injection outcome’. M Bedaiwy . *Gyne Endocr* 2012. 28 (1) p. .
- [Al-Dujaily and Alnakash ()] ‘Effect of pentoxifylline on the outcome of artificial inseminations’. S S Al-Dujaily , A H Alnakash , Al-Bayati , S . *Iraqi Post Med J* 2006. 5 (3) p. .
- [Wen et al. ()] ‘Estradiol, progesterone, testosterone profiles in human follicular fluid and cultured granulosa cells from luteinized preovulatory follicles’. X Wen , D Li , A J Tozer . *Reprod Biol Endocrinol* 2010. 8 p. 117.
- [Revelli et al. ()] ‘Follicular fluid content and oocyte quality: from single biochemical markers to metabolomics’. L D Revelli , S Casano , E Molinaril , Massobrio M Rinaudo , P . *Reprod Biol and Endocr* 2009. 7 p. .
- [Homologous intrauterine insemination in controlled ovarian hyperstimulation cycles: A comparison among three different regimens] ‘Homologous intrauterine insemination in controlled ovarian hyperstimulation cycles: A comparison among three different regimens’. *European J. Obstet Gynecol Reprod Bio* 2006. 129 p. .
- [Andersen et al. ()] ‘Hormonal characteristics of follicular fluid from women receiving either GnRH agonist or hCG for ovulation induction’. C Y Andersen , P Humaidan , H Bredkjaer . *Hum Reprod* 2006. 21 (8) p. .
- [Dickey et al.] *Insemination technique and insemination complications*, R P Dickey , P R Brinsden , R Pyrzak .
- [Verhulst et al. ()] ‘Intra-uterine insemination for unexplained subfertility’. S M Verhulst , B J Cohlen , E Hughes , Te Velde , E Heineman , MJ . *Cochrane Database of Systematic Reviews* 2006. (4) .
- [Intrauterine insemination and ovulation induction ()] *Intrauterine insemination and ovulation induction*, 2010. p. .
- [Koli et al. ()] ‘Intrauterine insemination: a retrospective review on determinants of success’. P Koli , M Anil , K Patil . *Int J Reprod Contracept Obstet Gynecol* 2013. 2 (3) p. .
- [Glover and Mitchel ()] *Introduction to Biostatistic. 2 nd ed*, T Glover , K Mitchel . 2008. 4.
- [Abdelrazik and Agrawal ()] ‘L-carnitine and assisted reproduction’. H Abdelrazik , A Agrawal . *Arch Med Sci* 2009. 5 p. .
- [Laboratory manual for the examination and processing of human semen ()] *Laboratory manual for the examination and processing of human semen*, 2010. Geneva, Switzerland. 5. WHO (th ed.)
- [Dohle ()] ‘Male infertility in cancer patients: Review of the literature’. G R Dohle . *Int.J.Urol* 2010. 17 (4) p. .
- [Leroy et al. ()] ‘Metabolic changes in follicular fluid of the dominant follicle in high-yielding dairy cows early postpartum’. J L Leroy , T Vanholder , J R Delanghe . *Theriogenology* 2004. 62 p. .
- [Machado et al. ()] ‘Mobilisation of Ca<sup>2+</sup> stores and flagellar regulation in human sperm by S-nitrosylation: a role for NO synthesised in the female reproductive tract’. G Machado , L Lefievre , C Ford . *Dev J* 2008. 135 (22) p. .
- [Aitken et al. ()] ‘New insights into sperm physiology and pathology’. R J Aitken , M A Baker , B Nixon . *Hand book Pharmacol* 2010. 198. p. .
- [Michael et al. ()] ‘Ovarian 11 beta hydroxysteroid dehydrogenase activity is inversely related to the outcome of in vitro fertilization-embryo transfer treatment cycles’. A Michael , L Gregory , E Piercy . *Fertil.Steril* 1995. 64 (3) p. .
- [Cooper (ed.) ()] *Physiology of Sperm Maturation and Fertilization*, T G Cooper . Nieschlag E BH (ed.) 2010. Berlin Heidelberg: Springer-Verlag. p. . (Andrology: Male Reprod Health and Dysfunction. 3rd ed)
- [Merviel et al. ()] ‘Predictive factors for pregnancy after intrauterine insemination’. P Merviel , M Heraud , N Grenier . *Ferti Steril* 2008. 10 p. .
- [Vm and Wilson ()] ‘Pregnancy outcomes after assisted reproductive technology’. Allen Vm , R D Wilson . *J Obst Gyne* 2006. 28 (3) p. .
- [Guidobaldi et al. ()] ‘Progesterone from the cumulus cells is the sperm chemoattractant secreted by the rabbit oocyte cumulus complex’. H A Guidobaldi , M E Teves , D R Unates . *PL J* 2008. 3 p. .
- [Eisenbach and Giojalas ()] ‘Sperm guidance in mammals -an unpaved road to the egg’. M Eisenbach , L C Giojalas . *Nat Rev Mol Cell Biol* 2006. 7 (4) p. .
- [Bahat and Eisenbach ()] ‘Sperm thermotaxis’. A Bahat , M Eisenbach . *Mol Cell Endocrinol* 2006. 252 (1-2) p.