



GLOBAL JOURNAL OF MEDICAL RESEARCH: F  
DISEASES

Volume 14 Issue 2 Version 1.0 Year 2014

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Scurvy in a Child with Cerebral Palsy- The Forgotten Vitamin Deficiency: A Case Report

By Dr. Raghavendra. K, Dr. Suresh Babu., Dr. Basavanthappa., Dr. Srinivas.V,  
Dr. Srinivas Murthy. C.L, Dr. Rajath Pejaver & Dr. Nikil. P. T

*Basaveshwara Medical College and Hospital, India*

*Abstract-* Scurvy was the first vitamin deficiency disease to be described. However it is seldom seen in the pediatric age group. It is often missed, especially amongst physically and mentally disabled patients who form a high risk group for this disease. Scurvy can present with a varied spectrum of signs and symptoms. Here we present a case of scurvy in a child with cerebral palsy.

*GJMR-F Classification : NLMC Code: WD 140, WD 100, WD 105*



*Strictly as per the compliance and regulations of:*



© 2014. Dr. Raghavendra. K, Dr. Suresh Babu., Dr. Basavanthappa., Dr. Srinivas.V, Dr. Srinivas Murthy. C.L, Dr. Rajath Pejaver & Dr. Nikil. P. T. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Scurvy in a Child with Cerebral Palsy- The Forgotten Vitamin Deficiency: A Case Report

Dr. Raghavendra. K <sup>α</sup>, Dr. Suresh Babu. <sup>σ</sup>, Dr. Basavanthappa. <sup>ρ</sup>, Dr. Srinivas. V <sup>ω</sup>,  
Dr. Srinivas Murthy. C.L <sup>¥</sup>, Dr. Rajath Pejaver <sup>§</sup> & Dr. Nikil. P. T <sup>x</sup>

**Abstract-** Scurvy was the first vitamin deficiency disease to be described. However it is seldom seen in the pediatric age group. It is often missed, especially amongst physically and mentally disabled patients who form a high risk group for this disease. Scurvy can present with a varied spectrum of signs and symptoms. Here we present a case of scurvy in a child with cerebral palsy.

## I. INTRODUCTION

A deficiency of Vitamin C (ascorbic acid) results in the clinical presentation of Scurvy, the oldest nutritional deficiency to be recognized. A disease that was once rampant is now rarely seen, more so in the pediatric age group.

Scurvy presents with swelling of joints with characteristic radiological changes, gum bleeds, anemia, petechiae (perifollicular), muscle weakness, fractures and poor wound healing. Here we report a case of scurvy in a child with cerebral palsy and developmental delay.

## II. CASE REPORT

A 18 month-old boy with quadriplegic cerebral palsy and pseudobulbar palsy due to perinatal asphyxia was admitted to the department of Pediatrics with history of swelling and pain of right knee joint with gum bleeds of 7 days duration. Child also had excessive irritability especially when picked up, along with fever. There was no history of trauma. The child was on a predominant milk based diet, with minimal intake of fruits and vegetables. He was on long-term phenytoin and phenobarbitone therapy for seizures.

On examination, the child had acute malnutrition (wt= 6.5kg, IAP grade 3) and microcephaly (HC- 34 cm). He was febrile, pale and had no hepatosplenomegaly or lymphadenopathy. The right knee joint was swollen and tender, with the skin on the joint appearing shiny, red and warm. There was minimal movement of the right lower limb. A possibility of septic arthritis was considered and intravenous antibiotic therapy initiated. Orthopedic opinion was sought and the limb was immobilized with a POP cast. The laboratory data results were as follows:

Hb:10.6 g/dl ; TC:7200 cells/mm<sup>3</sup> ; DC: P55%,L42%,E3%; ESR: 45 mm/hr;RBC Count: 4.7 million/mm<sup>3</sup>;MCV:68.5 fl;MCH:22.6 pg; MCHC:32.9 %; Platelet :2.31 Lakh/mm<sup>3</sup> ; Calcium: 9.2 mg/dl; S.Alkaline Phosphatase:102 IU/dl ;S.Phosphate: 3.7mg/dl.

The radiograph of the knee (Figure 1) showed: Ground glass appearance of the shaft of the tibia, fibula and femur. White line of Frankel (irregular, thickened white line at the metaphysis) and a characteristic zone of rarefaction under the white line at the metaphysis (Trummerfeld zone). A lateral prolongation of the white line at the cortical ends, known as Pelkan spur was seen. Subperiosteal elevation suggestive of a subperiosteal hemorrhage was seen at the lower end of the femur. All radiological features pointed towards scurvy. The diagnosis was confirmed with serum levels of vitamin C being less than the lower limit of normal. The child also had low levels of vitamin D possibly due to lack of sunlight exposure, dietary deficiencies and chronic anticonvulsant therapy. ( Vitamin C: 0.5mg/dl ; 25 OH-Vitamin D: <3micg/ml ).



Fig 1 : Child with POP cast , malnutrition and microcephaly

Author <sup>α</sup> <sup>σ</sup> <sup>ρ</sup> <sup>ω</sup> <sup>¥</sup> <sup>§</sup> <sup>x</sup>: Department of Pediatrics, Basaveshwara Medical College & Hospital, Chitradurga. – 577501, India.  
e-mail: rajath.pejaver@gmail.com



Fig. 3,4 : X-Ray AP & Lateral of knee joint showing Pelkan spur & white line of Frankel (arrows), Trummerfeld zone

### III. DISCUSSION

The diagnosis of scurvy was made, and the child was treated with 250mg of vitamin C daily. Vitamin D 6 lakh IU was also administered. His mother was educated about dietary modification. Two weeks after vitamin C administration, the child's general condition and joint swelling improved. Repeat X-ray of the knee joint showed features suggestive of healing.



Fig. 4 : X ray of knee joint showing healing

Scurvy is less common in the pediatric population, but case reports still appear [1–3]. A review of the literature by Noble et al. reveals twenty three case

reports of scurvy in children with behaviourally restricted diets including children with autism, mental retardation and cerebral palsy [4]. Scurvy is common in children with cerebral palsy as they subsist on predominant milk based diets (due to pseudobulbar palsy and difficulty swallowing solids) and boiled cows milk is a very poor source of vitamin C. deficiencies may be noted in preterm babies who are on prolonged TPN therapy, children with malnutrition and those with acute illnesses. Musculoskeletal manifestations are present in 80% of patients with scurvy and are prominent in pediatric population [3, 5]. Musculoskeletal manifestations include sub-periosteal hemorrhages leading to bone pain and musculoskeletal complaints such as limb pain, limping, swelling over long bones, and progressive leg weakness and fractures [6]. Dermatological manifestations include petechiae, ecchymoses, hyperkeratosis, and perifollicular hemorrhage [3, 7]. Oral symptoms include gingival disease characterized by swelling, bleeding gums, and loosening of teeth [3, 6, 8]. Systemic symptoms of scurvy in children include lassitude and fatigue, failure to gain weight, loss of appetite, and irritability [6]. In addition to these symptoms, deficiency of ascorbic acid may lead to a hypochromic microcytic anemia because of decreased absorption of iron, bleeding, and dietary deficiencies [3, 6].

The diagnosis of scurvy is based on history of poor dietary intake of vitamin C, classic clinical features and radiological findings and response to treatment with vitamin C. [3, 14]. Weinstein et al. [3] recommend oral doses of 100 to 300 mg of vitamin C daily until body stores are replenished per serum levels. Daily fruit and vegetable intakes should include a good source of

vitamin C such as citrus fruits, berries, green leafy vegetables and vegetables of brassica and crucifera family. Once a regimen of vitamin C is begun, improvement of symptoms usually begins in 24 hours, with pain diminishing in two to four days, and gingival lesions recovering in two to three weeks [6]. With vitamin C supplementation, metaphyseal abnormalities of scurvy will completely resolve [9]. The large shells of periosteal bone are common radiographic findings particularly during the healing phase of disease [12].

Various factors contribute to nutritional deficiencies in non ambulant children with severe spastic cerebral palsy like poor intake, oral motor dysfunction, feeding problems, and use of antiepileptic drugs [13].

11. D. Front, R. Hardoff, J. Levy, and A. Benderly, "Bone scintigraphy in scurvy," *Journal of Nuclear Medicine*, vol. 19, no. 8, pp. 916–917, 1978.
12. D. Resnick, *Hypervitaminosis and Hypovitaminosis. Diagnosis of Bone and Joint Disorders*, WB Saunders, Philadelphia, Pa, USA, 4th edition, 2002.
13. R. C. Henderson, P. P. Lin, and W. B. Greene, "Bone-mineral density in children and adolescents who have spastic cerebral palsy," *Journal of Bone and Joint Surgery. American*, vol. 77, no. 11, pp. 1671–1681, 1995.
14. *Nelson Textbook of Pediatrics*, 19th Edition, page no 198 -200.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. M. Larralde, A. S. Muñoz, P. Boggio, V. Di Gruccio, I. Weis, A. Schygiel, "Scurvy in a 10-month-old boy," *International Journal of Dermatology*, vol. 46, no. 2, pp. 194–198, 2007.
2. P. Rosati, R. Boldrini, and R. Boldrini, "A child with painful legs," *The Lancet*, vol. 365, no. 9468, p. 1438, 2005.
3. M. Weinstein, P. Babyn, and S. Zlotkin, "An orange a day keeps the doctor away: scurvy in the year 2000," *Pediatrics*, vol. 108, no. 3, p. E55, 2001.
4. J. M. Noble, A. Mandel, and M. C. Patterson, "Scurvy and rickets masked by chronic neurologic illness: revisiting "psychologic malnutrition"?" *Pediatrics*, vol. 119, no. 3, pp. e783–e790, 2007.
5. K. Rajakumar, "Infantile scurvy: a historical perspective," *Pediatrics*, vol. 108, no. 4, p. E76, 2001.
6. D. Popovich, A. McAlhany, A. O. Adewumi, and M. M. Barnes, "Scurvy: forgotten but definitely not gone," *Journal of Pediatric Health Care*, vol. 23, no. 6, pp. 405–415, 2009.
7. C. J. Burk and R. Molodow, "Infantile scurvy: an old diagnosis revisited with a modern dietary twist," *American Journal of Clinical Dermatology*, vol. 8, no. 2, pp. 103–106, 2007.
8. S. Chatproedprai and S. Wananukul, "Scurvy: a case report," *Journal of the Medical Association of Thailand*, vol. 84, no. 1, pp. S106–S110, 2001.
9. C. P. Duggan, S. J. Westra, and A. E. Rosenberg, "Case 23-2007: a 9-year-old boy with bone pain, rash, and gingival hypertrophy," *The New England Journal of Medicine*, vol. 357, no. 4, pp. 392–400, 2007.
10. F. G. Riepe, D. Eichmann, H. C. Oppermann, et al., "Special feature: picture of the month. Infantile scurvy," *Archives of Pediatrics and Adolescent Medicine*, vol. 155, pp. 607–608, 2001.