Artificial Intelligence formulated this projection for compatibility purposes from the original article published at Global Journals. However, this technology is currently in beta. *Therefore, kindly ignore odd layouts, missed formulae, text, tables, or figures.*

¹ Use of Transcatheter Arterial Embolization in High Risk Patients ² with Upper Gastrointestinal Bleeding

Asiri Arachchi

Received: 8 December 2013 Accepted: 3 January 2014 Published: 15 January 2014

6 Abstract

3

Δ

Introduction-Upper gastro intestinal bleeding is a common presentation to the Emergency 7 Department; approportion of these bleeds are secondary to duodenal ulcers. The management 8 of these young, fit patients who can tolerate a second bleed physiologically is well 9 established. However in the elderly frail patient a second bleed may be terminal. Options for 10 treatment for these blee ding duodenal ulcers (apart from resuscitation, including transfusion 11 if need be and medical management with intra venous Proton Pump Inhibitors), include 12 endoscopic control, laparotomy with under running of the bleeding artery and endovascular 13 approaches. 98 percent of patients can be successfully managed with endoscopy and 14 epinephrine injection, coagulation and / or clipping of the bleeder. The dilemma exists in 15 those patients whom are high risk for surgical intervention (laparotomy) should a re-bleed 16 occur. In these patients transcatheter arterial embolization has become a favourable 17 treatment modality and especially in high risk patients, selective transcatheterembolisationhas 18 been established as a safe approach (1,2,3,5,9). 19

21 Index terms—

20

²² 1 Introduction

pper gastro intestinal bleeding is a common presentation to the Emergency Department; aproportion of these bleeds are secondary to duodenal ulcers. The management of thesein young, fit patients who can tolerate a second bleed physiologically is well established. However in the elderly frail patient a second bleed may be terminal.

Options for treatment for these bleeding duodenal ulcers (apart from resuscitation, including transfusion if need be and medical management with intra venous Proton Pump Inhibitors), include endoscopic control, laparotomy with under running of the bleeding artery and endovascular approaches. 98 percent of patients can be successfully managed with endoscopy and epinephrine injection, coagulation and / or clipping of the bleeder. The dilemma exists in those patients whom are high risk for surgical intervention (laparotomy) should a re-bleed occur. In these patients transcatheter arterial embolization has become a favourable treatment modality and especially in high risk patients, selective transcatheterembolisationhas been established as a safe approach (1,2,3,5,9).

In this case report we discuss the management of a 98 year old male with a duodenal ulcer with stigmata of recent bleed who presented to our department in a rural setting in Victoria, Australia.

35 **2 II.**

36 3 Case Study

Mr A, a 98 year old male (with a background history of hypertension, a newly inserted pacemaker and not on any anti-coagulation), presented to our Emergency Department with epigastric pain, melena and a Haemoglobin of 68. He was cardiovascularly unstable and following resuscitation (including transfusion) was taken for endoscopy which demonstrated a large duodenal ulcer with a fresh clot and a visible artery. This was managed with peri ulcer injectionof (adrenaline). Attempts to place a clip on the artery was unsuccessful due to the position. 42 Author ? ?: Department of General Surgery Bairnsdale Regional Health Services PO Box 474, Bairnsdale 43 VIC 3875. AsiriArachchi MBBS. PDipSurg Anatomy. Msc (surgery), Servaise De Kock. MBBS. FRACS. He was 44 closely observed clinically and his haemoglobin was followed daily. His haemoglobin dropped again on day three 45 and he was again transfused and taken for endoscopy which now demonstrated a healed ulcer with no stigmata 46 of rebleed (photo 2).He was again managed conservatively and on day eight again dropped his haemoglobin.

49 the gastro duodenal artery, (image 3), with good outcome.

50 **4** III.

51 Discussion and Literature Review Our patient was undoubtedly not fit for a laparotomy and in these patients 52 with multiple medical comorbidities embolization therapy is a safe, effective treatment modality and has a good 53 prognosis.

Since its introduction in 1972 embolotherapy has become a possible modality in treatment of patients with upper gastrointestinal bleeding. Angiography involves either a brachial or femoral approach. Identification of the appropriate anatomy is important; particularly the coeliac axis, the common hepatic and delineating the anatomy involving the gastro duodenal artery, splenic, short gastric, left gastric. In general dynamic angiography via use of selective catheters can denote extravasation of contrast and thus ability to use either coils or particles to catheterize bleeding points as was the case with our patients (image 4 and 5). This illustrates a dynamic angiogram demonstrating the coeliac axis, regional anatomy and post embolization images as described in the

61 literature. (3,6,10).

There have been repeated discussions with importance of placing clips via initial endoscopy at the vicinity of the ulcer to aid the radiologist to identify the location of the bleeding vessel. This is an important pre adjuvant to formal transcatheterembolization and even though the artery may not be assessable to clipping itself in our case, would have been helpful if placed close to the ulcer and in future we would attempt to do this in selective

66 cases.

There is extensive literature denoting the success with clinical documented rates of close to 100 percent, with very minimal procedure related complications.Procedure complications relate to type of device used for transcatheterembolization.Main complications (1 to 8%) relate to in advent coiling of other vessels which can result in either re bleeding or secondary duodenal stenosis due to embolisation of muscular branches. (8,11,12).

Our patient had coiling of the gastro duodenal artery, but particles and spongesare also adjuvant options which interventional radiologists do use.

Retrospectively it was thought that the gastroduodenal artery was not noted on the second endoscopy because it was hiding behind the ridge of the ulcer. Therefore it is recommended to attempt to inspect the whole ulcer bed. If this was noted at this time we would have referred the patient earlier for embolization.

76 IV.

77 5 Conclusion

Within a rural hospital setting with no onsite access to endovascular embolization early recognition of the unwell,
high risk patient that needs embolization is pertinent. Endoscopy still plays a major role in the management
of these patients. It is important to inspect the whole ulcer bed for a visible vessel and be aware that it may

be hiding behind the ridge of the ulcer. Placement of a clip near the ulcer can be helpful to the interventional
 radiologist to identify the gastro-duodenal artery.

Advances in arterial embolization procedures have improved the outcome of these high risk patients and access to transport measures in a rural setting is of upmost importance.

Early recognition of the high risk bleeding ulcer in the high risk patient is important to enable timely referral for selective angiography and embolization.¹



Figure 1: Image 1:

86

 $^{^{1}}$ © 2014 Global Journals Inc. (US)

	assessment. J Vascular Interventional Radiology.			
	2001; 12:195-200.			
	2. Encarnacion CE, Kadir S, Beam CA, Payne CS.			
	Gastrointestinal	bleeding:		
	gastrointestinal arterial embolization. Radiology.	0		
	1992: 183:505-508.			
	3. Eriksson LG, Liungdahl M, Sundbom M, Nyman R.			
	Transcatheter arterial embolization versus surgery in			
	the treatment of upper gastrointestinal bleeding			
	after therapeutic endoscopy failure. J Vasc			
	Interventional Radiology, 2008: 19:1413-1418.			
	4. Goldman ML, LAND WC, Bradley EL, Anderson RT.			
Ye	ea Franscatheter therapeutic embolization in the management of massive u	pper gastr	ointestinal blee	ding. R
20)14	ppor 80001	0111005011101 0100	
	(2011) Empiric transcatheter arterial embolization for			
	massive bleeding from duodenal ulcers: efficacy			
	and complications. J Vascular Interventional			
	Radiology 22:911-916.			
	6. Lieberman DA, Keller FA, Katon RM et al (1984)			
	Arterial	emboliza	tion for	
	gastrointestinal tract bleeding in poor surgical	onnooniza		
	candidates I Gastroenterology 86:876-885			
	7 Loffroy B Bao P Ota S et al (2010) Embolization of			
	acute	nonvai	riceal	unner
	hemorrhage resistant to endoscopic treatment.	1101174	licear	upper
	results and predictors of recurrent bleeding			
	Cardiovascular		Interventional	
	1088-1100		meer ventional	
т	8 P. Chevallier, S. Novellas, G. Vanhiervliet et al. "Transcatheter embe	lization fo	r endosconicall	y unma
()	o. 1. Chevanier, 5. Novenas, G. Vanbiervilet et al., Transcatheter embe		r endoscopican.	y umma
)	gastrointestinal		hemorrhage."	
	Radiologie, vol. 88, no. 2, pp. 251-258, 2007.			
	9 K	Palmer	"Acute	upper
	haemorrhage."	·	110400	appor
	naomorrago,			

Figure 2:

5 CONCLUSION

- [Aina et al.] Arterial embolotherapy for upper gastrointestinal hemorrhage: outcome, R Aina , V L Oliva , E Therasse , P Perreault , B T Bui , M P Dufresne , G Soulez .