



GLOBAL JOURNAL OF MEDICAL RESEARCH: A
NEUROLOGY AND NERVOUS SYSTEM
Volume 14 Issue 3 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Cerebral Metastasis: Case Series Cerebrum as A Safe Haven

By Dr. Ashfaq ul Hassan, Ghulam Hassan, Dr. Pervez Shah, Dr. Masood Tanver,
Dr. Obaid MD Radiology, Dr. Muneeb ul & Dr. Zahida Rasool
Medicine SMHS Hospital Srinagar Kashmir, India

Abstract- Cerebral metastasis may be a common manifestation of systemic diseases. Cerebral metastasis are reported in 15-25 percent of patients with Brain tumors. Parenchymal blood flow is an important regulator of metastasis and most of the metastatic lesions are found in the cerebrum .most brain metastasis arise by hematogenous dissemination followed by dessimination through CSF.

Keywords: *metastasis, cerebral, cerebellum, MRI, disseminate, microinvasion.*

GJMR-A Classification : *NLMC Code: WS 342*



Strictly as per the compliance and regulations of:



Cerebral Metastasis: Case Series Cerebrum as A Safe Haven

Dr. Ashfaq ul Hassan ^α, Ghulam Hassan ^σ, Dr. Pervez Shah ^ρ, Dr. Masood Tanver ^ω,
Dr. Obaid MD Radiology [¥], Dr. Muneeb ul [§] & Dr. Zahida Rasool ^χ

Abstract- Cerebral metastasis may be a common manifestation of systemic diseases. Cerebral metastasis are reported in 15-25 percent of patients with Brain tumors. Parenchymal blood flow is an important regulator of metastasis and most of the metastatic lesions are found in the cerebrum .most brain metastasis arise by hematogenous dissemination followed by dissemination through CSF.

Keywords: metastasis, cerebral, cerebellum, MRI, disseminate, microinvasion.

I. INTRODUCTION

Meninges are the layers of connective tissue covering the brain and spinal cord The meninges consist of three membranous layers. Duramater: outer most, Arachnoid: middle layer: Piamater: inner most. Duramater is the thickest and toughest membrane covering the brain and consists of two layers. Endosteal layer: outer. Serves as internal periosteum, (endocranium). Meningeal layer: inner:

provides the protective membrane to brain. These two layers are fused to each other except where venous sinuses are enclosed between them.the cerebral metastatic lesions can be defin

Supratentorial and infratentorial. The supratentorial usually cause seizures, cognitive defects and headaches. The infratentorial lesions usually cause ataxia, Diplopia, dysarthria or brain herniation.

II. TEXT

a) Case 1

A 45 year old male presented in emergency Department with gradual onset headache ,. There was no past history of hypertension, diabetes, Sinusitis, pyrexia. The patient was able to move her extremities, Cranial nerves were normal on examination and Brain Stem Reflexes were normal. CT showed metastatic lesions in Occipital lobe

Symptom	Percentage of Patients / 30 patients
• Headache:	65
• Seizure:	12
• Ataxia:	5
• Weakness:	7
• Nausea/vomiting:	25
• Papilledema:	8
• Visual changes:	6
• Others(Syncope,dizziness):	10

Chart 1 : Showing Symptomatology Percentage of Presentation

Author^α: Clinical Anatomy Sheri Kashmir Institute of Medical Sciences College Bemina, Srinagar Kashmir India.

e-mail: ashhassan@rediffmail.com

Author^σ: Clinical Anatomy Sheri Kashmir Institute of Medical Sciences College Bemina, Srinagar Kashmir India.

Author^ρ: Medicine SMHS Hospital Srinagar Kashmir, India.

Author^ω: SMHS Hospital Srinagar Kashmir, India.

Author[¥]: SKIMS Soura.

Author[§]: Physician Directorate of Health Services, Srinagar Kashmir India.

Author^χ: Medical Consultant IUST Awantipora, Srinagar, Kashmir, India.

b) Case 2

A 45 year old woman presented with Breast Cancer and After several months with headache,

anorexia. Cranial nerves were normal on examination and Brain Stem Reflexes were normal. CT showed metastasis in occipital and frontal lobe.




Chart 2 : Showing Primary Sites

c) Case 3

A 48 year old man with thyroid cancer with CT showed metastasis in occipital lobe.




Chart 3 : Supratentorial Vs Infratentorial

III. EPIDEMIOLOGY

Brain Metastasis are common tumors of Brain.¹ Any tumor can spread to the brain. But the most common tumors spreading to the brain have been noticed to be lung, breast, melanoma, renal cell carcinoma, lymphoma, leukemias, thyroid, colorectal, from unknown sources.^{2,3,4,4,6,7,8}

The high rates of pulmonary disease spreading to Brain has been well documented. The tumor reaches the brain via hematogenous route. The tumor cells circulate in blood, to the left side of the heart and are carried by way of the carotid vessels to the circle of Willis from where they metastasize to different parts depending upon the dynamics of blood flow. The zones where the blood flow is considerably reduced along with narrowing of vessel diameter causes aggregation of tumor cells in these areas and represents an embolic source of disease. The parts of brain effected are the cerebrum, cerebellum and the brain stem.

In addition to the blood flow the soil seed theory of a cancer proliferation from the site of lodgement of malignant cells away from a primary source where the tumor cells grow and disseminate is important in case of cerebral metastasis and here the mitotic activity of cells is very important for rates of growth. The big and large metastasis are due to rapid cell growth in contrast to smaller metastatic lesions. The micromechanisms involved follow a cascade of intravasation, dissemination, extravasation and colonization.^{9,10,11}

In a survey conducted from multiple hospitals from the state it was seen that most of the cases of metastasis were from lungs, others from breast in case of females and from thyroid. A large number of cases were from other cancers.

The patients usually present with headache, seizures, vomiting, alterations of mental status, visual alterations. however patients can be asymptomatic as well. Headache is the most common symptom in case of cerebral metastasis. In adults the lung cancer followed by breast cancer and melanoma are the most common sources of primary tumors. Interestingly in many cases the intracranial involvement may be the first presentation. The rates of dissemination of the tumors in pediatric age group are significantly lower than that found in adults. The main problem with the cerebral metastasis is that the tumour size is not of much significance as even small lesions can cause considerable neurological sequelae. Sometimes the patients can also present with a haemorrhage inside the metastasis. This may produce a sudden headache, coma or a severe focal neurodeficit. Metastasis from melanomas, thyroid cancers and choriocarcinomas are particularly prone for hemorrhage. Usually in about one third of cases, patients with metastasis are diagnosed on routine investigations.

IV. RADIOLOGY

At present MRI and CT remain to be the most cost effective and non invasive techniques for detection of cerebral metastasis. Contrast enhancement can add finer details in the form of disruption of Blood Brain Barrier.¹⁴

Typically the metastasis are well demarcated from the surrounding parenchyma. There may be peritumoral edema as well. The radiographic features differ and there is a lot of variability among these tumors. the metastasis may be solitary or multiple. Mostly the metastasis are multiple.

On Pre contrast imaging CT Scanning may show iso dense or hypodense lesions. there may be vasogenic edema. following administration of contrast media, the enhancement may be nodular, punctate or ring enhancing. the main site of primary is the lung, breast, melanoma, renal cell carcinoma.¹⁵

The metastasis should be differentiated from primary brain tumors, cerebral abscesses, stroke, radiation necrosis, granulomatous brain lesions, demyelination and infarcts. The clinical implication in general is that the prognosis in a patient with cerebral metastasis is generally poor. Over the past few years whole brain radiotherapy has been considered as a standard treatment. Stereotactic radiotherapy has been introduced lately and is proving to be effective. surgery in the form of resecting is used for a selective group of patients.

V. CONCLUSION

Brain metastasis are common following solid cancers. A look out for cerebral metastasis should be essentially carried out as the impact on survival remains serious. Early diagnosis and aggressive therapy can be beneficial for the patient.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Walker AE, Robbins M, Weinfeld FD. Epidemiology of brain tumors: the national survey of intracranial neoplasms. *Neurology* 1985;35:219–26.
2. Carcangiu ML, Zampi G, Pupia A, Castagnoli A, Rosai J. Papillary carcinoma of thyroid: A clinicopathological study of 241 cases treated at university of Florence, Italy. *Cancer*. 1985;5:805–28.
3. Dorairajan N, Pandiarajan R, Yuvaraja S. A descriptive study of papillary thyroid carcinoma in a teaching hospital in Chennai, India. *Asian J Surg*. 2002;25:300–3.
4. Hoie J, Stenwig AE, Kullmann G, Lindegaard M. Distant metastases in papillary thyroid cancer: A review of 91 patients. *Cancer*. 1988;61:1–6.
5. Patel JK, Didolkar MS, Pickren JW, Moore RH. Metastatic pattern of malignant melanoma. A study

- of 216 autopsy cases. Am J Surg. 1978;135: 807-810.
6. Sampson JH, Carter JH Jr, Friedman AH, Seigler HF. Demographics, prognosis, and therapy in 702 patients with brain metastases from malignant melanoma. J Neurosurg. 1998;88: 11-20.
 7. Floyd CE, Stirling CT, Cohn I Jr. Cancer of the colon, rectum and anus: review of 1,687 cases. Ann Surg 1966;163:829-37.
 8. Patanaphan V, Salazar OM. Colorectal cancer: metastatic patterns and prognosis. Southern Med J 1993;86:38-41.
 9. Stolp, H.B.; Dziegielewska, K.M. Review: Role of Developmental inflammation and blood-brain barrier dysfunction in neurodevelopmental and neurodegenerative diseases. Neuropathol. Appl. Neurobiol. 2009, 35, 132-146.
 10. Brightman, M.W.; Reese, T.S. Junctions between intimately apposed cell membranes in the vertebrate brain. J. Cell Biol. 1969, 40, 648-677.
 11. Oldendorf, W.H.; Cornford, M.E.; Brown, W.J. The large apparent work capability of the blood-brain barrier: A study of the mitochondrial content of capillary endothelial cells in brain and other tissues of the rat. Ann. Neurol. 1977, 1, 409-417.
 12. The Blood-Brain Barrier, Biology and Research Protocols; Nag, S., Ed.; Humana Press: Totowa, NJ, USA, 2003; p. 572.
 13. Ricci PE. Imaging of adult brain tumors. Neuroimaging Clin N Am 1999;9:651-69.
 14. Geijer B, Holtas S. Diffusion-weighted imaging of brain metastases: their potential to be misinterpreted as focal ischaemic lesions. Neuroradiology 2002;44:568-73 Medline.
 15. Hiwatashi A, Kinoshita T, Moritani T, et al. Hypointensity on diffusion-weighted MRI of the brain related to T2 shortening and susceptibility effects. AJR Am J Roentgenol 2003;181:1705-09 Medlin.




Figure 1 and 2 : Showing Intracerebral Metastasis and Supratentorial/Infratentorial metastasis




Figure 3 and 4 : Showing Multiple Intracerebral Metastasis and Supratentorial/Infratentorial metastasis