

Magnitude of Malaria Infection in Ethiopia

Belayneh Regasa¹

¹ Arba Minch University

Received: 6 December 2013 Accepted: 3 January 2014 Published: 15 January 2014

Abstract

Background: Malaria is a major public health problem in worldwide and causes high morbidity and mortality. Studying its prevalence is necessary to implement effective control measures. Therefore, the aim of this study was to determine the prevalence of malaria in Arba Minch hospital, Ethiopia. Method: A cross sectional study was conducted from January to April, 2010. A well designed and structured questionnaire and Laboratory investigation were used to collect data. Data was processed and analyzed with SPSS version 16.0. Results: A total of 400 patients clinically suspected to have malaria were examined with overall prevalence of malaria was 7

Index terms—

1 Introduction

Malaria is a life-threatening infectious disease caused by the protozoan parasite called Plasmodium. The World Health Organization (WHO) estimated 660,000 deaths in 2011 directly attributed to malaria, approximately half of the world's population being at risk of infection [1]. Four main species of malaria infect humans: Plasmodium falciparum (P. falciparum), Plasmodium vivax, Plasmodium malariae and Plasmodium ovale. Plasmodium falciparum is the most highly virulent species and is responsible for almost all of the 1.7-2.5 million deaths worldwide caused by malaria [2,3]. It is a leading public health problem in Ethiopia where an estimated 68% of the population lives in malarious areas and threequarters of the total land mass is regarded as malarious [4] This makes malaria the number one health problem in Ethiopia with an average of 5 million cases per year [5]. The disease causes 70,000 deaths each year and accounts for 17% of outpatient visits to health institutions [6]. The aim of this study is to assess prevalence of malaria in Arba Minch hospital.

Author: Department of Medical Microbiology, Arba Minch University, Arba Minch, Ethiopia. e-mail: belayjanimen@gmail.com II.

2 Methods and Materials

During the period January to April 2010 a total of 400 patients with typical symptoms of the disease, such fever (>37.8 °C), headache and back and joint pain consistent with malaria was included in this study.

3 a) Sample collection

Two milliliters of venous blood samples were collected into an Ethylene diamine tetra acetic acid (EDTA) containing bottles for the study, using vein puncture technique [7].

b) Blood examination III.

4 Result

A total of 400 patients clinically suspected to have malaria in Arba Minch Hospital were participated, of these, 238 (59.5%) were males and 162 (40.5%) were females (Table 1). The overall prevalence of malaria was 7% (28 malaria cases out of 400 patients) of which 18 (64.3%) were positive for Plasmodium falciparum and 7 (25%) for

Plasmodium vivax; the remaining 3 (10.7%) showed mixed infections of Plasmodium falciparum and Plasmodium vivax (Table 2).

Laboratory analysis was carried out after thin and thick blood films prepared according to technique outlined by Cheesbrough [7] and examined microscopically for malaria parasites under the microscope. IV.

5 Discussion

Malaria is a major public health problem in Ethiopia. Over the past years, the disease has been consistently reported as the first leading cause of outpatient visits, hospitalization, and death in health facilities across the country [8]. In this study the overall prevalence rate of malaria was 28 (7%). This result was lower than similar studies done in Ethiopia [9,10,11] but higher than study conducted in other area [12]. This difference might be due to altitude variation and climatological differences that may contribute to a great role for breeding of Anopheles vector. The predominant Plasmodium species detected was Plasmodium falciparum, followed by Plasmodium vivax. This was in agreement with other previous studies [13][14][15][16][17]. But other studies reported that the most prevalent species was Plasmodium vivax, followed by Plasmodium falciparum [18, 19].

Males were more infected than females, which was statistically significant ($p < 0.005$). This is in line with the other previous studies [9,20,21]. The higher prevalence rate might be due to the fact that males engage in activities which make them more prone to infective mosquito bites as compared to females. In all age groups, malaria was reported in the study area. However, significantly affected age groups were 15-19 years old, followed by 20-29 years old. This might be associated with their daily activities. Farming is extensive in Arba Minch. Because of high temperature in this area, daily activities are accomplished especially during night. This may expose them to the bite of mosquitoes.

The occurrence of malaria depends on adequate rainfall and temperature. In areas with a temperate climate, transmission of malaria is commonly limited to months in which the average temperature is above the minimum required for sporogony [22].

V.

6 Conclusion

This study showed that there is high prevalence of malaria. Malaria has statistically significant association with sex and age. Therefore, health planners and administrators need to give intensive health education for the community on prevention and control of malaria.

7 VI.

Volume XIV Issue VII Version I Males 16 (4%) were more infected than females 12 (3%). Gender had statistically significant association with malaria infection ($p < 0.005$) [Table 3].

In this study all age groups were infected but high prevalence observed in age groups 15-19, followed by 20-29 years old (Table 4).

VII.



Figure 1:

1

Variables	Frequency (percent)
participants (n=400)	
Gender	
Male	238 (59.5)
Female	162 (40.5)
Age	
15-19	30 (7.5)
20-29	190(47.5)
30-39	110 (27.5)
40-49	50(12.5)
50-60	20(5)

Figure 2: Table 1 :

2

Malaria parasite	Frequency	Percent (%)
Plasmodium falciparum	18	64.3
Plasmodium vivax	7	25
Mixed infections (Plasmodium falciparum and Plasmodium vivax)	3	10.7

Figure 3: Table 2 :

3

Sex	Number of examined	Number infected	Percentage	P-value
Male	238	16	4	0.001
Female	162	12	3	
Total	400	28	7	

Figure 4: Table 3 :

4

Age	Number of examined	Number infected	Percent (%) infected
15-19	30	12	3
20-29	190	8	2
30-39	110	4	1
40-49	50	3	0.75
50-60	20	1	0.25
Total	400	28	7

Figure 5: Table 4 :

.1 Acknowledgements

The author would like to thank those who were involved in this research.

.2 Competing Interest

The author declared that there is no any relevant competing interest to disclose in this research.

[Starzengruber et al.] , P Starzengruber , H Fuehrer , B Ley , K .

[Mandel et al. ()] , B K Mandel , E G L Wilkins , E M Dunbar , R M White . Lecture Notes on the Infective Disease 1984. Blackwell Scientific Publications. (4th edition)

[World Health Organization, World Malaria Report ()] , *World Health Organization, World Malaria Report* 2005. WHO

[Asnakew et al. ()] , K Y Asnakew , G Sucharita , T H Afework , O D Dereje , P P Hrishikesh . *Spatial analysis of Journal of Health Geographics* 2009. 8 p. .

[World Health Organization: World Malaria Report ()] , <http://www.webcitation.org/query.php?url=http://www.who.int/malaria/2875-13-16>. http://www.who.int/malaria/publications/world_malaria_report_2012/wmr2012_no_profiles.pdfwebcite *World Health Organization: World Malaria Report* 2012.

[Woyessa et al. ()] ‘An indigenous malaria transmission in the outskirts of Addis Ababa, Akaki Town and its environs’. A Woyessa , T Gebre-Michael , A Ali . *Ethiopian Journal of Health Development* 2004. 18 p. .

[Ramos et al. ()] ‘Change in epidemiology of malaria infections in a rural area in Ethiopia’. J M Ramos , F Reyes , A Tesfamariam . *Journal of Travel Medicine* 2005. 12 (3) p. .

[Gabriel and James ()] ‘Developing malaria early warning system for Ethiopia’. S Gabriel , V James . *25th Annual ESRI International User Conference*, (San Diego, Calif, USA) 2005. National Center for EROS (Paper no. UC2409)

[Cheesbrough ()] *District Laboratory Practice in Tropical Countries. Part 2. 2 nd ed. Cambridge University, M* Cheesbrough . 2006. p. .

[Thriemer et al. ()] ‘High prevalence of asymptomatic malaria in southeastern Bangladesh’. P Thriemer , V Swoboda , Elisabeth . *Malaria Journal* 2014. 13 p. 16.

[Ghebreyesus et al. ()] ‘Household risk factors for malaria among children in the Ethiopian highlands’. T A Ghebreyesus , M Haile , K H Witten . *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2000. 94 (1) p. .

[Aikawa ()] ‘Human cerebral malaria’. M Aikawa . *The American Journal of Tropical Medicine and Hygiene* 1988. 39 (1) p. .

[Solomon et al. ()] ‘Malaria pattern observed in the highland fringe of Butajira, Southern Ethiopia: a ten-year retrospective analysis from parasitological and metrological data’. T Solomon , B Yeshambel , T Takele , M Girmay , M Tesfaye , P Beyene . *Malaria World Journal* 2012. 3 (5) .

[Fmoh ()] ‘National Five Year Strategic Plan for Malaria Prevention and Control in Ethiopia’. Fmoh . *Ministry of Health*, (Addis Ababa, Ethiopia) 2006-2010. 2006.

[President’s Malaria Initiative. Malaria Operational Plan (MOP) Ethiopia. FY ()] *President’s Malaria Initiative. Malaria Operational Plan (MOP) Ethiopia. FY*, 2008.

[Karunamoorthi and Bekele ()] ‘Prevalence of malaria from peripheral blood smears examination: a 1-year retrospective study from the Serbo Health Center’. K Karunamoorthi , M Bekele . *Journal of Infection and Public Health* 2009. 2 (4) p. .

[Karunamoorthi and Bekele ()] ‘Prevalence of malaria from peripheral blood smears examination: a 1-year retrospective study from the Serbo Health Center’. K Karunamoorthi , M Bekele . *Journal of Infection and Public Health* 2009. 2 (4) p. .

[Woyesa et al. ()] ‘Prevalence of malaria infection in Butajira area’. A Woyesa , W Deressa , A Ali , B Lindtjorn . *South central Ethiopia. Malaria journal* 2012. 11 p. 84.

[Abebe et al. ()] ‘Ten year trend analysis of malaria prevalence in Kola Diba’. A Abebe , M Dagnachew , M Mikrie , A Meaza , G Melkamu . *Parasites and Vectors*, (North Gondar, Northwest Ethiopia) 2012. 5 p. 173.

[Molineaux ()] ‘The epidemiology of malaria as an explanation of its distribution, including some implications for its control’. L Molineaux . *Malaria Principles and Practice of Malariology*, W Wernsdorfer (ed.) (Churchill Livingstone, Great Britain, UK) 1988. p. .

[Bray and Sinden ()] ‘The sequestration of Plasmodium falciparum infected erythro-cytes in the placenta’. R S Bray , R E Sinden . *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1979. 73 (6) p. .

[Sintasath ()] ‘The state of Ministry of Health of Eritrea’. D Sintasath . *Activity Report 2000-2001*. 2004. 134. (National malaria survey)

[Deressa et al. ()] ‘Treatment seeking of malaria patients in east Shewa of Oromia’. W Deressa , D Olana , S Chibsa . *Ethiopian Journal of Health Development* 2003. 17 p. .