

Oxidative Stress Induced Carbonyl Group Incorporation to Human RBC Membrane: Role in Vivo Senescence of Erythrocyte

Arghya Sur¹, Hemontika Chakraborty² and Arindam Basu³

¹ Oxidative Stress Induced Carbonyl Group Incorporation to Human RBC Membrane: Role in vivo Senescence of Erythrocyte.

Received: 15 December 2013 Accepted: 31 December 2013 Published: 15 January 2014

Abstract

This study was focused to identify the link between oxidative stress and senescence of erythrocytes in vivo. To elucidate this mechanism, various modification of RBC membrane proteins and lipids were analyzed invitro, exposing them to oxidative stress and the results were compared with the changes observed in erythrocytes undergoing senescence in vivo. The other objective was to confirm the mechanism of autoantibody mediated removal of aged RBCs. Our results established that increased lipid peroxidation products, followed by the enhanced damage of RBC membrane protein caused increased RBC membrane protein carbonylation, to normal red cells exposed to the in vitro Fe²⁺ ascorbate induced oxidative stress. It was presumable that these changes were mediated by hydroxyl (·OH) radicals. Further, similar changes were also seen in percoll gradient age fractionated high density aged RBCs.

Index terms— human RBC membrane, protein carbonylation, oxidative membrane damage, senescence of RBC, oxygen free radical, ROS, autoantibody to RBC.

1 Introduction

The term 'senescence' in the context of normal blood cells implies that the cells are removed from the circulation in an age dependent manner. Life span of erythrocyte varies between species, is exceptionally constant within a species [1]. Over the last few decades several age related alterations of erythrocyte membrane and mechanisms of damaged or aged RBCs have been investigated [1, 2]. Out of these oxidative damage to erythrocyte membrane proteins and lipids is presently thought to play key role during senescence of normal RBCs as well as accelerated senescence of pathological red cells seen in thalassemia, sickle cell anaemia etc. The oxidative damage is initiated probably by oxygen free radicals (ROS) and other oxidants produced endogenously [3, 4, 5, 6]. In vitro characterization of density separated red cells also has provided cumulative oxidative damage [7].

Dense human red cells show moderately increased amounts of methaemoglobin [8]. Exposing intact erythrocytes or ghosts to oxidizing systems in vitro showed various modifications e.g. lipid peroxidation, cross-linking and fragmentation of membrane cytoskeletal proteins, binding of hemichrome and heinz body to the inner surface of the membrane, clustering of band-3 protein, loss of membrane free sulphhydryl groups, incorporation of carbonyl content in the membrane protein etc [9, 10, 11, 12, 13, 14, 15, 16, 17]. Some of these changes have also been studied in vitro density separated or biotinylated aged erythrocytes and also in pathological red cells of accelerated ageing [1, 2, 18, 19, 20, 21]. Many of these changes are seen in stored RBC in blood bank [22, 23]. A possible reason for cumulative oxidative damage to the red cell membrane would be a reduction in the activity of oxidative defense enzymes like, superoxide dismutase (SOD), glutathione peroxidase, glutathione reductase etc [24, 25]. Superoxide radicals (·O₂⁻) undergo dismutation reaction spontaneously (2 ·O₂⁻ + 2H⁺ + 2H₂O₂) or catalyzed by SOD, produces H₂O₂ (2 ·O₂⁻ + 2H⁺ + SOD → H₂O₂) [26]. Again, ·O₂⁻ and H₂O₂ in presence of Fe

44 $2+ / \text{Cu} + / \text{Mn} 2+$ can produce more reactive $\cdot\text{H}$ radical by Haber Weiss reaction ($\cdot\text{O}_2 + \text{H}_2\text{O}_2 \xrightarrow{\text{Fe}^{2+} / \text{Cu}^{+} / \text{Mn}^{2+}} \cdot\text{H} + \text{OH}^-$). In biological system, simple salt of $\text{Fe}^{2+} / \text{Cu}^{+} / \text{Mn}^{2+}$ and H_2O_2 can also produce
45 $\cdot\text{H}$ radical via Fenton reaction ($\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \cdot\text{H} + \text{OH}^-$) [27] Fe^{3+} to Fe^{2+} may promote
46 metal catalyzed Haber Weiss reaction, to produce more damaging $\cdot\text{H}$ radicals [28,29]. Hence, Ascorbate has
47 dual role -anti oxidant and pro-oxidant, where latter is exhibited in presence of transition metals [27,30,31,32,33].
48 Again, it has been shown that membrane of high density red cell is highly susceptible to peroxidative damage
49 exclusively to poly unsaturated fatty acids of membrane in a chain reaction [27, 34,35,36,15]. The signals
50 that trigger the removal of aged erythrocytes and the actual mechanisms of such removal, i.e. elucidation of link
51 between oxidative membrane damage and removal of aged cells from circulation are still controversial. However
52 it has been suggested that recognition and removal of aged, infected or damaged RBC involve multiple pathways;
53 mainly antibody independent phagocytosis, antibody dependent phagocytosis and removal by splenic sinuses
54 [37]. In antibody independent removal, oxidatively damaged erythrocytes are phagocytosed by macrophages
55 through scavenger receptors for low density lipo protein (LDL) in absence of opsonizing antibodies [38,39]. In
56 antibody dependent removal of aged RBC, auto antibodies (anti band-3, antigalactosyl antibodies or antibodies
57 to malonaldehyde-protein adduct) bind to the surface of aged RBCs leading to phagocytic removal of aged,
58 oxidatively damaged and pathologically damaged red cells by macrophages [40,41,42, 43, 43,37,23]. But the
59 nature of such antibodies has been subjected to debate [40,41,42,37,18, 44].

61 It has been proposed that impaired deformability provides a major route of the destruction of senescent red
62 cells [44,45]. Poor deformability or increased rigidity due to oxidatively modified red cells as well as aged RBC
63 membrane proteins and lipids retard the red cells' movement in the spleen and they may be entrapped in splenic
64 sinuses for destruction [14,18,46,47].

65 It was clear that although oxidative damage to RBC membrane protein and lipid play key role in the
66 mechanism of senescence of erythrocyte; but there were important lacunae in our present state of knowledge.
67 This investigation led to fill up that lacunae and which in the long run might lead to identification of suitable
68 agents to prolong the survival of endogenous or transfused RBCs in the circulation.

69 2 II.

70 3 Aims and Objectives

71 The specific objectives of this work were-a) study of various oxidative modifications of human erythrocyte
72 membrane lipids and proteins exposed in vitro to physiologically relevant oxidizing system like iron and ascorbate.
73 b) study of effects of free radicals and metal chelators on the above system. c) to investigate if oxidative damage to
74 RBC membrane like increased carbonylation led to enhanced binding of immunoglobulins (IgG) to cell surface. d)
75 to study whether the oxidative damage to RBC membrane observed in vitro were also present in aged erythrocytes
76 obtained from the circulation by density gradient method.

77 III.

78 4 Materials and Methods

79 Blood obtained from healthy human male and female volunteers of age group 20 to 50 years was collected in 3.2%
80 sodium citrate solution. Institute's research ethical committee approval was obtained. All the healthy blood
81 donors were informed the purpose of the study prior to drawing of blood.

82 5 Materials

83 Percoll, Alpha cellulose, Microcrystalline cellulose, Catalase, Superoxide dismutase (SOD), Diethylene triamine
84 penta acetic acid (DETAPAC), Butylated hydroxy toluene (BHT), Bovine serum albumin (BSA), Polyvinyl
85 difluoride (PVDF) membranes, Alkaline phosphatase conjugated goat anti rabbit IgG, Polyclonal anti-DNP
86 antibody, 4-(2-hydroxy ethyl)-1-piperazine ethane sulphonic acid (HEPES), Phenyl methyl sulphonyl fluoride
87 (PMSF), Tween20(Polyoxyethylene sorbitan mono laurate, Sigma ultra) and Ferritin (Type1: from horse spleen)
88 were purchased from Sigma Chemical Co., USA., Biogel P-6 were obtained from Bio Rad, USA., 5-bromo-4
89 chloro-3-indolyl-phosphate/ nitro blue tetrazolium (BCIP/NBT), Tetra methyl benzidine (TMB), Protein-A-
90 horse radish peroxidase conjugate, protein -A-agarose column and Human IgG purification kit were obtained from
91 Bangalore Geni Pvt. Ltd., India. Thiobarbituric acid (TBA), Trichloro acetic acid (TCA) were from E.Merck,
92 Germany. 2-Deoxy ribose (DR), Mannitol, Dimethyl sulphoxide (DMSO), L-Ascorbic acid, 2,4-dinitrophenyl
93 hydrazine (DNPH) and all chemicals for Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-
94 PAGE) were purchased from Sisco Research Laboratory, Mumbai, India. All other chemicals used were of
95 analytical grade.

96 6 Methods a) Preparation of human erythrocyte ghosts

97 Human blood collected in 3.2% sodium citrate solution made free from WBC and Platelets by passing through
98 a column bed of microcrystalline cellulose and alpha cellulose (1:3) as adopted from Beutler et al (1975) [48].
99 . Briefly, microcrystalline cellulose (mean size 50 μm , Sigma cell-50) and α -cellulose were taken in 1:3 ratio by
100 weight and suspended in phosphate buffered saline (PBS), pH 7.4 and packed in a 10 ml plastic syringe with an

101 inner diameter of 1.5 cm, so that 5ml column was 1.5 cm by 5 cm. The packed column was washed with PBS, pH
102 7.4. The blood (approx 5 ml) was centrifuged at 1,000 x g for 5 min and the supernatant plasma was removed
103 along with the buffy coat. The packed red blood cells were poured over the cellulose column and washed down
104 with ice cold isotonic PBS, pH 7.4. The eluate was collected (approx. 40-50 ml for 5 ml of blood) and centrifuged
105 at 1,000 x g for 10 min to obtain the erythrocyte pellet. The erythrocytes were routinely recovered from the
106 column up to 80% and the preparations were practically free of leucocytes (> 99.5%) and platelets (> 95%) as
107 verified microscopically. The erythrocyte ghosts were prepared by hypotonic lysis [49]. The pelleted erythrocytes
108 were lysed in 60 volumes of 5 mM sodium phosphate buffer, pH 8.0 followed by centrifugation at 20,000 x g to
109 obtain haemoglobin free white ghosts. The creamy white ghosts were suspended in 50 mM phosphate buffer, pH
110 7.4 and kept frozen at -20°C for subsequent experiments.

111 **7 b) Estimation of protein**

112 Protein content of erythrocyte membrane ghosts was estimated by the method of Lowry et al (1951) [50] after
113 solubilizing in 1% SDS and using BSA as standard. c) 2-Deoxy ribose (DR) degradation assay Hydroxyl radical
114 generation in the incubated mixture of ascorbate and/or Fe 2+ has been measured by DR degradation assay [51].
115 The reaction mixture in 50 mM phosphate buffer, pH 7.4 contained 1mM DR, 0.5 mM ascorbate, 0.2 mM FeSO
116 4 or 5.6 ?M Ferritin with or without other additions like mannitol (20mM) or 20 mM DMSO in a total volume of
117 600 ?l. Incubation was terminated after 1 h by addition of 1.4ml of 2.8% TCA to each tube followed by addition
118 of 0.6ml of 1% (w/v) TBA. Then all the tubes were heated for 10 min in a boiling water bath. The tubes were
119 cooled briefly and absorbance taken at 532 nm.

120 **8 d) SDS-PAGE of erythrocyte membrane ghosts**

121 Freshly prepared erythrocyte ghosts with Fe 2+ (0.2mM) and ascorbate (0.5mM) were incubated at 37°C for
122 2 h with or without other additions. Incubation was terminated by addition of electrophoresis sample buffer
123 containing 3% SDS, and 5% ?-mercapto ethanol followed by heating in boiling water bath for 3 min. The
124 samples were immediately applied for discontinuous SDS-PAGE in reducing condition using 10% separating gel
125 following the method of Laemmli (1970) [52]. Slab gels were stained with coomassie brilliant blue-R and destained
126 in 5% methanol & 7.5% glacial acetic acid mixture.

127 **9 e) Lipid peroxidation study**

128 Lipid peroxidation in incubated ghosts was estimated by measuring the production of malonaldehyde (MDA) as
129 described by Ohkawa et al (1979) [53,51]. Erythrocyte ghosts incubated with or without ascorbate (0.5mM)
130 and /Fe 2+ (0.2mM) at 37°C up to 2 h with other additions like 0.5 mM BHT, mannitol (20mM) , DMSO
131 (20mM) or catalase (50 ?g/ml) in a total volume of 200 ?l. To each 200 ?l of incubation mixture were added 100
132 ?l of 8.0% SDS, 750 ?l of 20% acetic acid (pH 3.5) and 750 ?l of 8.0% aqueous TBA. The samples were heated at
133 100°C for 15 min. The tubes were briefly cooled and 2.5 ml of a mixture of nbutanol and pyridine (15:1, v/v)
134 was added to each tube and vortexed thoroughly. The samples were centrifuged at 4,000 r.p.m. for 10 min. The
135 organic layer was collected and absorbance measured at 532 nm. Amount of MDA was expressed in n moles/mg
136 protein using a molar extinction co-efficient of $1.56 \times 10^5 \text{ M}^{-1} \cdot \text{cm}^{-1}$ [54].

137 **10 f) Estimation of carbonyl content**

138 Erythrocyte ghosts were incubated with or without ascorbate (0.5mM) and Fe 2+ (0.2mM) at 37°C up to 2 h
139 with other additions in a total volume of 200 ?l.

140 To each tube was added 500 ?l of TCA followed by centrifugation at 3,000 r.p.m. to obtain a precipitate.
141 To each precipitate 500 ?l of 10 mM DNPH in 2 M HCl was added. The tubes were allowed to stand at room
142 temperature for 10 min with occasional vortexing. Then to each sample 500 ?l of 10% TCA was added followed by
143 vortexing and centrifugation at 10,000 x g for 5 min. The supernatant was discarded and the pellet was washed
144 3 times with 1ml ethanol-ethyl acetate mixture (1:1) to remove excess reagent. Precipitated protein as pellet
145 was solubilized in 1ml protein dissolving solution containing 2% SDS, 0.05% EDTA in 0.8 M sodium phosphate
146 buffer, pH 8.0. Appropriate sample and reagent blanks were kept for this assay. Absorbance was taken at 370
147 nm and the carbonyl content of each sample was calculated using a molar absorption coefficient of $22,000 \text{ M}^{-1} \cdot \text{cm}^{-1}$
148 and the results were expressed in terms of n moles/mg protein [55,56,57,58].

149 **11 g) Estimation of sulphydryl group**

150 Erythrocyte ghosts, in presence or absence of ascorbate (0.5 mM) and Fe 2+ (0.2mM) with other additions
151 were incubated for 2 h at 37°C in 50 mM phosphate buffer, pH 7.4 in a total volume of 200 ?l. At the end
152 of the incubation, erythrocyte membrane proteins were solubilized by the addition of 1 ml of protein dissolving
153 solution containing 2% SDS, 0.05% EDTA in 0.8 M sodium phosphate buffer, pH 8.0. To each sample 50 ?l
154 of DTNB solution (40mg DTNB in 10 ml of 0.1 M sodium phosphate buffer, pH 8.0) was added. A set of
155 reagent blanks were made and absorbance taken at 412 nm after 15 min, The thiol content was estimated using

14 K) SEPARATION OF YOUNG AND AGED ERYTHROCYTES

156 a molar absorption coefficient of thiols/mg of protein [13,57]. containing IgG was passed through a Bio-gel P-6 column
157 equilibrated in PBS, pH 7.4. IgG was stored in aliquots in presence of 0.1% sodium azide at 4°C.

158 12 i) Binding of autologous IgG to oxidatively damaged red 159 blood cells

160 Erythrocyte ghosts were incubated with or without Fe²⁺ (0.2 mM) /ascorbate (0.5mM) in presence or absence
161 of mannitol (20 mM), DMSO (20 mM) or catalase (50 ?g/ml). After the incubation at 37°C for 2h the red cell
162 ghosts were pelleted down. The supernatant from each tube was aspirated and the pellet suspended in PBS,
163 pH 7.4 containing 1% BSA. Samples (100 ?l) of red cell ghost suspension (control, oxidized and oxidized with
164 inhibitors) were taken in micro centrifuge tubes pre-coated with 1% BSA. To each tube was added 100 ?l of IgG in
165 PBS -1% BSA and the tubes were left overnight at 4°C for IgG binding. Appropriate blanks (without membrane
166 suspension) and negative controls (without added IgG) were kept. After the incubation, to each tube was added
167 0.8 ml of PBS-0.2% BSA followed by centrifugation at 10,000 x g for 10 min. The supernatant was removed and
168 the pellet was washed twice with 1 ml of PBS-0.2% BSA. 100 ?l of protein-A-HRP conjugate (diluted 1:10,000)
169 was added to each tube and kept at 4°C for 1 h followed by centrifugation at 10,000 x g for 10 min after addition
170 of 0.8 ml of PBS-0.2% BSA. The pellet was washed twice with PBS-0.2% BSA to remove completely the excess
171 unbound Protein-A-HRP. Subsequently, 100 ?l of the substrate solution TMB/H₂O₂ was added to each tube
172 and mixed thoroughly. The tubes were kept in dark for 30 min during which a blue colour was developed. The
173 reaction was stopped by addition of 200 ?l of 2 M H₂SO₄ which also changed the colour of the reaction mixture
174 from blue to yellow. The absorbance was taken at 450 nm. The absorbance of the blank and the appropriate
175 negative control were subtracted from the absorbance of each test sample. The net absorbance of each test
176 sample was a measure of IgG bound to red cell ghosts. IgG bound to oxidized ghosts with or without inhibitors
177 was expressed as ratio to IgG bound to control unoxidized ghosts after appropriate corrections for differences in
178 protein content in individual samples [13,43].

179 13 j) Purification of band-3 protein from human erythrocyte

180 Human erythrocyte ghosts obtained were suspended in a solution containing 1 mM EDTA, 5 mM ?-mercapto
181 ethanol, 0.03 mM PMSF, which was adjusted to pH 7.5 with 1 M NaOH and stirred gently for 18 h at 4°C. The
182 membranes were recovered by centrifugation at 1,00,000 x g for 1 h and re suspended in 50 mM sodium phosphate
183 buffer, pH 7.4. The membrane pellet was subjected to SDS-PAGE as described earlier. The protein visible in
184 the gel was predominantly band-3 and this alkali treated pellet was used as partially purified band-3 protein.
185 Autologous IgG (1,5 mg/ml) was mixed with an equal volume of suspension of partially purified band-3 protein
186 (alkali treated erythrocyte pellet, protein content 2 mg/ml) and kept at 4°C for 2 h followed by centrifugation at
187 1,00,000 x g for ½ h. The supernatant recovered was used as autologous IgG, depleted of anti band -3 antibodies
188 (IgG -depleted) and used for IgG binding assay [43].

189 14 k) Separation of young and aged erythrocytes

190 Young and aged erythrocyte of human were isolated by isopycnic centrifugation through gradient of "Percoll",
191 a commercially available polyvinyl-pyrrolidonecoated colloidal silica following the method of Rennie et al (1979)
192 [59]. A continuous density gradient of "Percoll" (0-100%) was prepared with the help of a gradient maker and
193 a peristaltic pump and using two stock solutions: one containing 5.263% BSA in water and another containing
194 5.263% BSA (w/v) in percoll. 19 volumes of each of these solutions were separately mixed with 1 volume of
195 solution (final pH 7.4) containing 2.66 M NaCl, 0.09 M KCl and 200 mM HEPES, pH 8.5. Thus, resultant
196 mixtures were isotonic containing 5% BSA. 0.8 ml of red cell suspension was brought to room temperature and
197 carefully layered on the top of the continuous percoll gradient (8 ml). Age fractionation of red cell was achieved
198 by centrifugation at 1,100 x g for 9 min at room temperature followed by slow deceleration of the rotor. The cell
199 fractions were then sequentially aspirated from the top of the gradient using a Pasteur pipette attached to the
200 peristaltic pump. Top 20% of the fractionated cells were taken as young cells and bottom 20% of the cells used as
201 aged red cells. The cells were finally washed to free percoll by 3 washes in PBS, pH 7.4 [59]. 1) After preparation
202 of young and aged erythrocyte membrane, protein carbonyl content, free thiol groups, protein cross-linking etc
203 were measured by previously mentioned methods.

204 m) Immunoblot analysis for protein carbonyls in young and aged erythrocyte membrane ghosts

205 Freshly prepared young and aged erythrocyte membrane ghosts were taken in two separate tubes each
206 containing 100 ?l of RBC membrane ghosts and to each of them was added 50 ?l of 18% SDS to solubilized
207 the protein followed by addition of 200 ?l 20 mM DNPH in 10% TFA. A yellow colour was developed after
208 incubating each mixture for 10 min. The samples were then neutralized to orange red colour by slow addition
209 of 2 M Tris in 30% glycerol and 19% ?-mercapto ethanol. The samples were subjected to SDS-PAGE followed
210 by electroblotting in a semi dry electroblotter unit, using a constant current of 2 m.A./cm² for 45 min. The
211 transfer of proteins to PVDF membrane was confirmed by staining one portion of the blotted Volume XIV Issue
212 VII Version I (D D D D)

213 Year 2014 C containing 1% BSA and kept for 2 h for blocking the nonspecific sites. The blocked membrane was
214 incubated at 37°C with shaking in presence of anti-DNP antibody (diluted 1:1000) for 1 h. After the incubation,
215 the PVDF membrane was washed in a washing solution containing 0.1% (v/v) Tween-20 in PBS followed by
216 further incubation for 1 h with alkaline phosphatase conjugated goat anti rabbit IgG. The bands in the PVDF
217 membrane were detected by the addition of BCIP/NBT [55,32].

218 15 n) Quantitation of cell bound IgG of young and aged 219 erythrocytes

220 The young and aged red cells were washed thoroughly in PBS followed by preparation of corresponding membrane
221 ghosts as described earlier.

222 Membranes of young and aged erythrocytes were taken in separate microcentrifuge tubes pre-coated with 1%
223 BSA. To each tube was added autologous IgG in PBS-1% BSA and the tubes were left overnight at 4°C for
224 IgG binding. Bound IgG was quantitated as mentioned earlier. IgG bound to aged RBC membrane ghosts was
225 expressed as a ratio of IgG bound to control young RBC membrane ghosts [43].

226 16 o) Statistical Analysis

227 All experiments were checked for reproducibility and statistical significance worked out. All the values in Tables
228 and Charts of result section were presented as mean \pm standard error of mean (SEM). Significance of difference
229 between groups were determined following Student's 't' test for paired and unpaired observations [60].

230 IV. ascorbate (0.5 mM) for 2 h at 37°C. d) A significant increase in protein carbonyl content in red cell
231 membrane ghosts exposed to iron and ascorbate for a period of 2 h was observed (Fig- ??).

232 17 Results

233 18 a) Results presented in

234 The phenomenon was inhibited by hydroxyl radical scavengers like mannitol, DMSO and also by antioxidant
235 enzyme, catalase (Fig- ??). The inhibition was statistically significant. e) When erythrocyte ghosts (protein
236 content 0.6 to 1.0 mg/ml) were incubated with Fe 2+ (0.2 mM) and ascorbate (0.5 mM) at 37°C, MDA produced
237 per mg protein per 2 h were significantly higher with respect to control erythrocyte ghosts. Lipid peroxidation
238 was significantly inhibited (about 90%) by 0.5 mM of BHT, while mannitol (20 mM), DMSO (20 mM) and
239 catalase (50 ?g/ml) did not inhibit MDA production (Table-3). f) g) In order to elucidate the link between
240 oxidative stress and enhanced IgG binding to red cell ghosts several radical scavengers and antioxidant enzymes
241 were used in the binding assay. Table-5 showed that catalase and hydroxyl radical scavengers like mannitol (20
242 mM) and DMSO (20 mM) prevented significantly enhanced binding of autologous IgG to oxidized erythrocyte
243 ghosts. h) The parameters of oxidative stress were then measured in density separated erythrocytes in a percoll
244 gradient. There was a statistically significant rise (1.5 fold) in protein carbonyl content in aged RBC membrane
245 ghosts with respect to young red cell membrane ghosts (Fig- ??).

246 (1971) [49]. Incubation of erythrocyte membranes with Fe 2+ (0.2 mM) and ascorbate (0.5 mM) did not
247 however produce any cross-linking or protein fragmentation (Lane-b, c & d).

248 i) Immunoblot analysis of young and aged erythrocyte ghosts using anti-DNP antibody also showed increased
249 carbonylation of proteins in aged erythrocyte membrane (Fig- ??).

250 blotted membrane was incubated with PBS enzyme catalase (50 ?g/ml) upto 58.9%. Again, it

251 Table-4 showed that a very significant increase (about 4 fold) in the binding of auto logous IgG occurred
252 in oxidized ghosts compared to control. The binding of auto logous IgG to ghosts was quantitated by an
253 immunoassay using Protein-A-HRP conjugate and TMB (substrate). Since it was difficult to estimate the
254 number of IgG molecules bound per ghost cell, the net absorbance at 450 nm for each sample normalized to
255 a protein concentration of 1 mg/ml was calculated and taken as a measure of bound IgG. The net absorbance for
256 each sample was derived by subtracting from the total absorbance, the values for the blank and the appropriate
257 negative control (ghost incubated without IgG). This increased binding of oxidized ghosts was however not seen
258 when anti-band-3 depleted IgG was used for binding assay.

259 membrane after it was dried. The other half of the j) Fig- ?? showed that there was no significant decrease in
260 protein thiol content in aged RBC membrane ghosts compared to young RBC membrane ghosts. k) SDS-PAGE
261 of young and aged erythrocyte ghosts showed that there was no such remarkable difference in the band pattern
262 of membrane proteins (Fig- ??), and especially no cross-linked high molecular weight proteins or evidence of
263 protein fragmentation was apparent in aged erythrocyte. l) As presented in Table-6, autologous IgG binding to
264 aged erythrocyte ghosts was more than 2.5 fold higher than that of young erythrocyte ghosts, when incubation
265 was carried out with IgG concentration of 1 mg/ml. Raising IgG concentration to 2.5 mg/ml, IgG binding to
266 aged red cell ghosts was about 3.7 times more than that of young erythrocyte ghosts.

267 V.

268 19 Discussion

269 Erythrocyte membrane encounters oxidative stress both from cell interior and exterior [3]. Different investigators
270 have studied red cell membrane alterations by subjecting intact RBC or RBC membrane ghosts to different
271 oxidizing systems like ascorbic acid and iron /copper, H_2O_2 /iron, ADP/Fe³⁺, phenyl hydrazine, xanthine /
272 xanthine oxidase etc have been reported [47,31, 2004,43, 2001, 2002]. Many of these changes have also been noticed
273 in aged red cells [1, 2, 3, 14, 15] or in damaged RBCs undergoing accelerated senescence in some pathological
274 conditions [19, 2002, 2002,21,15,16] or in stored RBC in blood bank [22,23].

275 In this study, mixture of ascorbic acid and iron (Fe²⁺) has been used to induce oxidative stress, because both
276 these components are physiologically relevant in the context of oxidative injury to erythrocytes in vivo. Several
277 discrete iron compartments in red cell have been suggested [21]. This iron is bioactive and can valance-cycle
278 between Fe³⁺ and Fe²⁺ states is capable of generating hydroxyl radicals [31,33,27,29]. Reducing agents like
279 ascorbate can potentiate this mechanism and thereby promote the formation of hydroxyl radicals via Fenton's
280 mechanism [30,29]. The elaborate transport and recycling of ascorbate by red blood cells in vivo have been
281 highlighted by many investigators [28,33].

282 Results from Table-1 indicate that a mixture of ascorbate and iron (Fe²⁺) is an active source of hydroxyl
283 radicals generated by Fenton's reaction as evident from the inhibition of 2-deoxyribose degradation by catalase
284 and hydroxyl radical scavengers like mannitol and DMSO. The oxidation of Fe²⁺ to Fe³⁺ produces $\cdot OH$ radicals,
285 which by dismutation produces H_2O_2 . H_2O_2 produce H_2O_2 [33]. This autoxidation of ascorbate is
286 $\cdot OH$ mediated and metal dependent as evident from its inhibition by SOD and DETAPAC [33]. The fact that
287 ascorbate can interact with ferritin to generate $\cdot OH$ radicals as shown in Table-1 implies that a similar reaction
288 between ascorbate and native or denatured haemoprotein in red cells may also lead to the formation of toxic
289 reactive oxygen species(ROS).

290 ROS can induce various forms of protein damage such as cross-linking of polypeptide chains, oxidation of
291 protein thiol (-SH) groups, incorporation of carbonyl groups into protein etc had been reported [57,43,55,22].
292 In hydroxyl radical generating system like iron/ascorbate, we have examined some of these modifications which
293 are considered as hallmark for oxidative modification [55,57,32]. Under our experimental conditions protein
294 cross-linking or fragmentation were not observed (Fig- 1). Some earlier studies however, showed cross-linking of
295 isolated cytoskeletal proteins of erythrocyte membrane when incubated for prolonged time with haemoglobin /
296 H_2O_2 [46] or with xanthine/ xanthine oxidase [11] etc.

297 Oxygen free radicals also cause aggregation of proteins, converting the side chains of cysteine, methionine,
298 histidine and tyrosine and form disulphide bonds as the consequence of oxidation of free thiol groups [13,22]. The
299 present results showed no significant decrease in protein thiol content, which was in conformity with the result
300 of Beppu et al (1990) [43]. Beppu et al (1989) also observed from their result of SDS-PAGE of RBC membrane
301 proteins in reducing condition no significant cross-linking of RBC membrane protein exposed to oxidative stress
302 with respect to control ghosts [13] which was in conformity of our result (Fig- 1). However, it has been observed
303 by the same observers that SDS-PAGE of RBC membrane proteins in non-reducing condition formed disulphide
304 bond and cross-linking of protein during oxidative stress [13,43].

305 The metal catalyzed oxidation of proteins can introduce carbonyl group at lysine, arginine, proline and
306 threonine residues in a 'site-specific' manner [61]. Free radical damage to amino acid residues and/or reaction with
307 aldehydes or both derived from lipid peroxides could contribute towards generation of protein carbonyl [61]. The
308 carbonyl bearing residues have not been completely identified but gamma-glutamyl semialdehyde appeared to be
309 the major residue [55]. In different purified and crude extracts of protein several workers have reported changes
310 in carbonyl content as a consequence of oxidative modification [55, 62.57], but on erythrocyte membranes
311 such reports are scanty. The results obtained from this study (Fig- 2) further reacts with Fe²⁺ to give rise to
312 $\cdot OH$ radicals by Fenton's mechanism [31,26,33]. Ascorbate potentiates the reaction by reductive cycling of iron
313 from Fe³⁺ to Fe²⁺ [29]. Further ascorbate undergoes autoxidation to ascorbate induced oxidative stress to
314 RBC membranes caused a significant rise in red cell membrane protein carbonyl content which was inhibited by
315 catalase and hydroxyl radical scavengers-mannitol, and DMSO (Fig- 2) which indicated that the increase in
316 protein carbonyl content was mediated by hydroxyl radical attack.

317 In crude tissue preparation increased carbonyl content as a result of oxidant stress may also be secondary to
318 associated lipid peroxidation [57]. During lipid peroxidation, lipid peroxide or hydroperoxide or peroxy radicals
319 are formed [27]. These preformed lipid peroxidation products which broke down to yet more radical species by
320 reacting with transition metal like iron [27] in turn caused oxidative damage to membrane proteins [57]. This type
321 of damage could not be inhibited by $\cdot OH$ radical scavengers like mannitol, DMSO or anti oxidant enzyme catalase;
322 but lipid soluble chain breaking antioxidants like α -tocopherol and BHT were effective [57]. In this study however,
323 protein carbonylation has been inhibited significantly by hydroxyl radical scavengers like mannitol, DMSO and
324 catalase (Fig- 2); but lipid peroxidation of RBC membrane was not inhibitable by these agents (Table -3). This
325 implied that in this system protein oxidation was not secondary to lipid peroxidative damage. The enhanced
326 production of MDA in red cell membranes by Fe²⁺ and ascorbate during in vitro incubation was presumably the
327 result of breakdown of pre-formed lipid peroxides or hydro peroxides by Fe²⁺ (Table -3). Ascorbate potentiated
328 such iron catalyzed break down of peroxides and hydro peroxides keeping the iron in the reduced (Fe²⁺)
329 state [57]. That has been observed in other systems also [27, 2002, 2002, 2003]. However, the consequence of such
330 peroxidative damage to membrane was controversial [28,34].

331 Autologous IgG binding to red cell surface has been studied by many investigators [40,18,42], but there were
332 several controversies regarding the binding site and antigenic specificity of bound autologous antibody [37,18,42].
333 The signal that led to enhanced IgG binding to red cell surface also were controversial.

334 [3]. Clustering of band-3 protein or proteolytic cleavage of band-3 have been suggested by different workers as
335 trigger for enhanced IgG binding to RBC membrane [40,18,41,23,62]. Beppu et al (1990) [43] and later on other
336 scientist [??] have shown that in vitro enhanced binding of autologous Ig G to red cell surface occurred following
337 an oxidant stress. We have tried to confirm that finding. Our results confirmed that increased binding of anti
338 band-3 immunoglobulins to red cell surface following an oxidative stress by iron and ascorbate . However that
339 was prevented by hydroxyl radical scavengers like mannitol, DMSO and anti oxidant enzyme catalase (Table-5).

340 Results presented in Fig- 1 indicated that oxidant stress to red cell ghosts under our experimental conditions
341 did not lead to protein cross-linking binding of autologous IgG to red cell surface following an oxidative stress .
342 That obviously implied that the two phenomena were inter-related. It was presumable that the bound IgG in our
343 in vitro assay systems were auto antibodies generated earlier in vivo against oxidatively modified proteins with
344 increased carbonylation.

345 Various types of membrane alterations have been reported to aged red cells, which were linked to oxidative
346 damage [??, ??,18,19]. In the present study increased carbonyl content to aged RBC membrane, observed both
347 in spectrophotometric and immunodetection assay system (Fig- ??, Fig- ??) with respect to that of young red
348 cell membrane ghosts represented that the aged red cells have been subjected to oxidative stress in vivo which
349 had been considered as the hall mark of oxidative protein damage [55]. That result seemed interesting, as no such
350 information was available in aged erythrocytes except for an isolated study on stored RBC in blood bank [22].
351 Again, we have observed no significant alteration in membrane protein thiol content of aged red cells compared
352 to young cells (Fig- ??). That was in conformity with our in vitro result, where oxidative stress and ascorbate
353 failed to cause measurable change in protein thiol content of RBC membrane ghosts after incubation up to 2 h
354 (Table-2). Our result [63].

355 Increased binding of autologous IgG to aged red cell membrane have been well documented [40,18,37], although
356 disputes were there regarding precise nature of the auto-antibodies binding to RBC surface [37]. Our results
357 have confirmed this (Table -6 Table-2). On the other hand ?H radical scavengers like mannitol, DMSO and
358 anti oxidant enzyme catalase prevented both the increased incorporation of carbonyl groups to membrane protein
359 (Fig- ??) and enhanced in our study supported the observation of Kannan et al (1991) [64] and the result was also
360 consistent with elevated IgG binding seen in red cells of other species like mouse, dog etc, using different methods
361 for No cross-linking or polymerization of red cell membrane polypeptides were noticed in reducing condition
362 discontinuous SDS-PAGE pattern of aged red cell membranes in comparison to young red cell membranes (Fig-
363 ??). These results were also in accordance with the in vitro data where oxidant treatment of RBC membrane
364 ghosts failed to produce any cross-linking or fragmentation of proteins (Fig- 1). However Rettig et al [??1999] [2]
365 have reported a covalent, non-reducible, non disulphide cross-linking of globin subunits of denatured haemoglobin
366 with membrane at the end of the RBC's life span. Our failure to observe any cross-linking of membrane proteins
367 in SDS-PAGE of aged erythrocytes membrane might be related to running the electrophoresis of aged RBC
368 membrane ghosts in reducing condition [13] or species difference or method of isolation of aged red cells. by Fe
369 2+ was in agreement with the result of Piccinini et al [??1995] separating aged red cells from young red cells [65,
370 ??].

371 Since oxidant stress in vitro led to enhanced binding of autologous IgG to red cell surface (Table-4), it might be
372 implicated that the increased binding of autoantibodies to aged erythrocyte membrane was the result of oxidative
373 injury to red cell membrane during in vivo ageing. This had also been suggested by Beppu et al (1990) [43] and
374 Fujino et al (2000) [??]. The nature of the auto antibody binding to aged red cells have not been elucidated in our
375 study. However, since increase in lipid peroxidation products followed by increased damage of RBC membrane
376 protein as increased carbonyl group incorporation was the only noticeable change in the membrane proteins of
377 aged erythrocytes compared to that of control young erythrocyte, it would be tempting to speculate that auto-
378 antibodies had been directed against such altered membrane proteins. The antibody dependent phagocytosis
379 of red cells had been considered as an important mechanism of removal of old and damaged red cells from the
380 circulation [40,18, ??, ??]. The oxidative modification of membrane protein during ageing of erythrocytes was
381 therefore, directly linked with the subsequent removal of those cells from circulation.

382 20 VI.

383 21 Conclusion


384 From the results and discussion of our study, it can be concluded that in vitro oxidant stress to red blood
385 cell ghosts by iron and ascorbate, led to increased lipid peroxidation products followed by enhanced damage to
386 RBC membrane protein, caused increased membrane protein carbonylation; presumably mediated by hydroxyl
387 radicals and that phenomenon was directly linked to enhanced binding of autologous IgG to oxidized ghosts
388 under similar condition of incubation. It was presumable that bound autologous immunoglobulins to oxidized
389 red blood cell membranes represented auto-antibodies generated in vivo against carbonylated proteins. Further,
390 in vivo senescence of red cells was associated with increased membrane protein carbonylation and enhanced
391 binding of autologous IgG to red cell surface. Since removal of aged red cells from the circulation in large

392 measure depended upon antibody dependent phagocytosis of red cells by macrophages, our results indirectly
393 pointed out that oxidative stress induced modification of RBC membrane carbonylated proteins triggered the
394 latter mechanism for removal of RBC by subsequent formation of auto-antibodies.

395 22 6.53

396 Incubation of erythrocyte ghosts was carried out at 37°C for 2 h in phosphate buffer (50 mM, pH 7.4) with or
397 without addition of FeSO₄ (0.2 mM) and ascorbate (0.5 mM) in presence or absence of other additions like
398 BHT (0.5 mM), mannitol (20 mM), DMSO (20 mM) or catalase (50 ?g/ml) followed by measurement of lipid
399 peroxidation as described in the methods. Results presented were the mean ± SEM of 5 observations (N=5),
400 followed by Student's 't' test (paired). Value marked '*' indicated significant rise of lipid peroxidation with
401 respect to control (erythrocyte ghosts incubated alone with buffer). *p < 0.001. Value marked ' ? ' showed
402 significant inhibition by BHT. ? p < 0.001. Value marked '#' showed no significant inhibition when compared
403 with the value marked '*'.

404 Erythrocyte ghosts incubated with or without FeSO₄ (0.2 mM) and ascorbate (0.5 mM) for 2 h at 37°C and
405 binding of unfractionated autologous IgG (1mg/ml) and anti band-3 depleted autologous IgG (0.8 mg/ml) were
measured as described in the methods. Results presented were mean ± SEM of 6 observations (N=6).¹



25

Figure 1: 25 h

406 2
407

¹© 2014 Global Journals Inc. (US)

²© 2014 Global Journals Inc. (US) VII.





Figure 2:



29

Figure 3: 29 /




Figure 4: Figure 1 :



2339420146

Figure 5: Figure 2 :Figure 3 : 39 Figure 4 :) 2014 CFigure 6 :




Figure 6:




Figure 7:

O₂) or by the action of selenium dependent glutathione peroxidase (GSH glutathione)

catalase ? 2H₂O +

Figure 8:

1 showed that the mixture of Fe²⁺ (0.2 mM) and ascorbate (0.5 mM) produced hydroxyl radicals by Fenton reaction as shown by 2-deoxyribose (DR) degradation assay. The phenomenon was inhibited by hydroxyl radical scavenger like mannitol (20mM) upto 62.4% and DMSO (20 mM) up to 77.4% or by antioxidant was also observed that ferritin (5.6 μM) can also produce OH radical when incubated with 0.5 mM ascorbate (Table-1). Hence iron and ascorbate were physiologically relevant to erythrocyte in vivo and mixture of these components was used as a model system to oxidative stress in vitro.

b) Fig-1 showed that SDS-PAGE analysis of membrane proteins of erythrocyte ghosts incubated with Fe

[Note: Fe²⁺ and ascorbate. In control sample (Lane a) different protein bands of red cell membranes were visible and named according to Fairbanks et al c)]

Figure 9: Table -

1

1. Suzuki T., Dale G.L., Membrane proteins in senescent erythrocyte. *Biochem. J.*, 1989, 257, 37-41.
2. Rettig M.P., Low P.S., Gimm J.A., Mohandas N., Wang J., Christian J.A., Evaluation of biochemical changes during in vivo erythrocyte senescence in dog. *Blood*, 1999, 93, 376-384.
3. Clark M.R., Senescence of red blood cells: Progress and problems. *Physiol. Rev.*, 1988, 68, 503-554.
4. Sauberman N., Fortier N.L., Joshi W., Piotrowski J., Snyder M., Spectrin-haemoglobin cross-linkages associated with in vitro oxidant hypersensitivity in pathologic and artificially dehydrated red cells. *Br. J. Haematol.*, 1983, 54, 15-28.
5. Rice-Evans C., Hochstein P., Alterations in erythrocyte membrane fluidity by phenyl hydrazine. *Res. Commun.*, 1981, 100, 1537-1542.
6. Hebbel R.P., Eaton J.W., Balasingam M., Steinberg M. H., Spontaneous oxygen radical generation by sickle erythrocytes. *J. Clin. Invest.*, 1982, 70, 1253-1259.
7. Fujino T., Ando K., Beppu M., Kikugawa K., Enzymatic removal of oxidized protein aggregates from erythrocyte membranes. *J. Biochem.*, 2000, 127, 1081-1086.
8. Celedon G., Rodriguez I., Espana J., Escobar J., Lissi E., Contribution of haemoglobin and membrane constituents modification to human erythrocyte damage promoted by peroxy radicals of different charge and hydrophobicity. *Free. Radic. Res.*, 2001, 34, 17-31.
9. Takeshita M., Tamura M., Yubisui T., Yoneyama Y., Exponential decay of cytochrome b5 and cytochrome b5 reductase during senescence of erythrocytes: relationship increased methaemoglobin content. *J. Biochem.*, 1983, 93, 931-934.
10. Chakrabarti S., Sonaye B., Naik A. A., Nadkarni P. P., Erythrocyte membrane protein damage by

Figure 10: Table 1 :

2

Year 2014
1 33
Volume XIV Issue VII Version I
D D D D) C
(

[Note: Values were not significantly different from corresponding control values as observed from Student's 't' test (paired).]

Figure 11: Table 2 :

3

Incubation mixture

Figure 12: Table 3 :

4

Year
Volume XIV Issue VII Version I
()

Incubation mixture	Autologous IgG binding (ratio to control)
Oxidized ghost + Unfractionated IgG	4.26 ± 0.175
Oxidized ghost + Anti band-3 depleted IgG	0.98 ± 0.188

[Note: 2014 C © 2014 Global Journals Inc. (US)]

Figure 13: Table 4 :

5

Incubation mixture	Autologous IgG binding (ratio to control)
Oxidized ghost + IgG	4.26 ± 0.176
Catalase treated oxidized ghost + IgG	$0.85 \pm 0.155 *$
Mannitol treated oxidized ghost + IgG	$0.91 \pm 0.159 *$
DMSO treated oxidized ghost + IgG	$0.85 \pm 0.149 *$

Figure 14: Table 5 :

6

Incubation mixture Aged RBC membrane ghost + IgG Aged RBC membrane ghost + IgG	Concentration of IgG mg/ml	1	2.5	0.174	?	3.72 ± 0.183 *	Autologous IgG binding (ratio to control)	2.70 ±	Year 2014 1 35 Volume XIV Issue VII Version I D D D D) C (
--	----------------------------	---	-----	-------	---	----------------	---	--------	--

[Note: Year () 2014 C © 2014 Global Journals Inc. (US)]

Figure 15: Table 6 :

.1 Acknowledgement

- 408
409 Thanks to DST, New Delhi for financial assistance. We also acknowledge to the Calcutta University & Department
410 of Biochemistry, University College of Medicine (IPGME & R, Kolkata), 244B, A.J.C. Bose Road, Kolkata-700
411 020, India, for their kind oxidation products of phenyl hydrazine. *Biochem. Mol. Biol. Int.*, 1995, 35, 255-
412 263. 11. Girotti A. W., Thomas J. P., Jordan J. E., Xanthine oxidase-catalyzed cross-linking of cell membrane
413 proteins. *Arch. Biochem. Biophys.*, 1986, 251, 639-653.
- 414 [T] , T .
- 415 [Winterbourn et al. ()] , C C Winterbourn , B M Mcgrath , R W Carrell . *Biochem. J* 1976. 155 p. .
- 416 [Proc. Natl. Acad. Sci. USA ()] , *Proc. Natl. Acad. Sci. USA* 1989. 86 p. .
- 417 [Galili et al. ()] ‘A unique natural human IgG antibody with anti- β -galactosyl specificity’. U Galili , E A
418 Rachmilewitz , A Peleg , I Flechner . *J. Exp. Med* 1984. 160 p. .
- 419 [Chakraborty et al. ()] ‘Age-related oxidative inactivation of Na⁺, K⁺-ATPase in rat brain crude synaptosomes’.
420 H Chakraborty , P Sen , A Sur , U Chatterjee , S Chakrabarti . *Exp. Gerontol* 2003. 38 p. .
- 421 [Sur et al. ()] ‘Ascorbate induced cross-linking of oxyhemoglobin sub units’. A Sur , K Pal , S Sen , ChakrabartiS
422 . *Ind. J. Exp. Biol* 2000. 38 p. .
- 423 [Frei et al.] *Ascorbate is an outstanding antioxidant in human blood plasma*, B Frei , L England , B N Ames .
- 424 [Hochstein and Jain ()] ‘Association of lipid peroxidation and polymerization of membrane proteins with
425 erythrocyte ageing’. P Hochstein , S K Jain . *Fed. Proc* 1981. 40 p. .
- 426 [Beppu et al. ()] ‘Binding of anti -band 3 autoantibody to oxidatively damaged erythrocytes’. M Beppu , A
427 Mizukami , M Nagoya , K Kikugawa . *J. Biol. Chem* 1990. 265 p. .
- 428 [Levine et al. ()] ‘Carbonyl assays for determination of oxidatively modified proteins’. R L Levine , J A William
429 , E R Stadtman , E Shacter . *Methods. Enzymol* 1994. 233 p. .
- 430 [Margetis et al. ()] ‘carbonylation of red blood cell membrane in diabetic retinopathy’. P I Margetis , M H
431 Antonelou , I K Petropoulos . *Exp. Mol.Pathol* 2009. 87 p. .
- 432 [Laemmli ()] *Cleavage of structural proteins during assembly of head of bacteriophage-Nature*, U K Laemmli .
433 1970. 227 p. .
- 434 [Schluter and Drenckhahn ()] ‘Co-clustering of denatured hemoglobin with band-3: Its role in binding of auto
435 antibodies against band 3 to abnormal and aged erythrocytes’. K Schluter , D Drenckhahn . *Proc. Natl. Acad.*
436 *Sci. USA* 1986. 83 p. .
- 437 [Sorette et al. ()] ‘Comparison of serum anti-band 3 and anti-gal antibody binding to density separated human
438 red cells’. M P Sorette , U Galili , M R Clark . *Blood* 1991. 1989. 77 p. .
- 439 [Amici et al. ()] ‘Conversion of amino acid residues in proteins and amino acid homopolymers to carbonyl
440 derivatives by metal-catalyzed oxidation reactions’. A Amici , R L Levine , L Tsai , E R Stadtman . *J.*
441 *Biol. Chem* 1989. 264 p. .
- 442 [Shaklai et al. ()] ‘Cross-linking of isolated cytoskeletal proteins with hemoglobin: A possible damage inflicted
443 to the red cell membrane’. N Shaklai , B Frayman , N Fortier , M Snyder . *Biochim. Biophys. Acta* 1987. 915
444 p. .
- 445 [Levine et al. ()] ‘Determination of carbonyl content in oxidatively modified proteins’. R L Levine , D Garland ,
446 C N Oliver , A Amici , I Climent , A G Lenz , B W Ahn , D Shaltiel , E R Stadtman . *Methods. Enzymol*
447 1990. 186 p. .
- 448 [Mendez et al. ()] ‘Differential carbonylation of cytoskeletal proteins in blood group O erythrocytes: potential
449 role in protection against severe malaria’. D Mendez , M L Hernaez , A N Kamali , A Diez , A Puyet , J M
450 Bautista . *Infect. Genet. Evol* 2012. 12 p. .
- 451 [Jarolim et al. ()] ‘Effect of hemoglobin oxidation products on the stability of red cell membrane skeletons and
452 associations of skeletal proteins: correlation with a release of hemin’. P Jarolim , M Lahav , S C Liu , J Palek
453 . *Blood* 1990. 76 p. .
- 454 [Fairbanks et al. ()] ‘Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane’. G
455 Fairbanks , T L Steck , D F H Wallach . *Biochemistry* 1971. 10 p. .
- 456 [Singer et al. ()] ‘Erythrocyte homeostasis: antibody mediated recognition of the senescent state by
457 macrophages’. J A Singer , L K Jennings , C W Jackson , M E Dockter , M Morrison , W S Walker .
458 *Proc. Natl. Acad. Sci. USA* 1986. 83 p. .
- 459 [Puppo and Halliwell ()] ‘Formation of hydroxyl radicals from hydrogen peroxide in the presence of iron: Is
460 haemoglobin a biological Fenton reagent?’. A Puppo , B Halliwell . *Biochem. J* 1998. 249 p. .
- 461 [O’connel et al. ()] ‘Formation of hydroxyl radicals in the presence of ferritin and haemosiderin: Is haemosiderin
462 formation a biological protective mechanism?’. M O’connel , B Halliwell , C P Moorhouse , O I Aruoma , H
463 Baum , T J Peters . *Biochem. J* 1986. 234 p. .

- 464 [Halliwell and Gutteridge ()] 'Formation of thiobarbituric acid-reactive-substance from deoxyribose in the pres-
465 ence of iron salts: The role of superoxide and hydroxyl radicals'. B Halliwell , J M C Gutteridge . *FEBS. Lett*
466 1981. 128 p. .
- 467 [Halliwell and Gutteridge (ed.) ()] *Free Radicals in biology and Medicine*, B Halliwell , J M C Gutteridge .
468 Halliwell B., Gutteridge J.M.C (ed.) 1989. Oxford: Clarendon Press. (2nd edition)
- 469 [Margaritis et al. ()] 'Glutathione metabolism in red cell ageing'. L H Margaritis , I S Papassideri , H Imanishi ,
470 T Nakai , T Abe , T Takino . *Mech. Ageing. Dev* 1985. 32 p. . (Increased protein 24)
- 471 [Rennie et al. ()] 'Human erythrocyte fractionation in "percoll" density gradient'. C M Rennie , S Thompson , A
472 C Parker , A Maddy . *Clin. Chim. Acta* 1979. 98 p. .
- 473 [Kannan et al. ()] 'Isolation and partial characterization of antibody and globin-enriched complexes from
474 membranes of dense human erythrocytes'. R J Kannan , J Yuan , P S Low . *Biochem. J* 1991. 278 p.
475 .
- 476 [Leaverton ()] P E Leaverton . *Statistical Inference: Tests for Statistical Significance*, P E Leaverton (ed.)
477 (Boston) 1995. p. . (5th edition Little Brown and Company)
- 478 [Chakraborty et al. ()] 'Lipid peroxidation associated protein damage in rat brain crude synaptosomal fraction
479 mediated by iron and ascorbate'. H Chakraborty , S N Ray , S Chakrabarti . *Neurochem. Int* 2001. 39 p. .
- 480 [Chakrabarti and Shankar ()] 'Lipid peroxidation in developing rat brain during undernutrition'. S Chakrabarti
481 , R Shankar . *Neurosci. Lett* 1984. 48 p. .
- 482 [Kay ()] 'Localization of senescent cell antigen on band 3'. M M B Kay . *Proc. Natl. Acad. Sci. USA*, (Natl.
483 Acad. Sci. USA) 1984. 81 p. .
- 484 [Beppu et al. ()] 'Macrophage recognition of periodate-treated erythrocytes: Involvement of disulfide formation
485 of the erythrocyte membrane protein'. M Beppu , H Ochiai , K Kikugawa . *Biochim. Biophys. Acta* 1989. 979
486 p. .
- 487 [Campwala and Desforges ()] 'Membrane bound hemichrome in density separated cohorts of normal (AA) and
488 sickled (SS) cells'. H Q Campwala , J F Desforges . *J. Lab. Clin. Med* 1982. 99 p. .
- 489 [Lutz ()] 'Naturally occurring auto antibodies in mediating clearance of senescent red blood cells'. H U Lutz .
490 *Adv. Exp. Med. Biol* 2012. 750 p. .
- 491 [Hernandez-Munoz et al. ()] 'Oxidant status and lipid composition of erythrocyte membranes in patients with
492 type 2 diabetes, chronic liver damage, and a combination of both pathologies'. R Hernandez-Munoz , M
493 Olguin-Martinez , I Aguilar-Delfin , L Sanchez-Sevilla , N Garcia-Garcia , M Diaz-Mufioz . *Oxid. Med. Cell.*
494 *Longev* 2013. 2013. p. .
- 495 [Mazzulla et al. ()] 'Oxidation of human red blood cells by a free radical initiator: Effects on rheological
496 properties'. S Mazzulla , A Schella , D Gabriele , N Baldino , S Sesti , E Perrotta , A Costabile , B De
497 Cindio . *Clin. Hemorheol. Microcirc* 2014. (In press)
- 498 [Piccini et al. ()] 'Oxidation state of glutathione and membrane proteins in human red cells of different age'. G
499 Piccini , G Minetti , C Balduini , A Brovelli . *Mech. Ageing. Dev* 1995. 78 p. .
- 500 [Shinar et al.] 'Oxidative damage to human red cells induced by copper and iron complexes in the presence of
501 ascorbate'. E Shinar , A Rachmilewitz , A Shifter , E Rahamin , P Saltman . *Biochim. Biophys. Acta*
- 502 [Szebeni et al. ()] 'Oxidative interactions between haemoglobin and membrane lipid: A liposome model'. J
503 Szebeni , C C Winterbourn , R W Carrell . *Biochem. J* 1984. 220 p. .
- 504 [Rinalducci et al. ()] 'Oxidative stress and Capase-mediated fragmentation of cytoplasmic domain of erythrocyte
505 band 3 during blood storage'. S Rinalducci , E Ferru , B Blasi , F Turrini , L Zolla . *Blood Transfus* 2012. 10
506 (2) p. .
- 507 [Halliwell and Gutteridge] 'Oxygen toxicity, oxygen radicals, transition metals and disease'. B Halliwell , J M C
508 Gutteridge . *Biochem. J.* , ! 984 p. .
- 509 [May et al. ()] 'Protection and recycling of α -tocopherol in human erythrocytes by intracellular ascorbic acid'. J
510 M May , Z C Qu , S Mendiratta . *Arch. Biochem. Biophys* 1998. 349 p. .
- 511 [Dumaswala et al. ()] 'Protein and lipid oxidation of banked human erythrocytes: Role of glutathione'. U J
512 Dumaswala , L Zhuo , D W Jacobsen , S K Jain , K A Sukalski . *Free. Radic. Biol. Med* 1999. 27 p. .
- 513 [Lowry et al. ()] 'Protein measurement with Folin phenol reagent'. O H Lowry , N J Rosebrough , A L Farr , R
514 J Randall . *J. Biol. Chem* 1951. 193 p. .
- 515 [Habeeb ()] 'Reaction of protein sulfhydryl groups with Ellman's reagent'. A F S A Habeeb . *Methods. Enzymol*
516 1972. 25 p. .
- 517 [Reactions involving superoxide and normal and unstable oxidatively damaged red blood cells in mice Blood ()]
518 'Reactions involving superoxide and normal and unstable oxidatively damaged red blood cells in mice'. *Blood*
519 2000. 95 p. .

- 520 [Chancerelle et al. ()] 'Recognition and elimination of senescent erythrocytes: Implication of antibodies specific
521 for malonic dialdehyde-protein adducts, as demonstrated by flow cytometry'. Y Chancerelle , J Mathieu , J
522 F Kergonou . *Biochem. Mol. Biol. Int* 1994. 34 p. .
- 523 [Sambrano et al. ()] 'Recognition of oxidatively damaged erythrocytes by a macrophage receptor with specificity
524 for oxidized low density lipo protein'. G R Sambrano , S Parthasarathy , D Steinberg . *Proc. Natl. Acad. Sci.*
525 *USA* 1994. 91 p. .
- 526 [Terpstra et al. ()] 'Scavenger receptors on liver Kupffer cells mediate the in vivo uptake of 53'. V Terpstra , J C
527 Vanberkel , H Ohkawa , N Oshini , K Yagi . *Anal. Biochem* 1979. 95 p. . (Assay for lipid peroxides in animal
528 tissues by thiobarbituric acid reaction)
- 529 [Weed ()] 'The importance of erythrocyte deformability'. R I Weed . *Am. J. Med* 1970. 49 p. .
- 530 [Hodgson and Fridovich ()] 'The interaction of bovine erythrocyte superoxide dismutase with hydrogen peroxide:
531 inactivation of the enzyme'. E K Hodgson , I Fridovich . *Biochemistry* 1975. 14 p. .
- 532 [Williams and Morris ()] 'The internal viscosity of the human erythrocyte may determine its life span in vivo'.
533 A R Williams , D R Morris . *Scand. J. Haematol* 1980. 24 p. .
- 534 [Browne et al. ()] 'The molecular pathobiology of cell membrane iron:The sickle red cell as a model'. P Browne
535 , O Shalev , R P Hebbel . *Free. Radic. Biol. Med* 1998. 24 p. .
- 536 [Beutler et al. ()] 'The removal of leukocytes and platelets from whole blood'. E Beutler , C West , K G Blume
537 . *J. Lab. Clin. Med* 1975. 88 p. .
- 538 [Low et al. ()] 'The role of haemoglobin denaturation and band-3 clustering in red blood cell ageing'. P S Low ,
539 S M Waugh , K Zinke , D Drenckhahn . *Science* 1985. 227 p. .