Online ISSN: 2249-4618
Print ISSN: 0975-5888

# Global Journal

## of Medical Research: K

## Interdisciplinary



Effect of Aerobic Training

Impact of Low Frequency

**Highlights**

Gestation Diabetic Detection

Lymphatic Drainage on Triglycerides

Discovering Thoughts, Inventing Future

VOLUME 15 ISSUE 6 VERSION 1.0

© 2001-2015 by Global Journal of Medical Research , USA

Volume 15 Issue 42 was originally issued as K6

Global Journal of Medical Research: K
Interdisciplinary

Volume 15 Issue 42
Originally issued as K6

# Global Journal of Medical Research: K
# INTERDISCIPLINARY

Volume 15 Issue 6 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Medical Research . 2015.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

**Incorporation No.: 0423089**
**License No.: 42125/022010/1186**
**Registration No.: 430374**
**Import-Export Code: 1109007027**
**Employer Identification Number (EIN):**
**USA Tax ID: 98-0673427**

## Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

*Sponsors: Open Association of Research Society*
*Open Scientific Standards*

### *Publisher's Headquarters office*

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
*USA Toll Free: +001-888-839-7392*
*USA Toll Free Fax: +001-888-839-7392*

### *Offset Typesetting*

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

### *Packaging & Continental Dispatching*

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

### *Find a correspondence nodal officer near you*

To find nodal officer of your country, please email us at *local@globaljournals.org*

### *eContacts*

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

### *Pricing (Including by Air Parcel Charges):*

*For Authors:*
22 USD (B/W) & 50 USD (Color)
*Yearly Subscription (Personal & Institutional):*
200 USD (B/W) & 250 USD (Color)

# Integrated Editorial Board
## (Computer Science, Engineering, Medical, Management, Natural Science, Social Science)


**John A. Hamilton,"Drew" Jr.,**
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

**Dr. Henry Hexmoor**
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

**Dr. Osman Balci, Professor**
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

**Yogita Bajpai**
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

**Dr. T. David A. Forbes**
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

**Dr. Wenying Feng**
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

**Dr. Thomas Wischgoll**
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

**Dr. Abdurrahman Arslanyilmaz**
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

**Dr. Xiaohong He**
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

**Burcin Becerik-Gerber**
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley
& Istanbul University

**Dr. Bart Lambrecht**
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

**Dr. Carlos García Pont**
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

**Dr. Fotini Labropulu**
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

**Dr. Lynn Lim**
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

**Dr. Mihaly Mezei**
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

**Dr. Söhnke M. Bartram**
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

**Dr. Miguel Angel Ariño**
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

**Philip G. Moscoso**
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

**Dr. Sanjay Dixit, M.D.**
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

**Dr. Han-Xiang Deng**
MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University
Feinberg School of Medicine

**Dr. Pina C. Sanelli**
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

**Dr. Roberto Sanchez**
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

**Dr. Wen-Yih Sun**
Professor of Earth and Atmospheric SciencesPurdue University Director
National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

**Dr. Michael R. Rudnick**
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

**Dr. Bassey Benjamin Esu**
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

**Dr. Aziz M. Barbar, Ph.D**.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

## President Editor (HON.)

**Dr. George Perry, (Neuroscientist)**
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

## Chief Author (HON.)

**Dr. R.K. Dixit**
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

## Dean & Editor-in-Chief (HON.)

**Vivek Dubey(HON.)**
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

**Sangita Dixit**
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

**Suyash Dixit**
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

**Er. Suyog Dixit**
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

**Pritesh Rajvaidya**
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

**Luis Galárraga**
J!Research Project Leader
Saarbrücken, Germany

# Contents of the Issue


i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue

1. A Gestation Diabetic Detection Technique using Muscle Energy Derived from Surface EMG. *1-11*
2. Effect of Aerobic Training on Ventricular Remodeling After Percutaneous Coronary Intervention *13-18*
3. The Impact of Low Frequency Ultrasound and Lymphatic Drainage on Triglycerides. *19-28*
4. MRI-T1 and T2 Image Fusion for Brain Image using CDF Wavelet based on Lifting Scheme. *29-33*

v. Fellows
vi. Auxiliary Memberships
vii. Process of Submission of Research Paper
viii. Preferred Author Guidelines
ix. Index

Global Journal of Medical Research: K
Interdisciplinary
Volume 15 Issue 6 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

# A Gestation Diabetic Detection Technique using Muscle Energy Derived from Surface EMG

By Anjaneya L. H, Mallikarjun S. Holi & Dr. S. Chandrasekhar

*Abstract-* Electromyogram (EMG) is one among the important biopotential signal reflecting the human skeletal muscle activity. EMG signals can be used for many biomedical applications pertaining to diagnosis and therapy of musculoskeletal and rheumatological problems. EMG signals are complex in nature and require advanced techniques for analysis, such as decomposition, detection, processing, and classification. Diabetes mellitus is a chronic metabolic disorder characterized by elevated levels of blood glucose. The musculoskeletal system can be affected by diabetes in a number of ways. The main aim of the paper is to identify the diabetic patient and show the classification performance of the proposed framework. In this paper EMG signal is investigated by feature extraction and are classified into normal and diabetic for comprehension of EMG signal. The primary point of this work is to recognize the diabetes utilizing different elements and to demonstrate the performance of the proposed framework. The obtained results demonstrate that the extracted feature in proposed framework displays better performance for classification the EMG signal contrasted with alternate elements. Based on the impacts of features on the EMG signal classification, different results were obtained through analysis of the SVM Classification. Experimental study shows that the proposed method's classification accuracy is 98.98%.

*Keywords:* *feature extraction; electromyography (EMG) signal; SVM classifier.*

*GJMR-K Classification:* *NLMC Code: QZ 4*




*Strictly as per the compliance and regulations of:*

© 2015. Anjaneya L. H, Mallikarjun S. Holi & Dr. S. Chandrasekhar. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# A Gestation Diabetic Detection Technique using Muscle Energy Derived from Surface EMG

Anjaneya L. H [α], Mallikarjun S. Holi [σ] & Dr. S. Chandrasekhar [ρ]

*Abstract-* Electromyogram (EMG) is one among the important biopotential signal reflecting the human skeletal muscle activity. EMG signals can be used for many biomedical applications pertaining to diagnosis and therapy of musculoskeletal and rheumatological problems. EMG signals are complex in nature and require advanced techniques for analysis, such as decomposition, detection, processing, and classification. Diabetes mellitus is a chronic metabolic disorder characterized by elevated levels of blood glucose. The musculoskeletal system can be affected by diabetes in a number of ways. The main aim of the paper is to identify the diabetic patient and show the classification performance of the proposed framework. In this paper EMG signal is investigated by feature extraction and are classified into normal and diabetic for comprehension of EMG signal. The primary point of this work is to recognize the diabetes utilizing different elements and to demonstrate the performance of the proposed framework. The obtained results demonstrate that the extracted feature in proposed framework displays better performance for classification the EMG signal contrasted with alternate elements. Based on the impacts of features on the EMG signal classification, different results were obtained through analysis of the SVM Classification. Experimental study shows that the proposed method's classification accuracy is 98.98%.

*Keywords: feature extraction; electromyography (EMG) signal; SVM classifier.*

## I. Introduction

For the evaluation and administration of patient health, observing of physiological and physical signal is crucial. Electromyogram is a critical health pointer. EMG is likewise utilized as a part of numerous sorts of exploration labs, incorporating those included in engine control, neuromuscular physiology, biomechanics, development issue, postural control, and exercise based recuperation. Physiological and anatomical properties of muscles are presented by the signals; an EMG sign is the electrical action of a muscle's engine units, which comprise of two sorts: surface EMG, and intramuscular EMG [1]. Surface EMG and intramuscular EMG signs are recorded by non-intrusive cathodes and obtrusive terminals, separately. Nowadays, surface-identified signs are ideally used to acquire data about the time or power of shallow muscle enactment [2].

Electromyography (EMG) signs are viewed as most valuable as electrophysiological signs in both medical and engineering fields. The essential strategy for comprehension the human body's practices under typical and neurotic conditions is given by the recording of EMG signs. At whatever point an EMG sign is being recorded from the muscle, different sorts of clamors defile it. In this way, dissecting and characterizing the EMG signs is exceptionally troublesome on account of the complicated pattern of the EMG, particularly when EMG movement happens. EMG is controlled by sensory system and relies on upon anatomical and mental properties of muscles. It is an electrical sign gained from diverse organs. EMG is typically an element of time, discussed in terms of amplitude, phase and frequency. Electromyography (EMG) signs presents to the electrical movement of a muscle amid compression [1].

These signs may be recognized from skin surface anodes or from needles set specifically inside of the muscle. These two sorts of recordings are utilized for diverse purposes, with needle recording used to recognize the conduct of individual muscle motor and fiber units while surface recordings are utilized to distinguish general muscle action specifically positions or activities. Surface EMG is not a typical clinical technique; however it might be utilized as a part of restoration (rehabilitation). Needle electromyography is utilized to figure out if there is harm to nerve filaments to individual muscles. At the point when nerve sends the sign to start muscle constriction a potential is produced over the muscle because of the stream of particles all through muscle cells (i.e. developments of electrolytes) and this ionic current is changed over into electronic current with Ag-AgCl anodes put on the surface of the skin of the contracting muscle. Recently a surface electromyography is progressively utilized for recording from shallow muscles in clinical conventions, where intramuscular cathodes are utilized for profound muscle only [3].

The innovation of EMG is moderately new. There are still restrictions in characterization and detection of EMG sign, phase estimation, procuring suitable data because of induction from typicality. Conventional framework remaking calculations have different impediments and extensive computational complexity quality and numerous show high differences.

*Author α: Biomedical-Engineering, BIET, Davangere, India. e-mail: anjudvg@yahoo.com*
*Author σ: Electronics & Instrumentation Engineering, U.B.D.T., VTU, India. e-mail: msholi@yahoo.com*
*Author ρ: Emergency Medicine, J.J.M. Medical College, Davangere, India. e-mail: drschandru@yahoo.com*



© 2015 Global Journals Inc. (US)

Recently in innovations of signal handling and numerical models have made it to create progressed EMG location and examination strategies [4]. In this way, look into and broad researches have been made in the zone, growing better algorithms, overhauling existing approaches, enhancing recognition methods to diminish clamor, and to gain exact EMG signals. It is very imperative to do an examination to group the real issues of EMG signs investigation and legitimize the acknowledged measures. Higher-order factual routines may be utilized for analyzing the EMG signal because of the one of a kind properties of measurable techniques connected to arbitrary time arrangement. This paper identifies with the redesigning existing procedures filtering, processing, decomposition and modeling of EMG. In our proposed work our principle expect to recognize and order the diabetic patient for that EMG signal information has been considered to do the work. By utilizing the EMG signal it can be dissected what nerves have been harmed and how broad that harm is. An electromyography (EMG) test is often done in conjunction with a Nerve Conduction Velocity (NCV) test. It shows how well muscles are receiving signals from the nerves. Damaged nerves won't send clear or consistent messages.

Reminder of this paper is organized as follows: Section 2 presents the related work in this area. Section 3 presents proposed method for classifying EMG diabetic signal. Section 4 gives detailed description about the result obtained. And the paper is concluded in Section 5.

## II. Related Work

This section provides the previous researches in the field of EMG signal processing i.e. feature extraction, noise removal, filtering and classification.

Nishikawa and Kuribayashi et al. [1], Used neural system to segregate hand movements for EMG-Controlled Prostheses. Here the neural system was utilized to take in the connection between EMG signal's energy spectrum and the movement errand craved by the incapacitated subjects.

*Xiao Hu;* Qun Yu et al. [2] presented a novel and basic technique to extricate the general element of two surface EMG sign examples: lower arm supination surface EMG flag and lower arm pronation surface EMG signal. The system decays surface EMG signal into 16 Frequency groups $(FB)$ by wavelet bundle change $(WPT)$, and afterward wavelet coefficient entropy $(WCE)$ of two picked FBs is ascertained. The two WCEs were utilized to recognize FS surface EMG signals from lower arm supination surface EMG signals. The outcome demonstrates that $WCE$ is a powerful technique for removing the component from surface EMG signal.

*kouchaki, s.;* Boostani et al. [3] proposed a compelling combinational feature to upgrade the accuracy of classification among the control group and subjects with neuropathy and myopathy illnesses. All EMG signs were create and simulated artificially, by fusing measurable and morphological properties of every group into their sign models, in the EMG lab of Waterloo University. To characterize the subjects by the proposed system, in the first place, EMG signs are deteriorated by observational mode decay or Empirical mode decomposition(EMD) to its regular subspaces, then number of subspaces is adjusted through every windowed sign, and Kolmogorov Complexity (KC) and other informative component are resolved to uncover the measure of anomaly inside of every subspace. Finally, these elements are connected to support vector machine (SVM).

*ZhaojieJu;* Gaoxiang Ouyang et al. [4] proposed and assessed systems for nonlinear feature extraction and nonlinear classification to recognize distinctive hand controls taking into account surface electromyography (sEMG) signals. The nonlinear measures are accomplished in light of the repeat plot to represent to dynamical attributes of sEMG amid hand movements. Fuzzy Gaussian Mixture Models $(FGMMs)$ are proposed and utilized as a nonlinear classifier to perceive distinctive hand handles and close by controls caught from diverse subjects. Different trials are led to assess their execution by looking at 14 individual elements, 19 multifeatures and 4 distinct classifiers. The test results exhibit the proposed nonlinear measures give vital supplemental data and they are key to the great execution in multifeatures.

*Artug, N.T.;* Goker et al. [5] proposed another system for feature extraction. In this study another dataset are prepared for neuromuscular sicknesses utilizing checking EMG strategy and four new components are extricated. These components are described as duration of phase, maximum amplitude the maximum amplitude, and maximum amplitude times phase duration, and number of peaks. By utilizing factual values, for example, mean and variance, number of elements has expanded up to eight. This dataset was characterized by utilizingk-nearest neighbors calculation $(k-NN)$, radial basis function networks $(RBF)$, support vector machines $(SVM)$, andmulti-layer perceptron $(MLP)$.

## III. Proposed System

As a result of the different clamors and antiques identified among EMG signs, obliged data remains an amalgam inside the raw EMG signals. Then again, if these raw signals are utilized as a data as a part of sEMG order, the proficiency of the classifier reduces. To enhance the performance of the classifier, researchers have been utilizing distinctive sorts of EMG elements as

© 2015 Global Journals Inc. (US)

an information to the classifier. To accomplish ideal order execution, the properties of EMG highlight space should be taken into consideration. There are three sorts of EMG components in diverse spaces: time area, frequency area and time-frequency space features. Hudgins et al. created time area elements of the sEMG. They utilized mean absolute value (MAV), mean absolute value slope, slope sign changes (SSC), waveform lengths (WL) and zero crossing (ZC) for presenting to myoelectric examples. These components are termed as 'the Hudgins highlight'. A deliberately chose set of info components gives a higher grouping rate than the crude sign. In the journey to enhance, the precision of myoelectric sign example grouping Englehart et al. looked at time space (TD) components utilized by Hudgins with the time recurrence area highlights (TFD). Time-recurrence area elements are viable capabilities particularly for transient myoelectric sign example characterization. Because of the high dimensionality and high-determination issue of time-frequency representation, dimensionality lessening is regularly an important supplement to highlight extraction. Every one of them should be possible progressively and electronically and it is straightforward for usage. Feature in this group are regularly utilized for onset identification, muscle constriction and muscle action recognition.





*Figure 1 :* Overall System Architecture

Above given fig1 presents the overall architecture of the framework. To collect the EMG data surface electrodes are being used, this data is called raw data. To perform the desired operation on this data we need to do some preprocessing on this data i.e. amplification, filtering and sampling. The next step is to perform segmentation on the data. After this step the data is divided into samples which are compatible for feature extraction. In our proposed framework various features are extracted which are described below.

Fifteen time domain features and frequency domain features are described in this section. Thirteen time domain variables are measured as a function of time. Because of their computational simplicity, time domain features or linear techniques are the most popular in EMG pattern recognition. Integrated EMG, Mean absolute value, Modified mean absolute value 1, Modified mean absolute value 2, Mean absolute value slope, Willison amplitude Zero crossing, Root mean square, Slope sign change, Variance of EMG, Waveform length and Histogram of EMG are used to test the performance. Additionally, highlights in recurrence area are utilized to speak to the identify muscle weakness and neural anomalies, and at some point are utilized as a part of EMG example acknowledgment. Three customary and two adjusted elements in recurrence range are performed specifically autoregressive coefficients, mean and middle frequencies, changed mean and middle frequencies. A while later, the assessment techniques for two models that used to quantify the power property of the entire components



© 2015 Global Journals Inc. (US)

are presented. Finally the classification is done on the feature extracted data.

## IV. Feature Extraction in Time Domain

*a) Integrated EMG*

Integrated EMG ($IEMG$) is computed as the summation of the total estimations of the $sEMG$ signal amplitude. For the most part, $IEMG$ is utilized as an onset list to recognize the muscle movement that used to approaching the control charge of assistive control gadget. It is identified with the $sEMG$ sign arrangement terminating point, which can be communicated as

$$IEMG = \sum_{n=1}^{N} |x_n|, \tag{1}$$

where $N$ denotes the length of the signal and $x_n$ represents the $sEMG$ signal in a segment.

*b) Mean Absolute Value*

Mean Absolute Value ($MAV$) is like normal corrected quality. It can be figured utilizing the moving normal of full-wave amended EMG. It is ascertained by taking the normal of the total estimation of $sEMG$ sign. It is a simple route for location of muscle compression levels and it is a prominent element utilized as a part of myoelectric control application. It is characterized as

$$MAV = \frac{1}{N}\sum_{N=1}^{N} |x_n| \tag{2}$$

*c) Modified Mean Absolute Value 1*

It is an addition in MAV using weighting window function$w_n$. It can be defined as

$$MAV1 = \frac{1}{N}\sum_{n=1}^{N} w_n |x_n| \tag{3}$$

$$w_n = f(x) = \begin{cases} 1, if\ 0.25N \le n \le 0.75N \\ 0.5 \qquad\quad , otherwise \end{cases}$$

*d) Modified Mean Absolute Value 2*

Modified Mean Absolute Value 2 ($MMAV2$) is similar to $MMAV1$. In this method by using continuous weighting window function$w_n$ , the smooth window is improved. It is given by

$$MMAV2 = \frac{1}{N}\sum_{n=1}^{N} w_n |x_n| \tag{4}$$

$$w_n = \begin{cases} 1, if\ 0.25N \le n \le 0.75N \\ \dfrac{4n}{N}, if\ 0.25N > n \\ \dfrac{4(n-N)}{N}, if\ 0.75 < n. \end{cases}$$

*e) Mean Absolute Value Slope*

Mean Absolute Value Slope ($MAVSLP$) is a modified version of $MAV$. The differences between the $MAVs$ of adjacent segments are determined. It can be defined as

$$MAVSLP_i = MAV_{i+1} - MAV_i \tag{5}$$

*f) Variance of EMG*

Variance of EMG ($VAR$) uses the power of the $sEMG$ signal as a feature. Generally, the variance is the mean value of the square of the deviation of that variable. However, mean of EMG signal is close to zero. In consequence, variance of $EMG$ can be calculated by

$$VAR = \frac{1}{N-1}\sum_{n=1}^{N} x_n^2 \tag{6}$$

*g) Root Mean Square*

Root Mean Square (RMS) is modeled as amplitude modulated Gaussian random process whose RMS is related to the constant force and non-fatiguing contraction. It relates to standard deviation, which can be expressed as

$$RMS = \sqrt{\frac{1}{N}\sum_{n=1}^{N} x_n^2} \tag{7}$$

*h) Waveform Length*

It is a measurement of cumulative length of the waveform over time segment. $WL$is related to the waveform amplitude, frequency and time. It is given by

$$WL = \sum_{n=1}^{N-1} |x_{n+1} - x_n| \tag{8}$$

All of these features above, eq.1-eq.8 are computed based on sEMG signal amplitude. From the experimental results, the pattern of these features is similar. Hence, we selected the robust feature representing for the other features in this group. The results and discussion is presented in Section 4.

*i) Zero Crossing*

Zero intersection (ZC) is the quantity of times that the adequacy estimation of sEMG sign crosses the zero y-axis. In EMG highlight, the edge condition is utilized to go without the background clamor. This component gives a surmised estimation of recurrence space properties. It can be detailed as



© 2015 Global Journals Inc. (US)

$$ZC = \sum_{n=1}^{N-1} [\, sgn(x_n \times x_{n+1}) \cap |x_n - x_{n+1}| \geq threshold] \quad (10)$$

$$sgn(x) \begin{cases} 1, & if\ x \geq threshold \\ 0, & otherwise \end{cases}$$

*j) Slope Sign Change*

Slope Sign Change (SSC) is like ZC. It is another technique to present to the recurrence data of sEMG sign. The quantity of changes in the middle of positive and negative slant among three continuous portions is performed with the edge capacity for keeping away from the impedance in sEMG signal. The count is characterized as

$$SSC = \sum_{n=2}^{N-1} [f[(x_n - x_{n-1}) \times (x_n - x_{n+1})]]; \quad (11)$$

$$f(x) = f(x) = \begin{cases} -1, & if\ x \geq threshold \\ 0 & , \quad otherwise \end{cases}$$

*k) Willison Amplitude*

Willison amplitude (WAMP) is the quantity of times that the contrast between sEMG signal adequacy among two contiguous portions that surpasses a predefined limit to lessen commotion impacts same as ZC and SSC. The definition is as

$$WAMP = \sum_{n=1}^{N-1} f(|x_n - x_{n+1}|); \quad (12)$$

$$f(x) = \begin{cases} 1, & if\ x \geq threshold \\ x & , \quad otherwise \end{cases}$$

WAMP is related to the firing of motor unit action potentials (MUAP) and the muscle contraction level. The suitable value of threshold parameter of features in ZC, SSC, and WAMP is normally chosen between 10and 100 mV that is dependent on the setting of gain value of instrument. Nevertheless, the optimal threshold that suitable for robustness in sEMG signal analysis is evaluated and discussed in Section 4.

## V. Frequency Domain Feature Extraction

*a) Autoregressive Coefficients*

Autoregressive (AR) model described each sample of sEMG signal as a linear combination of previous samples plus a white noise error term. AR coefficients are used as features in EMG pattern recognition. The model is basically of the following form:

$$x_n = -\sum_{i=1}^{p} a_i x_{n-i} + w_n, \quad (13)$$

where $x_n$ is a sample of the model signal,$w_n$ is AR coefficients,$w_n$ is white noise or error sequence, and $p$ is the order of $AR$ model. The forth order AR was suggested from the previous research [19]. However, the orders of AR between the first order and the tenth order are found. The results are discussed in Section 4.

*b) Modified Median Frequency*

Modified Median Frequency (MMDF) is the frequency at which the spectrum is divided into two regions with equal amplitude. It can be expressed as

$$\sum_{j=1}^{MMDF} A_j = \sum_{j=MMDF}^{M} A_j = \frac{1}{2}\sum_{j=1}^{M} A_j, \quad (14)$$

where $A_j$ is the sEMG amplitude spectrum at frequencybin $j$.

*c) Modified Mean Frequency*

Modified Mean Frequency (MMNF) is the average frequency. MMNF is calculated as the sum of the product of the amplitude spectrum and the frequency, divided by the total sum of spectrum intensity, as in

$$MMNF = \sum_{j=1}^{M} \frac{f_j A_j}{\sum_{j=1}^{M} A_j}, \quad (15)$$

where $f_j$ is the frequency of spectrum at frequency bin $j$.

According to the proposed method for EMG classification the above given mathematical equation are used for the frequency and time domain feature extraction.

Because of the way of the EMG signal, there would be an extensive variety in the estimation of specific features between individuals. Numerous elements, for example, changes in terminal position, sign preparing will deliver changes in highlight values after some time. A suitable acknowledgment technique must have the capacity to oblige the normal individual contrast. SVM Classifier hypothesis assumes an imperative critical part in managing vulnerability when settling on choices in biomedical field of uses. The classifier proposed for the grouping of the EMG signs was executed by a basic methodology in view of Support Vector Machine is to arrange the EMG sign to one of the classifications, Diabetic or Non-Diabetic.



© 2015 Global Journals Inc. (US)

## VI. Results and Discussion

*a) Data Recording Stage*

The EMG records correspond to the activity of the uterine muscles and might therefore be used to predict the premature onset of labor. Records were collected from the general population as well as from the patients admitted to the hospital with the diagnosis of impending pre-term labor. One record pre-pregnancy was recorded. The records are of 30-minduration and consist of three channels. The sampling frequency, fs, was 20 Hz. The records were collected from the abdominal surface using four AgCl2 electrodes. The electrodes were placed in two horizontal rows, symmetrically under and above the navel, spaced7 cm apart. A special protocol was used during the attachment of the electrodes in order to improve the quality of the measurements [13]. According to the protocol, the resistance between the electrodes had to be lower than $100\ k\Omega$. The first acquired signal was measuredbetween the topmost electrodes (E2–E1), the second signal between the leftmost electrodes (E2–E3) and the third signal between the lower electrodes (E4–E3). Prior to sampling the signals were filtered using an analog three pole Butterworth filter with the bandwidth from 0 to 5 Hz. The resolution of the scanning system was 16 bits with the amplitude range $\pm 2.5$ mV.

In our experiment we have considered 297 users EMG signal data. Out of 297, 295users are not having diabetes and the remaining 2 users are diabetic affected.

The given table 1 shows the mean and standard deviation for all the features for diabetic patient.

*Table 1 :* Mean and standard deviation measurement of all the features

| FEATURES | MEAN | STD |
|---|---|---|
| Integrated EMG | -59.609 | 503.9653 |
| Mean Absolute Value | 246.3694 | 132.842 |
| Modified Mean Absolute Value 1 | 187.0663 | 102.1839 |
| Modified Mean Absolute Value 2 | 124.7915 | 69.9963 |
| Mean Absolute Value Slope | -3.313 | 36.7025 |
| Variance of EMG | 5.24E+06 | 7.33E+06 |
| Root Mean Square | 0.01 | 0.0059 |
| Waveform Length | 3.77E+04 | 2.08E+04 |
| Zero Crossing | 32.125 | 9.4582 |
| Slope Sign Change | 58.1528 | 6.9995 |
| Willison Amplitude | 97.4556 | 0.7376 |
| Autoregressive Coefficients | 2.58E+07 | 3.64E+07 |
| Modified Median Frequency | 1.68E+03 | 1.01E+03 |
| Modified Mean Frequency | 486.8818 | 11.2808 |

The given table 2 shows the mean and standard deviation for all the features for diabetic patient.

| FEATURES | MEAN | STD |
|---|---|---|
| Integrated EMG | 288.6233 | 852.8137 |
| Mean Absolute Value | 439.7441 | 237.7973 |
| Modified Mean Absolute Value 1 | 333.7233 | 180.5012 |
| Modified Mean Absolute Value 2 | 221.8945 | 119.6891 |
| Mean Absolute Value Slope | -1.441 | 75.8474 |
| Variance of EMG | 1.65E+07 | 1.83E+07 |
| Root Mean Square | 0.00116 | 0.0081 |
| Waveform Length | 6.97E+04 | 3.72E+04 |
| Zero Crossing | 31.767 | 13.7266 |
| Slope Sign Change | 5.94E+01 | 7.83E+00 |
| Willison Amplitude | 97.196 | 0.945 |
| Autoregressive Coefficients | 8.11E+07 | 8.92E+07 |
| Modified Median Frequency | 3.07E+03 | 1.62E+03 |
| Modified Mean Frequency | 4.81E+02 | 1.08E+01 |

© 2015 Global Journals Inc. (US)

## VII. Feature Extraction Stage

In this section the feature extraction and classification steps and their outcomes are presented





Fig.2 (a) Original Signal

Fig.2 (b) Segmented Signal

Fig 2. (c) Mean Absolute Value

Fig.2 (d) Slope of Mean Absolute Value

*Figure 2 :* Segmentation & Mean calculation of Raw EMG Signal

The given figure 2 shows the EMG signal processing and feature extraction using our proposed method. Fig 2(a) is the raw signal taken from the electrodes. Fig 2(b) is the segmentation of the raw signal in samples. Fig 2(c) presents the Mean absolute value. In fig 2(d) Slope of Mean Absolute Value.

The next step is to combine all the features in to one matrix for the classification. In our method for non-diabetic user we have assigned class as 0 and for diabetic user class 1 has been assigned. The extracted features combined into one matrix and passed to the SVM classifier for the training the data. After finishing the training step, data is parsed to the prediction stage to get the classification results.

Finally, confusion matrix of classification, precision and recall, specificity and sensitivity of the proposed system are calculated.



© 2015 Global Journals Inc. (US)





*Figure 3 :* Power Spectral Density and Weighted Mean calculation

The given figure 3 shows the EMG signal Power Spectral Density and Weighted Mean calculation. Fig 3(a) is the mean of power spectral density. Fig 3(b) is the amplitude of mean which is calculated from FFT. Fig 3(c) presents the power spectral density of the median data of the Raw EMG. In fig 3(d) presents the weighted mean of absolute values which is denoted by MMAV1.



© 2015 Global Journals Inc. (US)





Fig 4 (a) Frequency Spectrum Ratio

Fig 4(b) Median Amplitude

Fig 4(c) MMAV2

Fig 4(d) Waveform Length

*Figure 4 :* Frequency Spectrum, median amplitude, mean absolute value and Waveform length calculation

The given figure 4 shows the EMG signal Frequency Spectrum, median amplitude, mean absolute value and Waveform length calculation. Fig 4(a) is the frequency ratio of the Raw EMG. Fig 4(b) is the amplitude of median which is calculated from FFT. In fig 4(c) presents the weighted mean of absolute values which is denoted by MMAV2. Fig 4(d) is the waveform length.

## VIII. Classification Stage

This is the final stage of our framework in order to achieve the classification results. In this stage all the extracted feature are given to the SVM Classifier. The first step in the classification stage is to train the dataset based on the features extracted from our feature extraction method. After training the dataset one trained model is created based on the feature. This model contains labels of the dataset, indexes and support vectors.

After creating this process of training; this model is given to the SVM prediction step for the classification of the test dataset. In the prediction model the predicted labels are achieved according to the test and train data. Finally the statistics of SVM Classifier is calculated.



© 2015 Global Journals Inc. (US)

*Table 1 :* Confusion Matrix

| | |
|---|---|
| 292 | 3 |
| 0 | 2 |

The above given table represents the confusion matrix of the classifier. According to our dataset we have 297 user's EMG signals. Out of 297, 2 user are having diabetes and remaining are non-diabetes, as we discussed earlier section that for non-diabetes, the class is assigned as 0 and diabetes is 1. It is clear from the confusion matrix that the proposed system is able to identify the diabetes class. It is showing that 292 users are non-diabetic and 2 users are diabetic patient, the remaining 3 users are the misclassification of the approach.





*Figure 5 :* Sensitivity of the proposed system

In figure 5 another result analysis parameter is plotted which is the Sensitivity of the system. According to this framework, this approach is highly sensitive for the diabetic and non-diabetic class. The x-axis represents the number of class and y-axis represents the sensitivity of the classifier w.r.t the class.

*Table 2 :* Confusion matrix statistics

| RECALL | PRECISION | SPECIFICITY |
|---|---|---|
| 0.9898 | 1 | 1 |
| 1 | 0.4 | 0.9898 |

The above given table represents the recall, precision, specificity of the proposed frame work.

$$Precision = TP / (TP + FP)$$

$$Recall = TP / (TP + FN)$$

- Sensitivity (also called the true positive rate, or the recall in some fields) measures the proportion of positives that are correctly identified.
- Specificity (also called the true negative rate) measures the proportion of negatives that are correctly identified.

According to this the precision and recall is calculated for this method.

Total number of users in the dataset is 297 and correctly classified are 294, so the final classification result in terms of accuracy is 98.98%.

## IX. Conclusion

The study described the use of the electromyography pattern recognition method, which is very important in different applications, such as rehabilitation devices, prosthetic arm/leg control, assistive technology, symptom detection for neuromuscular disorder, and so on.

In case of a disease monitoring system (i.e. diabetes), two major criteria are applicable—one is robustness and reliability, and another is accuracy of detection. Based on these criteria, the SVM classifier was trained using the extracted features. The experimental results show that the proposed approach is able to classify the diabetes with a better accuracy of 98.9899%.

## References Références Referencias

1. Nishikawa, K. and Kuribayashi, K., "Neural network application to a discrimination system for EMG controlled prostheses.", IEEE /RSJ international workshop on intelligent robots and systems, pp. 231-236 (1991).
2. Xiao Hu; Qun Yu; Waixi Liu; Jian Qin, "Feature Extraction of Surface EMG Signal Based on Wavelet Coefficient Entropy," in Bioinformatics and Biomedical Engineering, 2008. ICBBE 2008. The 2nd International Conference on, vol., no., pp. 1758-1760, 16-18 May 2008
3. kouchaki, s.; Boostani, R.; Shabani, S.; Parsaei, H., "A new feature selection method for classification of EMG signals," in Artificial Intelligence and Signal Processing (AISP), 2012 16th CSI International Symposium on , vol., no., pp.585-590, 2-3 May 2012
4. ZhaojieJu; Gaoxiang Ouyang; Wilamowska-Korsak, M.; Honghai Liu, "Surface EMG Based Hand Manipulation Identification Via Nonlinear Feature Extraction and Classification," in Sensors Journal, IEEE , vol.13, no.9, pp.3302-3311, Sept. 2013
5. Artug, N.T.; Goker, I.; Bolat, B.; Tulum, G.; Osman, O.; Baslo, M.B., "Feature extraction and classification of neuromuscular diseases using scanning EMG," in Innovations in Intelligent Systems and Applications (INISTA) Proceedings, 2014 IEEE



© 2015 Global Journals Inc. (US)

International Symposium on , vol., no., pp.262-265, 23-25 June 2014
6. Yong Ning; Xiangjun Zhu; Shanan Zhu; Yingchun Zhang, "Surface EMG Decomposition Based on K-means Clustering and Convolution Kernel Compensation," in Biomedical and Health Informatics, IEEE Journal of , vol.19, no.2, pp.471-477, March 2015
7. Sivarit Sultornsaneea, Ibrahim Zeida, Sagar Kamarthia, " Classification of Electromyogram Using Recurrence Quantification Analysis", Proceediang of Computer Science, Elsiever, vol.6, pp (375–380), 2011.



© 2015 Global Journals Inc. (US)

This page is intentionally left blank

© 2015 Global Journals Inc. (US)

Global Journal of Medical Research: K
Interdisciplinary
Volume 15 Issue 6 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

# Effect of Aerobic Training on Ventricular Remodeling After Percutaneous Coronary Intervention

By Sarah Medhat Mohammed kamel, MSc, Prof. Dr. Zahra Mohammed Hassan Serry, PhD Dr. Hany Ezzat Obaya & Prof. Dr. Osama Al Sayed Abd El Moneem, PhD

*Cairo University, Egypt*

*Abstract-* *Background:* Coronary artery disease (CAD) is one of the most common causes of morbidity and mortality in different communities worldwide, and impair the patient's quality of life (QoL). Left ventricular ejection fraction (LVEF) as a clinical index of myocardial contractility and its pumping action is a well established predictor of mortality and long term prognosis in acute myocardial infarction (AMI). Exercise training in post event CAD patients could significantly improve not only the myocardial contractility in terms of LVEF but also could be effectively and safely used with low risk and moderate risk CAD patients. The aim of this study was to determine the effect of exercise training on ventricular remodeling and QoL after percutaneous coronary intervention (PCI).

*Keywords:* *aerobic exercise, left ventricular ejection fraction, percutaneous coronary intervention, quality of life.*

*GJMR-K Classification:* *NLMC Code: WB 377*




*Strictly as per the compliance and regulations of:*

© 2015. Sarah Medhat Mohammed kamel, MSc, Prof. Dr. Zahra Mohammed Hassan Serry, PhD Dr. Hany Ezzat Obaya & Prof. Dr. Osama Al Sayed Abd El Moneem, PhD. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Effect of Aerobic Training on Ventricular Remodeling After Percutaneous Coronary Intervention

Sarah Medhat Mohammed kamel [α], MSc, Prof. Dr. Zahra Mohammed Hassan Serry [σ], PhD
Dr. Hany Ezzat Obaya [ρ] & Prof. Dr. Osama Al Sayed Abd El Moneem [ω], PhD

*Abstract- Background:* Coronary artery disease (CAD) is one of the most common causes of morbidity and mortality in different communities worldwide, and impair the patient's quality of life (QoL). Left ventricular ejection fraction (LVEF) as a clinical index of myocardial contractility and its pumping action is a well established predictor of mortality and long term prognosis in acute myocardial infarction (AMI). Exercise training in post event CAD patients could significantly improve not only the myocardial contractility in terms of LVEF but also could be effectively and safely used with low risk and moderate risk CAD patients. The aim of this study was to determine the effect of exercise training on ventricular remodeling and QoL after percutaneous coronary intervention (PCI).

*Subjects and Methods:* Sixty patients of both sexes had been recruited from National Heart Institute (NHI), Cairo, Egypt. All patients were within the first year after PCI. They were randomly assigned to 2 groups equal in numbers. Study group was 30 patients (21 men and 9 women, mean age was 52.2 ± 4.9 years) that had been received aerobic moderate intensity exercise training on bicycle ergometer for 50 minutes, 3 times/week, day after day, for 3 months, while control group was 30 patients (20 men and 10 women, mean age was 53.4 ± 4.8 years) that had been received the traditional cardiac care without any exercise training in form of routine pharmacological therapy and lifestyle education. Doppler echocardiography was used to measure LVEF, left ventricle end diastolic diameter (LVEDD) and left ventricle end systolic diameter (LVESD), and Nottingham health profile (NHP) questionnaire was used to measure differences in QoL between both groups. Both measurements were done before and after the study.

*Results:* After completion of the study, a significant increase was observed in LVEF ($P < 0.05$), without any significant changes in LVEDD and LVESD, also, improvement in QoL were observed in the study group ($P < 0.05$) when compared to control group.

*Conclusion:* Aerobic moderate intensity exercise training could improve LVEF and QoL after PCI. Aerobic exercise is a good method that improves cardiac contractility and ejection fraction (EF), and did not have adverse effects on LVEDD and LVESD nor cause severe cardiovascular complications.

*Keywords:* *aerobic exercise, left ventricular ejection fraction, percutaneous coronary intervention, quality of life.*

## I. Introduction

Coronary artery disease is one of the most common causes of morbidity and mortality in different communities worldwide, and impair the patient's QoL.[1] Various echocardiographic parameters have been shown cardiac dysfunction in CAD patients, such as left ventricular volumes and EF which are strongly related to prognosis of cardiac diseases.[2] LVEF as a clinical index of myocardial contractility and its pumping action is a well established predictor of mortality and long term prognosis in AMI.[3] Many published studies of heart failure patients underwent PCI commonly have a reduced LVEF when compared with normal.[4]

Pooled data from clinical trials have shown significant improvement in LVEF after exercise training in patients after PCI.[5] The Exercise in left ventricular dysfunction study reported that 3 months exercise training improved LVEF significantly in patients after PCI, furthermore, this result was confirmed after 3 months exercise in patients after AMI who had undergone successful PCI. LVEF during exercise was significantly improved.[5] However, exercise training is the core component of cardiac rehabilitation (CR) and secondary prevention of CAD, there is an evidence regarding the effectiveness of exercise training on LVEF in patients with CAD who received PCI, an early (within 1 month post discharge) 12 weeks structured exercise training program in post event CAD patients could significantly improve not only the myocardial contractility in terms of LVEF but also could be effectively and safely used with low risk and moderate risk CAD patients.[4]

It has been widely shown that exercise based CR in patients with AMI and PCI has several beneficial effects on cardiovascular functional capacity, QoL, risk factors modification, psychological profile and mortality.[6]

*Author α: Physiotherapist, MSc, National Heart Institute, Cairo, Egypt.*
*e-mail: ali311pt@yahoo.com*
*Author σ: Assistant Professor of Physical Therapy, Department of Cardiopulmonary/Respiratory Disorder and Geriatrics, Faculty of Physical Therapy, Cairo University, Egypt.*
*Author ρ: Lecturer of Physical Therapy, Department of Cardiopulmonary/Respiratory Disorder and Geriatrics, Faculty of Physical Therapy, Cairo University, Egypt.*
*Author ω: Consultant of Cardiothoracic Surgery, National Heart Institute, Cairo, Egypt.*



© 2015 Global Journals Inc. (US)

The degree of ventricular remodelling was regarded as an important prognostic factor associated with cardiac function after AMI, and an increasing number of studies have shown that, in patients with AMI with left ventricular dysfunction, exercise training did not worse ventricular remodelling, and may even prevent this spontaneous deterioration.[7]

In cardiac patients, QoL was evaluated on the basis of objective clinical criteria such as mortality, morbidity, angina, complication rates, test results, or simple indicators such as return to work or repeated hospitalizations.[8] This approach ignored the fact that cardiac disease is a life threatening disease and the operation in itself leads, regardless of the result, to changes in patients' psychosocial and social functioning.[8] It also must be realized that QoL is not only an outcome indicator but may itself be a factor that affects health, illness and coping.[8] Poor QoL has been associated with poorer outcomes, such as lower survival rates, increases in the number of hospitalizations, decreased capacity to perform activities of daily living, and decreased compliance with treatments in other populations like cardiac patients.[9] The most commonly sensitive generic instruments in heart disease are the NHP (used in approximately 40% of studies).[10] The NHP is a well validated test, which has been used previously for assessment of QoL before and after cardiac events.[10]

## II. Patients and Methods

This study was conducted in physiotherapy department of NHI, Cairo, Egypt. 60 Patients of both sexes, their ages were 45-60 years old, within the first year after PCI, their BMI was 30- 34.9 Kg/m$^2$ (class I obesity), were selected and assigned to 2 equal groups in numbers. The study group (30 patients, 21 men and 9 women) that had been received aerobic moderate intensity exercise training on bicycle ergometer 3times/week for 3 months, while the control group (30 patients, 20 men and 10 women) that had been received the traditional cardiac care without any exercise training in form of routine pharmacological therapy and lifestyle education.

Exclusion criteria were patients with Post-infarction, residual myocardial ischaemia, severe ventricular arrhythmias, atrioventricular block, hypertrophic cardiomyopathy, valvular disease requiring surgery, pericarditis, acute systemic illness or fever, severe renal dysfunction (i.e. Creatinine > 2.5 mg/dl), severe orthopaedic problems, such as osteoarthritis of both knees, other metabolic problems, such as diabetes mellitus, acute thyroiditis, hypokalaemia, hyperkalaemia and hypovolaemia.

Before starting the study, a meeting was done for all patients to record demographic data, Doppler echocardiography was used to measure LVEF, LVEDD, LVESD, and NHP questionnaire was used to measure QoL presented with each patient. In that stage a face to face instructions and administration of NHP questionnaire to all participants was given, and if the patient was not sure whether to say "yes" or "no" to a problem, they were instructed to answer the one more true at that time. All patients were taking their prescribed medications by their cardiologists.

Participants in the CR program were requested to attend their exercise program 3 times/week, day after day, for a period of 3 months. Moderate intensity aerobic exercise is prescribed based on Borg's rating of perceived exertion (BRPE) scale. The scale is comprised of 15 points where a rating of 6 means no exertion and a rating of 20 means maximal exertion.[11] Patients are encouraged to achieve a rating between 11 (fairly light) and 14 (hard), with training heart rate (60-70% of maximal heart rate). Target heart rate using karvonen method, taking into account the resting heart rate, is calculated as follows: {[(220-age) or (210-age)-resting heart rate]x(60-70)}+ resting heart rate.[12] For participants in the CR program involved in this study, each exercise session is comprised of 5 to 10 minutes warming up and 5 to 10 minutes cooling down, with active phase of approximately 30 minutes of aerobic exercise training. Aerobic exercise was the dominant mode of exercise which implemented using a bicycle ergometer in the CR program. After 3 months, LVEF, LVEDD, LVESD were measured, and also NHP questionnaire were applied again.

Data were analyzed with SPSS software version 17. The level of significance was set at $P < 0.05$. Paired t-test was applied for each group to compare pre and post values within the same group. Unpaired t-test was applied to compare pre and post values between both groups of the study.

## III. Results

Base line measurements had shown no statistical significant differences between both groups ($P>0.05$). The baseline and final values of each group (Table 1) had shown highly significant increase in LVEF ($P<0.000$) in the study group but, LVEDD and LVESD did not change significantly ($P>0.05$). The control group did not show any significant changes in LVEF, LVEDD, and LVESD ($P>0.05$). Results of LVEF were improved significantly in the study group when compared with the control group after the program ($P<0.002$).

As shown in (table 2) the six domains of NHP questionnaire of the study group had improved highly significantly ($P<0.000$). The control group did show significant improves ($P>0.05$). All domains of NHP questionnaire were improved significantly in the study group when compared with the control group after the program ($P<0.05$).



© 2015 Global Journals Inc. (US)

*Table (1) :* Changes of Echo-parameters from baseline to the end of the study within each group and between groups:

| Variables | Study group | | | Control group | | | P value for both groups after program |
|---|---|---|---|---|---|---|---|
| | Pre program | Post program | P Value | Pre program | Post program | P Value | |
| | Mean ±SD | Mean ±SD | | Mean ±SD | Mean ±SD | | |
| LVEF (%) | 56.6±6.8 | 59.6±6.9 | 0.000* | 54.1±5.9 | 55.9±6.6 | 0.3 | 0.002* |
| LVEDD(cm) | 5.01±0.32 | 5.02±0.31 | 0.89 | 5.10±0.20 | 5.13±0.21 | 0.6 | 0.13 |
| LVESD (cm) | 3.49±0.43 | 3.44±0.42 | 0.13 | 3.55±0.47 | 3.47±0.43 | 0.15 | 0.71 |

*SD: standard deviation, LVEF: Left ventricular ejection fraction, LVEDD: Left ventricle end-diastolic diameter, LVESD: Left ventricle end-systolic diameter. Significant level: P<0.05*.*

*Table (2) :* Changes of 6 domains of NHP from baseline to the end of the study within each group and between groups:

| Variables | Study group | | | Control group | | | P value for both groups after program |
|---|---|---|---|---|---|---|---|
| | Pre program | Post program | P Value | Pre program | Post program | P Value | |
| | Mean ±SD | Mean ±SD | | Mean ±SD | Mean ±SD | | |
| Energy level | 51.44±16.06 | 34.24±15.02 | 0.000* | 47.94±15.99 | 44.77±15.23 | 0.44 | 0.009* |
| Pain | 40.40±21.50 | 22.17±12.17 | 0.000* | 38.42±23.44 | 31.95±18.09 | 0.22 | 0.02* |
| Emotional reaction | 26.01±13.26 | 13.75±7.22 | 0.000* | 23.67±13.97 | 21.07±13.78 | 0.48 | 0.02* |
| Sleep | 37.22±13.98 | 22.30±8.46 | 0.000* | 33.51±17.07 | 28.96±13.00 | 0.25 | 0.02* |
| Social isolation | 37.04±12.10 | 24.15±9.07 | 0.000* | 34.48±11.76 | 31.25±11.55 | 0.31 | 0.01* |
| Physical ability | 35.29±15.56 | 20.23±11.55 | 0.001* | 30.64±16.21 | 27.58±13.53 | 0.41 | 0.03* |

*SD=Standard Deviation, Significant level: P<0.05*.*

## IV. Discussion

In this study, the LVEF significantly increased from 56.6±6.8 % at the beginning of the study to 59.6±6.9 % after 3 months of CR program (P<0.05), while no significant changes were observed in the LVEDD and LVESD when compared between those at the onset and those at three months after rehabilitation.

Results of the study were supported by Mohammad et al. baseline LVEF in the study group was 46.9 ± 5.9 and in the control group was 47.9 ± 7.0, there was a significant improvement in LVEF after 12 weeks of exercise training in the study group (46.9 ± 5.9 to 61.5 ± 5.3) compared with the control (47.9 ± 7.0 to 47.6± 6.9) group (P=0.001), Also, he concluded that a structured individually tailored home based training program could be as effective as center based programs and safely used not only in low risk but also in moderate risk CAD patients.[4] Agreed with these results, Huan et al. had showed that 6 months exercise training increased LVEF in study group more than controls, and LVESD and LVEDD decreased in the exercise group but increased in the control group, which suggested that, to a certain extent, exercise could prevent ventricular remodeling in patients after AMI.[6]

These findings are in consistent with results from Haddadzadeh et al. that found similar effects with exercise training program in post event CAD patients that significantly improved the myocardial contractility in terms of LVEF.[13] Going with the same line Masoumeh et al. reported that CR had positive effects on patients with EF of about less than 50% that was improved significantly after CR program, Moreover, peak exercise capacity was significantly improved, Also, LVESD and LVEDD had no clinical or statistical change after the program.[2] The results show that, among the patients with LV dysfunction, exercise based rehabilitation is beneficial and has no detrimental effects on ventricular remodeling.[2] Agreed with this study results, Soleimannejad et al. demonstrated that LVEF improved significantly after PCI (with or without the CR exercise program), however, the effect of the CR exercise



© 2015 Global Journals Inc. (US)

program on chamber diameters, i.e. LVEDD, LVESD were neutral.[14]

Agreed with these results, Sherin et al. in NHI in Egypt, but in dilated cardiomyopathy patients, there was high statistical significant increase in peak VO2, EF, diastolic dysfunction, resting and maximal heart rates after intervention only in the training group.[15] There was no significant change in any parameter within the control group, as for comparison between both groups; there was high significant difference in peak VO2, resting heart rate and EF after intervention, and the number of patients in the training group with normal diastolic pattern was zero before training, while it was 8 (53.3%) after training.[15] Supporting this study results Giallauria et al. showed that 6-month exercise based CR induced a combined reverse left atrial and LV remodeling as well as significant improvement in exercise functional capacity, LVEF, and early LV diastolic filling.[16]

In contrast to this study results Chul et al. reported that, in a follow up observation of 70 patients diagnosed with AMI from the fourth day of onset for three years, 14 (20%) showed widening in diameter of the left ventricle, a phenomenon that may ultimately cause severe left ventricular failure.[17] Also, they reported that 12 week CR exercise on 13 patients with AMI of around 16 weeks after onset showed reduction in motility of the myocardial wall and LVEF in echocardiography.[17] Also, Kubo et al. investigated the effects of 3 months exercise training on ventricular remodeling after extensive anterior AMI with LVEF $<$ 45% and found that control group patients' LV diastolic volume index and LV systolic volume index improved, but there was no change in the rehabilitation group.[18] Conversely, Otsuka et al. reported that early exercise training did not deteriorate ventricular remodeling in mild, moderate and severe left ventricular dysfunction in his patients.[19] In a similar study by Jiang et al. they reported that LV diastolic diameter increased in the control group, but not in the exercise group, after 3 months' exercise training.[5]

The results obtained in the present study revealed a statistical significant differences in six domains of NHP variables (pain, physical mobility, emotional reactions, energy, social isolation and sleep) between control and study groups ($p < 0.05$) that reflected more improvement of QoL for the study group after CR program. Supporting these results, Marzieh et al. reported that after CR, scores of all physical domains of the short form-36 questionnaire (SF-36) including physical function, physical limitation, body pain, vitality in addition to general health were significantly improved in all patients compared to the baseline and showed that physical activity had influenced QoL, so that increasing physical activity improves QoL.[20] Going with the same line, Arrigo et al. had shown that a comprehensive CR improves QoL even one year after the program.[21]

Supporting these results Babaee et al. had shown significant difference in QoL (by SF-36 and NHP) between study and control group, significant improvements in QoL between two groups, as measured by the NHP, were seen in energy, pain, emotional reaction, sleep, physical mobility and total average quality of life.[22] Significant improvements in QoL between two groups, as measured by the SF-36, were seen in physical function, role limitations resulting from emotional status, role limitations resulting from physical status, mental health, vitality and total average QoL, and demonstrated that health education resulted in improved QoL for patients with CABG.[22] In consistent with the study results, Yohannes et al. investigated the long term effects of a 6 week CR on physical activity, psychological well being, and QoL in 147 cardiac patients. The results demonstrated the benefits of CR in improving HRQoL and physical activity, and in reducing anxiety and depression.[23] Furthermore, these benefits were maintained at 12 month follow up.[23]

However, there were some investigations with different findings; in Serber et al. study, the impact of CR on patients with severe psychological distress was more than others in physical, mental and social aspects of QoL, and showed that QoL was related to primary level of psychological distress of the patients and CR could improve QoL and anxiety just in these group of patients.[24] In contrast Worcester et al. suggested that CR is not sufficiently intensive to influence recovery of QoL.[25] Briffa et al. reported that CR was only found to affect physical function in a recent randomized controlled trial of an 18 session program which compared CR with usual care, but this may be due to short period of his program.[26]

There is a significant and positive relationship between CR and changes in QoL domains. Patients started the study with bad QoL scores and demonstrated significant improvement in QoL scores following the completion of the programme. Likewise, the contractility of the heart increased significantly, physical abilities had increased, they reported feeling less pain and were less limited in activities they did before. According to the reports of the investigators in fields related to the present study, it can be concluded that, all participants's QoL scores and LVEF had improved after the intervention. Exercise based rehabilitation is beneficial to improve cardiac contractility and has no detrimental effects on ventricular remodeling.

## V. Conclusion

It was concluded that aerobic training has a positive effects on improving LVEF in post PCI patients, also QoL domains of NHP questionnaire as pain, physical mobility, emotional reactions, energy, social isolation and sleep were improved, further more CR is a



© 2015 Global Journals Inc. (US)

good method that improve cardiac contractility and ejection fraction, and did not have adverse effects on LVEDD and LVESD nor cause severe cardiovascular complications.

## References Références Referencias


1. Hadaegh F., Harati H., Ghanbarian A. and Azizi F. Prevalence of coronary heart disease among Tehran adults: Tehran Lipid and Glucose Study. East Mediterr Health J 2009; 15: 157-166.
2. Masoumeh S., Mohammad G. Mahboobeh T., Nizal S., Hamidreza R., and Mohsen K. The Impacts of Cardiac Rehabilitation Program on Echocardiographic Parameters in Coronary Artery Disease Patients with Left Ventricular Dysfunction. Cardiology Research and Practice 2013; article ID 201713, 4 pages.
3. Dutcher J., Kahn J., Grines C. and Franklin B. Comparison of left ventricular ejection fraction and exercise capacity as predictors of two and five-year mortality following acute myocardial infarction. Am J Cardiol 2007; 99: 436- 441.
4. Mohammad H., Arun M., Padmakumar C., Bijan S. and Fardin M. Effect of Exercise Based Cardiac Rehabilitation on Ejection Fraction in Coronary Artery Disease Patients: A Randomized Controlled Trial. heart views 2011; 12: 51-57.
5. Jiang A., Zhang F., Gao W., Li Z., Zhao W. and Li X. et al. The impact of exercise rehabilitation on left ventricular remodeling and systolic function in acute myocardial infarction patients. Chin J Intern Med 2006; 45 (11): 904-906.
6. Huan Z., Ming L., Yi S., Yuan M. and Wenhui K. effects of 6 months exercise training on ventricular Remodelling and autonomic tone in patients with Acute myocardial infarction and percutaneou s coronary intervention. J Rehabil Med 2008; 40: 776-779.
7. Van T., Huijsmans R., Kroon D., Schothorst M. and Kwakkel G. Effects of exercise training on cardiac performance, exercise capacity and quality of life in patients with heart failure: a meta analysis. Eur J Heart Fail 2006; 8: 841-850.
8. Rumsfeld J., Michael P. and Magid D. et al. Predictors of Health-Related Quality of Life after Coronary Artery Bypass Surgery. Ann Thorac Surg 2004; 77: 1508-1513.
9. Havik O., Sivertsen B. and Relbo A. et al. Depressive symptoms and all-cause mortality after heart transplantation. Transplantation 2007; 84: 97-103.
10. Mcauley E. and Morris K. Advances in Physical Activity and Mental Health: Quality of Life. American journal of lifestyle medicine 2007; 1: 389- 396.
11. Borg G. and Linderholm H. Perceived exertion and pulse rate during graded exercise in various age groups. Acta Medica Scandinavia 1967; 181(472): 194-206.
12. Kennedy M., Haykowsky M., Daub B., Lohuizen K., Knapik G. and Black B. Effects of a comprehensive cardiac rehabilitation program on quality of life and exercise tolerance in women. Curr Controlled Trials Cardiovasc Med 2003; 4(1): 1. doi: 10.1186/1468-6708-4-1
13. Haddadzadeh M., Maiya A., Padmakumar R., Devasia T., Kansal N. and Borkar S. Effectiveness of cardiac rehabilitation on myocardial contractility in post-event coronary artery disease patients: A randomized controlled trial. Physiotherapy 2010; 8: 5-12.
14. Soleimannejad K., Nouzari Y., Ahsani A., Nejatian M. and Sayehmiri K. Evaluation of the Effect of Cardiac Rehabilitation on Left Ventricular Diastolic and Systolic Function and Cardiac Chamber Size in Patients Undergoing Percutaneous Coronary Intervention. J Teh Univ Heart Ctr 2014; 9(2):54-58.
15. Sherin H. Correlation between changes in diastolic dysfunction and health-related quality of life after cardiac rehabilitation program in dilated cardiomyopathy, Journal of Advanced Research 2013; 4: 189-200.
16. Giallauria F., Galizia G. and Lucci R. et al. Favourable effects of exercise-based cardiac rehabilitation after acute myocardial infarction on left atrial remodeling. International Journal of Cardiology 2009; 136 no. 3: 300-306.
17. Chul K., Duk Y. and Dong W. The Impact of Early Regular Cardiac Rehabilitation Program on Myocardial Function after Acute Myocardial Infarction. Ann Rehabil Med 2011; 35: 535-540.
18. Kubo N., Ohmura N., Nakada I., Yasu T., Katsuki T. and Fujii M. et al. Exercise at ventilatory threshold aggravates left ventricular remodeling in patients with extensive anterior acute myocardial infarction. Am Heart J 2004; 147: 113-120.
19. Otsuka Y., Takaki H., Okano Y., Satoh T., Aihara N. and Matsumoto T. et al. Exercise training without ventricular remodelling in patients with moderate to severe LV dysfunction early after acute myocardial infarction. Int J Cardiol 2003; 87: 237-244.
20. Marzieh S., Samaneh M., Hosein H. and Sepehr M. Effects of a comprehensive cardiac rehabilitation program on quality of life in patients with coronary artery disease, arya atheroscler 2013; 9: 179-185.
21. Arrigo I., Brunner H. and Lefkovits M. Comparative outcome one year after formal cardiac rehabilitation: the effects of a randomized intervention to improve exercise adherence. Eur J Cardiovasc Prev Rehabil 2008; 15: 306-311.
22. Babaee G., Keshavarz M., Hidarnia A. and Shayegan M. Evaluation of quality of life in patients with coronary artery bypass surgery using controlled clinical trial. Acta Medica Iranica 2007; 45: 69-76.




© 2015 Global Journals Inc. (US)

23. Yohannes A., Doherty P. and Bundy C. et al. The long-term benefits of cardiac rehabilitation on depression, anxiety, physical activity and quality of life. J Clin Nurs 2010; 19: 2806-2813.
24. Serber E., Todaro J. and Tilkemeier P. et al. Prevalence and characteristics of multiple psychiatric disorders in cardiac rehabilitation patients. J Cardiopulm Rehabil Prev 2009; 29: 161-168.
25. Worcester M., Murphy B. and Elliott P. et al. Trajectories of recovery of quality of life in women after an acute cardiac event. British Journal of Health Psychology 2007; 12: 1-15.
26. Briffa T., Eckermann S. and Griffiths A. et al. Cost-effectiveness of rehabilitation after an acute coronary event: a randomized controlled trial. Med J Aust 2005; 183: 450-455.



© 2015 Global Journals Inc. (US)

Global Journal of Medical Research: K
Interdisciplinary
Volume 15 Issue 6 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

# The Impact of Low Frequency Ultrasound and Lymphatic Drainage on Triglycerides

By Manar M. Badawy, Nesreen G. Elnahas, Zahra M. H Serry & Basem Alzareef Fouad

*Dammam University; Saudi Arabia & Cairo University; Faculty of physical therapy; Egypt*

*Abstract-* The aim of this study is to evaluate the effect of low frequency ultrasound plus lymphatic drainage on blood triglycerides in cardiac patients (chronic coronary atherosclerosis patients with high triglycerides and fat mass body composition. Forty female patients with age ranges from 40 to 50 years were selected from Palestine hospital and they were chronic atherosclerotic patients and were assigned into 2 groups according to their BMI based on the classification of the world health organization. Each patient in the two groups (Group A and Group B) was evaluated before and after 24 sessions treatment program by using the combination of ultrasound and lymphatic drainage machine. The assessment of blood serum triglycerides by UDICHEM-310 ANALYSER have been done before and after the end of 24 sessions and Re-assessment after 2 months from the last treatment session.

*Keywords: low frequency ultrasound; lymphatic system; lymphatic drainage; triglycerides; coronary atherosc-lerosis.*

*GJMR-K Classification: NLMC Code: WN 180, WH 700*




*Strictly as per the compliance and regulations of:*

© 2015. Manar M. Badawy, Nesreen G. Elnahas, Zahra M. H Serry & Basem Alzareef Fouad. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# The Impact of Low Frequency Ultrasound and Lymphatic Drainage on Triglycerides

Manar M. Badawy [α], Nesreen G. Elnahas [σ], Zahra M. H Serry [ρ] & Basem Alzareef Fouad [ω]

*Abstract-* The aim of this study is to evaluate the effect of low frequency ultrasound plus lymphatic drainage on blood triglycerides in cardiac patients (chronic coronary atherosclerosis patients with high triglycerides and fat mass body composition. Forty female patients with age ranges from 40 to 50 years were selected from Palestine hospital and they were chronic atherosclerotic patients and were assigned into 2 groups according to their BMI based on the classification of the world health organization. Each patient in the two groups (Group A and Group B) was evaluated before and after 24 sessions treatment program by using the combination of ultrasound and lymphatic drainage machine. The assessment of blood serum triglycerides by UDICHEM-310 ANALYSER have been done before and after the end of 24 sessions and Re-assessment after 2 months from the last treatment session. The collected raw data of the current patients were statistically analyzed to evaluate the results of the two groups to investigate the effect of using the combination of ultrasound and lymphatic drainage machine on blood serum triglycerides. In this study, the results are revealed statistically significant improvement of blood serum triglycerides before and after the treatment with more improvement had been achieved after 2 months after last session. Furthermore, the low frequency ultrasound technique plus lymphatic drainage technique improve blood serum triglycerides of chronic coronary atherosclerosis patients with high triglycerides and fat mass composition.

*Keywords: low frequency ultrasound; lymphatic system; lymphatic drainage; triglycerides; coronary atherosclerosis.*

## I. Introduction

Triglyceride (TG, triacylglycerol, TAG or triacylglyceride) is an ester derived from glycerol and three fatty acids. Triglycerides are a blood lipid that helps enable the bidirectional transference of adipose fat and blood glucose from the liver. There are many triglycerides depending on the oil source. Some of these are highly unsaturated in the human body; high levels of triglycerides in the bloodstream have been linked to atherosclerosis and by extension, the risk of heart disease and stroke [1].

Hypertriglyceridemia is a prevalent risk factor for cardiovascular disease (CVD) and increasingly important in the setting of current obesity and insulin resistance epidemics. High triglyceride (TG) levels are markers for several types of atherogenic lipoproteins. Patients who have hypertriglyceridemia may be at significant risk for CVD even if low-density lipoprotein cholesterol levels are at goal, and therefore warrant treatment that optimizes diet, reduces overweight, and promotes regular exercise [2].

High-risk patients with hypertriglyceridemia, such as those with diabetes, CVD, or metabolic syndrome, may benefit from additional drug treatment aside from a statin to address other lipid abnormalities. In this discussion, we review the role of hypertriglyceridemia and its associated atherogenic lipoproteins in the pathogenesis of atherosclerosis, the relevance of a high TG level as a predictor of CVD, the cardiovascular outcomes from TG-lowering intervention trials, and the current guidelines for treating hypertriglyceridemia [3].

Men and women who have high triglyceride levels >150 mg/dl and a low level of HDL cholesterol <40 mg/dL are characterized by a significantly increased cardiovascular risk. The high triglyceride/low HDL cholesterol phenotype is a hallmark of the metabolic syndrome. The metabolic syndrome is closely associated with insulin resistance and is highly associated with the risk of CHD. It has a greater impact on the incidence of CHD in women than in men [3].

Adipose cells which make up adipose tissue are specialized cells which contain and can synthesize globules of fat. This fat either comes from the dietary fat we eat or is made by the body from surplus carbohydrate or protein in our diet. Adipose tissue is mainly located just under the skin, although adipose deposits are also found between the muscles, in the abdomen, and around the heart and other organs. The location of fat deposits is largely determined by genetic inheritance. Thus it is not possible to affect where we store fat. Nor is it possible to influence from which area the body burns fat for energy purposes [4].

*Author α: Department of Physical Therapy, Faculty of Applied Medical Health, University of Dammam, Dammam.*
*e-mail: mmbadawy@uod.edu.sa*
*Author σ ρ ω: Faculty of Physical Therapy, Cardiovascular/ Respiratory Disorder and Geriatrics Department, Cairo University.*



© 2015 Global Journals Inc. (US)





*Fig. (1) :* Simple composition of adipose tissue

Respective of the location from which they are obtained, the fat cells in humans are composed almost entirely of pure triglycerides with an average density of about 0.9 kilograms per liter. Most modern body composition laboratories today use the value of 1.1 kilograms per liter for the density of the "fat free mass", a theoretical tissue composed of 72% water (density = 0.993), 21% protein (density = 1.340) and 7% mineral (density = 3.000) by weight [5].

Fat cells are not only energy depots, but are busy endocrine organs. They secrete cytokines, which regulate responses to infection, immune reactions, inflammation and trauma. In regards to inflammation regulation, fat cells secrete pro-inflammatory (TNF, IL-6, and C-reacting protein "C-RP") and anti-inflammatory (adiponectin) cytokines. Unfortunately, with visceral fat obesity accumulation, adiponectin levels are reduced, thus leading to a higher cardio metabolic disorders (e.g., heart disease and diabetes [3].





*Fig. (2) :* Adipocyte tissue

High fat masses especially abdominal obesity is associated with an accelerated atherosclerosis, atherogenic blood triglycerides and other compartments of lipid profile, elevated platelet counts in females with chronic inflammation and increased rates of cardiovascular death [6,7].





*Fig. (3) :* Abnormal adipose tissue and atherosclerosis [6,7].



© 2015 Global Journals Inc. (US)

Low intensity low frequency (LILFU) stands for low intensity, low frequency ultrasound. It is a new technique devised by the team of William Tyler from Arizona State University to manipulate neural circuits using ultrasounds. This could make in the future the need of intervention (surgical) neuromodulation unnecessary [8].





*Fig. (4) :* The relation between ultrasound frequency and penetration

Cavitation is the process in which a bubble in a fluid is forced to oscillate in size or shape due to some form of energy input, such as an acoustic field. Such cavitation is often employed in ultrasonic cleaning baths and can also be observed in pumps, propellers [8].

Lymph carries away large particles bacteria, Cell debris which can then be filtered out and destroyed by the lymph node Lymph capillaries in the interstitial spaces have same structure as blood capillaries but their walls are more permeable to inter stitial fluid constituents [9].





*Fig. (5) :* Diffusion of nutrients and waste products between capillaries and cells [9].

While lymph is in many instances, a fluid rich in emulsified lipids, it is interesting to note that fat deposition has a tendency to develop first along lymphatic structures, as it very clearly occurs around mesenteric lymphatics and lymph nodes (Figure (1)). Moreover, nearly all lymph nodes are embedded in adipose tissue and most peripheral adipose depots contain one or more lymph nodes as well.44. However, whether there is a causal relationship between lipid content in the lymph and fat deposition is not known [10].

What seems to become clear, however, is that lymph stasis and ⁄ or fluid leakage from lymphatic vessels may promote fat accumulation.45 In our experience, when intestinal lymphatic drainage was interrupted by heat cauterization of guinea pig mesenteric vessels, an obvious increase in mesenteric adipose tissue deposition occurred over a 28-day period.46. This observation is reminiscent of what occurs during lymphedema, where lymphatic drainage is disrupted or malfunctioning [11].

The fatty acids found within lymphocytes of lymph nodes upon stimulation come mainly from triacylglycerol's in the immediately adjacent perinodal adipose tissue, which contains more polyunsaturated fatty acids than fat further from lymph nodes [10].

The relationships between the lymphatic system, adipose tissue, lipids profile and immune response ⁄inflammation are undeniable. Our current understanding of the cross-talk between these systems and illustrates how disturbance of these interactions may contribute to the pathogenesis of many disease [11].



© 2015 Global Journals Inc. (US)





*Fig. (6) :* Proposed working model for the interactions between lymphatic, fat and inflammation [11].

As the lymphatic system is the platform for the immune system and lymph nodes are the sites of adaptive immune responses modulated by the surrounding adipose tissue, increased knowledge of how the lymphatic system contributes to triglycerides transport, distribution and metabolism and to the pathogenesis of chronic inflammatory conditions may provide the basis for the development of new therapeutic strategies and increased quality of life [11].

## II. Methods and Procedures

*a) Participants*

The study was carried on 40 female patients with age ranges from 40 to 50 years were selected from Palestine hospital, They were obese atherosclerotic patients and were assigned into 2 groups according to their BMI based on the classification of the world health organization, Each patient in the two groups (class I and class II) was evaluated before and after 24 sessions treatment program by using the combination of ultrasound and lymphatic drainage machine. The assessment of blood serum triglycerides by UDICHEM-300 Chemistry Analyzer have been done before and after the end of 24 sessions and Re-assessment after 2 months from the last treatment session.

*Group (A):* Class II obesity 20 patients with BMI ranged between (35 – 39.9) kg/m$^2$

*Group (B):* Class III obesity 20 patients with BMI ranged between (≥ 40) kg/m$^2$

All patients signed a consent sheet before participation in the study.

*b) Procedures*

i. *Evaluation equipment's and Procedure*

- Weight, height are measured by using weight, height scale
- Measurements were performed while the patients standing erect back and knees extended and both upper limb beside the body

*Measurement of fasting Blood sample:* Tourniquet, disposable plastic syringes to draw venous blood samples from antecubital vein in to polypropylene tubes, and then blood glucose and blood lipid profile were assessed through laboratory investigations, figure.



*Fig. (7) :* UDICHEM-300 Chemistry Analyzer



© 2015 Global Journals Inc. (US)

- Blood sample will be collected and investigated in the laboratory before and immediately after the treatment course and re-estimated two months late.

A three milliliter-sample of venous blood was drawn from the antecubital vein after 12- 14 hours fast from all patients before the initiation of the training program and after the completion of the study (i.e. at the end of the 24 sessions) and re-assessed 2 months after the end of the program to be assayed for measurement of serum level of fasting Triglycerides.



*Fig. (8) :* Polypropylene tubes showing its gel contents

ii. *Procedure of the Treatment*

*Preparation of the patient:*

- The purpose and the procedures of the study were thoroughly explained to the patient.
- Assessment of weight, height and serum triglycerides before and immediately after the treatment course and re-estimated two months later.
- A-Low Frequency Ultrasound Application
  - General application of the ultra sound including abdominal area, upper arms, thigh and hips from supine position , Scapular area , posterior aspect of the hips from prone position
  - Duration of the treatment was 20 minutes. The pulse duration was 200 msec. Freq. low frequency 28-31 KHz low intensity. Stimulation mode (20sec. stimulation, 20sec pause).
  - 22 rectangular electrodes (60 x 130)mm. Lip sound machine VLS3. Good stabilization of electrodes was obtained throughout the session. Usage of ultrasound gel for all electrodes.



*Fig. (9) :* Low Frequency Ultrasound Machine



*Fig. (10) :* Low Frequency Ultrasound Machine Electrodes



© 2015 Global Journals Inc. (US)

*Vacuum Pressure Machine for lymphatic drainage*

Vacuum pressure was applied for 20 minute (Starvac vacuum massage 866-312-7540 low power consumption 200W) cyclical vacuum starting from 80 mill bars, (roller methods). General application of the ultra sound including abdominal area, upper arms, thigh and hips from supine. Scapular area, posterior aspect of the hips from prone position. Duration of the treatment was 20 minutes.

Treatment session was applied 3 times a week for two month and each session included lipo sound lymphatic drainage technique.



*Fig. (11) :* Vacuum massage machine



*Fig. (12) :* Lymphatic drainage technique

## III. Main Results

*a) General Characteristics*

The 40 females participated in this study were classified into two groups of equal number:

*Class I:* include twenty female patients with age ranges from 40 to 50 years old with the mean value of (46 ± 3.77) yrs, the weight ranged from 80 to 100 Kg with the mean value of (98.5 ± 6.67) kg, the height ranged from 150 to 160 cm with the mean value of (158 ± 5.06) cm. The BMI ranged from 35 to 39.9 kg/m2 with the mean value of (38. 3 ± 1.98) kg/m2 as shown in Table (1) and Figure (1).

*Class II:* include twenty female patients with age ranges from 40 to 50 years old with the mean value of (45 ± 3.01) yrs, the weight ranged from 110 to 120 Kg with the mean value of (110.5 ± 5.55) kg, the height ranged from 155 to 165 cm with the mean value of (155 ± 4.44) cm. The BMI ranged from > 40 kg/m2 with the mean value of (44.5 ± 3.22) kg/m2 as shown in Table (3) and Figure (44).



© 2015 Global Journals Inc. (US)

*Table (1) :* The mean values within the two groups.

| Variable | Groups | Mean ± SD | median | Max - Min | t-value | p-value |
|---|---|---|---|---|---|---|
| Age (years) | Class I | 46 ± 3.77 | 46 | 40 – 50 | 0.56 | 0.889 NS |
| | Class II | 45 ± 3.01 | 45.5 | 40 - 50 | | |
| Weight (Kg) | Class I | 98.5 ± 6.67 | 90 | 80 - 100 | 1.011 | 0.765 NS |
| | Class II | 110.5 ± 5.55 | 114 | 110 - 120 | | |
| Height (cm) | Class I | 158 ± 5.06 | 155 | 150 – 160 | 1.209 | 0.433 NS |
| | Class II | 155 ± 4.44 | 160 | 155 – 165 | | |
| BMI (kg/m²) | Class I | 38. 3 ± 1.98 | 37 | 35 – 39.9 | 0.97 | 0.766 NS |
| | Class II | 44.5± 3.22 | 47 | > 40 | | |





*Fig. (13) :* The mean values within the two groups

*b) Total Body Triglycerides*

To determine the differences in the mean values of total body triglycerides among the groups, repeated measure analysis of variance (AVOVA F-Test) was performed as shown in Table (1) showed the f-test result.

*Table (2) :* Comparing the mean values of total body triglycerides among the two groups.

| | Total Body Triglycerides | Mean ± SD | ANOVA Table | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | Sum of Squares | DF | Mean Square | F | Sig |
| Class I | Pre treatment | 267.78 ± 34.34 | Between Groups | 8767.492 | 2 | 4383.746 | 3.973 | .024 |
| | Post treatment | 245.01 ± 32.79 | Within Groups | 62900.358 | 57 | 1103.515 | | |
| | Post II treatment | 240 ± 32.49 | Total | 71667.850 | 59 | | | |
| | | | | | | | | |
| | | | | Sum of Squares | DF | Mean Square | F | Sig |
| Class II | Pre treatment | 300.29 ± 66.12 | Between Groups | 15220.133 | 2 | 7610.0665 | 3.443 | .038 |
| | Post treatment | 268.39 ± 57.22 | Within Groups | 125999.91 | 57 | 2210.5248 | | |
| | Post II treatment | 264.89 ± 61.15 | Total | 141220.05 | 59 | | | |



© 2015 Global Journals Inc. (US)





*Fig. (14) :* Comparing the mean values of Total Body Triglycerides among the two groups

The results of the class I and class II groups revealed that, there is a significant improvement (reduction) in the Total Body Triglycerides for the subjects at the three stages of the measurements (Pre, Post, and Post II). Accordingly to ANOVA F-test should be followed by the Post-Hoc LSD method to identify which group has the significant differences.

In the following table, class I results shows that Pre-treatment is significantly different from Post treatment and Post II.

*Table (3) :* The mean value of the three measurement of Total body Triglycerides

| Variable | Mean Difference | Significance |
|---|---|---|
| Total Body Triglycerides | | |
| Pre, Post | 22.765* | (0.034) S |
| Pre, Post II | 22.78* | (0.011) S |
| Post, Post II | 5.015 | (0.635) NS |

In the following table, Class II results shows that Pre-treatment is significantly different from Post II treatment only.

*Table (4) :* The mean value of the three measurement of Total body Triglycerides

| Variable | Mean Difference | Significance |
|---|---|---|
| Total Body Triglycerides | | |
| Pre, Post | 31.9 | (0.098) NS |
| Pre, Post II | 25.049 | (0.029) S |
| Post, Post II | 3.5 | (0.854) NS |

## IV. Discussion

This study was conducted to assess the responses of serum Triglycerides to a treatment program by using the combination of ultrasound and lymphatic drainage machine in chronic atherosclerotic patients, The results showed that there were significant difference in Serum Triglycerides that is strongly correlated with the reduction of the body fat mass studied earlier in the first part of our research

With respect to Dennis Jones and Wang Min [12] with the emergence of lymphatic-specific markers, further characterization of the underlying molecular mechanisms for lymph angiogenesis may provide a therapeutic avenue for selective inhibition of lymphatic vessels in diseases such as cancer. On the other hand, stimulation of lymph angiogenesis may be beneficial in diseases of lymphatic insufficiency. Additional study of lymphatic vessel regulation will yield further insight into recent implications of their contribution to transplant rejection, obesity, hypertension, and other metabolic and inflammatory disorders.

Results of this study agreed also with Dennis Jones and Wang Min [12] who stated implications of lymphatic stimulation on obesity, hypertension, and other metabolic and inflammatory disorders. As Lymphatic drainage, techniques improved the level of Triglycerides and Fat mass in chronic atherosclerotic cases.



© 2015 Global Journals Inc. (US)

Moreover, Palumbo P1 et. al. [13] effects of a new low frequency, high intensity ultrasound technology on human adipose tissue ex vivo were studied. In particular, they investigated the effects of both external and surgical ultrasound-irradiation, in our experimental conditions, both transcutaneous and surgical ultrasound exposure caused a significant weight loss and fat release. This effect was more relevant when the ultrasound intensity was set at 100 % (~2.5 W/cm$^2$, for external device; ~19-21 W/cm2, for surgical device) compared to 70 % (~1.8 W/cm$^2$ for external device; ~13-14 W/cm2 for surgical device), the effectiveness of ultrasound was much higher when the tissue samples were previously infiltrated with saline buffer, in accordance with the knowledge that ultrasonic waves in aqueous solution better propagate with a consequently more efficient cavitation process. On the other hand, the overall effects of ultrasound irradiation did not appear immediately after treatment but persisted over time, being significantly more relevant at 18 h from the end of ultrasound irradiation. and a significant increase mainly of triglycerides and cholesterol.

The data obtained in our study revealed statistical significance changes in fat mass and the significant improvement (reduction) in the Total blood serum triglycerides for the group A (Class I) and, group B class II groups at the three stages of the measurements (Pre, Post, and Post II). revealed that the combination of lymphatic drainage technique with the low frequency ultrasound reduce the Triglycerides rather than using the low frequency ultrasound only and revealed that there is a positive significant correlation between the reduction of fat mass post-treatment and the reduction of the total body Triglycerides.

This also was consistence with Katsunori Nonogakia et. al. [14], they examined the effects of low-frequency and low-intensity ultrasound (800 kHz, 2mW/cm2), Subjects underwent ultrasound irradiation applied to the forearm for 30 min at 800 kHz and 2 mW/cm2 with 100% duty, or placebo irradiation. After treatment for 30 min, BP and pulse rate were again measured using a BP monitoring system, There were no significant differences in age, BMI, serum high-density lipoprotein-cholesterol (HDL-c), low-density lipoprotein-cholesterol (LDL-c), triglyceride (TG), high sensitive-C-reactive protein (hs-CRP), fasting plasma glucose (FPG), HbA1c, systolic and diastolic BP, pulse rate, pulse pressure, CO, TPR, or CI between the placebo controls and ultrasound treatment group Systolic and diastolic BP, pulse rate, pulse pressure, CO, and CI in the ultrasound treatment group, however, were significantly lower than the baseline values in subjects with hypertension, and lower than those of placebo controls.

The present study relayed on objective method of assessment as body composition analysis and Triglycerides evaluation which is considered one of the objective methods that increase the reliability and validity of the study. Results show reduction in Triglycerides that can improve many cardiac cases complications as hyper tension.

## V. Conclusion

Low frequency ultrasound plus lymphatic drainage technique can be considered a safe and well tolerated method with no life threatening side effects. LFUS plus LD technique improves the blood serum triglycerides fat mass, fat free mass, body water composition, in different types of cardiac obese patients.

Low frequency ultrasound plus lymphatic drainage as a technique could be used as an alternative to conventional exercise and alternative to many obesity surgery as Liposuction surgery and thus provide an opportunity to improve the quality of obese cardiac patient as many obesity surgery have a lot of hazards and sides effect that may affect the patient especially those with cardiac conditions, and that result could be achieved with the usage of LFUS plus LD technique in order to reduce the blood serum triglycerides, fat cells size and even destruction of adiposities with acceptable penetration and get rid of that cells out of the body preventing hazards and complication as thrombosis and many cardiovascular complicated that can be a result of high blood serum triglycerides and high total body fat mass.

## VI. Acknowledgments

The authors wish to thank all participants in the study for their cooperative participation and gratefully thanks the Palestinian hospital for allowing us to use the research center and Laboratory. Also the authors express their sincere thanks to referees for careful reading and helpful suggestions.

## References Références Referencias

1. Finn AV, Nakano M, Narula J, Kolodgie FD, Virmani R, Concept of vulnerable/unstable plaque, *Arterioscler. Thromb. Vasc. Biol*., 30 (7) (2010), 1282–92.
2. Luc G, Fievet C, Arveiler D, et al., Apolipoproteins C-III and E in apoB- and non-apoB-containing lipoproteins in two populations at contrasting risk for myocardial infarction: the ECTIM study. Etude Cas Temoins sur Infarctus du Myocarde. *J. Lipid. Res*., 37 (2009), 508–517.
3. Yarnell JW, Patterson CC, Sweetnam PM, et al., Do total and high density lipoprotein cholesterol and triglycerides act independently in the prediction of ischemic heart disease? Ten-year follow-up of Caerphilly and Speedwell Cohorts, *Arterioscler Thromb Vasc Biol*. ; 21(2010), 1340–1345
4. EJ, McNamara JR, Shah PK, et al., Elevated remnant-like particle cholesterol and triglyceride



© 2015 Global Journals Inc. (US)

levels in diabetic men and women in the Framingham Offspring Study, *Diabetes Care*., 25. (2002), 989–994
5. Jackson, AS; Stanforth, PR; Gagnon, J; Rankinen, T; Leon, AS; Rao, DC; Skinner, JS; Bouchard, C. True, *Int. J. Obesity*, 26 (6) (2008),789–96
6. Fantuzzi G., and Mazzone T. (2007): Adipose tissue and atherosclerosis; exploring the connection. Arterioscler ThrombVascBiol, 27: 996-1003
7. Hunter G. R., Brock D. W., Byrne N. M., Chandler-Laney P., Del Coral P., andGower B. A. Exercise training prevents regain of visceral fat for 1-year following weight loss, *Obesity*, 18 (4) (2010), 690–695
8. Waston, T., Influence of Therapeutic Ultrasound on Skeletal Muscle Regeneration Following Blunt Contusion, *Int. J. Sports Medicine*, 25 8(2006), 5-99.
9. Rose and Wilson, Anatomy & physiology in health and illness Chapter 6 The Lymphatic system (2006), 32-137
10. Reeder LB, Yang LH, Ferguson MK. Modulation of lymphatic spontaneous contractions by EDRF. J Surg Res 2009 56: 620–5
11. von der Weid P-Y, Crowe MJ, van Helden DF. Endothelium-dependent modulation of pacemaking in lymphatic vessels of the guinea-pig mesentery*, J Physiol*, 493 (2010), 563–75.
12. Dennis Jonesand Wang Min, An overview of lymphatic vessels and their emerging role in cardiovascular disease, *J. Cardiovasc Dis Res*., 2(3) (2011), 141–152.
13. Palumbo P1, Cinque B, Miconi G, La Torre C, Zoccali G, Vrentzos N, Vitale AR, Leocata P, Lombardi D, Lorenzo C, D'Angelo B, Macchiarelli G, Cimini A, Cifone MG, Giuliani M., Biological effects of low frequency high intensity ultrasound application on ex vivo human adipose tissue. *Int. J. Immunopathol Pharmacol*, 24(2) (2011), 411-22
14. Katsunori Nonogakia, Marina Suzukia, Marin Sanukia, Naohiko Nonogakib, Kouji Takedac, Nobuyuki Tsujitac, Shuichi Katohd, Nariki Kubotad, Low-frequency and very low-intensity ultrasound decreases blood pressure in subjects with hypertension, *Int. J. Cardiology*, 168 (2013), 1585–1586



© 2015 Global Journals Inc. (US)

Global Journal of Medical Research: K
Interdisciplinary
Volume 15 Issue 6 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

# MRI-T1 and T2 Image Fusion for Brain Image using CDF Wavelet based on Lifting Scheme

By A. Bengana, I. Boukli Hacene & M. A. Chikh
*University of Tlemcen, Algeria*

*Abstract-* In the field of medical imaging. Image fusion is an important application for extracting complementary information from different modality. In this work, we propose a fusion algorithm using CDF9/7 wavelet based on lifting scheme with specified fusion rules to combine pairs of multispectral Magnetic Resonance Imaging (MRI) such as T1, T2. The experimental results of brain tumor show that the proposed algorithm preserves both edge and component information and also increases the efficiency of tumor detection. The parameters like mutual information MI, entropy EN, and spatial frequency SF, standard deviation SD are calculated to evaluate performance of proposed algorithm. Finally the results are compared with existing methods.

*Keywords: image fusion, multispectral MRI-T1, MRI-T2, brain tumor, CDF9/7 wavelet, lifting scheme.*

*GJMR-K Classification: NLMC Code: WN 180*




*Strictly as per the compliance and regulations of:*

© 2015. A. Bengana, I. Boukli Hacene & M. A. Chikh. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# MRI-T1 and T2 Image Fusion for Brain Image using CDF Wavelet based on Lifting Scheme

A. Bengana [α], I. Boukli Hacene [σ] & M. A. Chikh [ρ]

*Abstract-* In the field of medical imaging. Image fusion is an important application for extracting complementary information from different modality. In this work, we propose a fusion algorithm using CDF9/7 wavelet based on lifting scheme with specified fusion rules to combine pairs of multispectral Magnetic Resonance Imaging (MRI) such as T1, T2. The experimental results of brain tumor show that the proposed algorithm preserves both edge and component information and also increases the efficiency of tumor detection. The parameters like mutual information MI, entropy EN, and spatial frequency SF, standard deviation SD are calculated to evaluate performance of proposed algorithm. Finally the results are compared with existing methods.

*Keywords: image fusion, multispectral MRI-T1, MRI-T2, brain tumor, CDF9/7 wavelet, lifting scheme.*

## I. Introduction

The medical image has seen an important development in the last two decades such as computed tomography (CT), magnetic resonance image (MRI), positron emission tomography (PET), and single photon emission computed tomography (SPECT). Now, the doctors need information from more than one modality for efficient disease detection and diagnosis. Medical image fusion is the technique of merging images belonging to different modality into a single resultant image that is provide complementary information for better analysis.

Image fusion can be grouped into three categories: Pixel level, Region level and Decision level. There are two approaches for pixel level method: spatial domain and transform domain **[4]**. Spatial domain fusion techniques are simple and fused image can be obtained by directly applying fusion rules on pixel values of source images such as Simple averaging, PCA (Principal Component Analysis) **[5]** and linear fusion **[6]**. But major disadvantage of spatial domain techniques are that they introduce spatial distortions in the fused image **[7]** and do not provide any spectral information. Transform domain techniques overcome the disadvantages of spatial domain fusion. Pyramid and wavelet transforms based techniques are the mostly used transform domain image fusion methods. Laplacian pyramid **[8],** contrast pyramid **[9].** These methods overcome the disadvantages of spatial domain techniques but suffer from blocking effect **[7]**. As a result wavelet transform based fusion **[11, 12]** approaches are used and it was found that no blocking effect occurred during fusion process. Discrete wavelet transform (DWT) [3] and integer wavelet transform [2] preserves different frequency information but may produce specularities along the edges and shift sensitivity, poor directionality and lack of phase information **[15, 16]**. The third type is new multiresolution methods are proposed, such as image fusion based on nonsubsampled contourlet transform (NSCT) **[17]**. Due to these limitations of real valued DWT. Another approach uses biorthogonal wavelet CDF 9/7 **[18]** based on lifting scheme provides better representation of fused image.

The objective of this research is to introduce CDF 9/7 wavelet based on lifting scheme to fuse more properties (edge, information) of different modalities such as MRI-T1 and MRI-T2 brain images (normal and tumor). The proposed algorithm is also uses the specified fusion rules of low and high frequency subbands. Experimental results on the Brain Web database show the usefulness of this wavelet family and clearly indicate its potential in medical image fusion.

The rest of paper organization of this is as follows, the section 2 we present the proposed fusion algorithm steps with fusion rules. The experimental results are shown in section 3 and we compare results obtained with the existing techniques. Finally the main conclusions are summarized in Section 4.

## II. Proposed Fusion Algorithm

The input Images can suffer from artifacts due to different factors. However the recent advances in acquisition protocol make it possible to acquire images with very limited artifacts. The proposed method is illustrated in Figure 1

*Author α σ ρ: Biomedical Laboratory, Department of electrics and Electronics, Technology Faculty. University of Tlemcen 13000, Algeria. e-mails: befatih80@yahoo.fr, ismaill80@yahoo.fr, mea_chikh@mail.univ-tlemcen.dz*



© 2015 Global Journals Inc. (US)





*Figure 1 :* Block diagram fusion using CDF9/7 lifting wavelet transform (LWT)

*a) Image de-noising using median filter (pre-processing)*

This proposed system describes the information of enhancement using weighted median filter for removing high frequency component which characterize the noise. These filters have the robustness and edge preserving capabilities with noise attenuation characteristics. After the pre-processing operations the input images are subjected to wavelet analysis which is described in the following section.

*b) Biorthogonal CDF9/7 wavelet based on lifting scheme [1]*

Sweldens from Bell Labs proposed a method which does not depend on the Fourier transform of ascension wavelet construction in the mid 1990's presented in [19, 20], where sweldens showed that the convolution based biorthogonal Wavelet Transform (WT) can be implemented in a lifting-based scheme. The lifting-based WT consists of splitting, lifting, and scaling modules and the WT is treated as prediction-error decomposition. It provides a complete spatial interpretation of WT.

In this article we deals with biorthogonal wavelet 9/7, these wavelets are part of the family of symmetric biorthogonal wavelet CDF. The low pass filters associated with wavelet 9/7 have p=9 coefficients in the analysis, p=7 coefficients to synthesize. The wavelets 9/7 have a great number of null moments for a relatively short support. They are more symmetrical and very close to orthogonality [21]. Were the first to show the superiority of the biorthogonal wavelet transform 9/7 for the decorrelation of natural images





*Figure 2 :* The Lifting -based wavelet transform

*c) Fusion rules*

The input medical image (A, B) are decomposed using CDF9/7 lifting wavelet transform to obtain the low frequency ($LL_A$, $LL_B$) and high frequency ($LH_A$, $HL_A$, $HH_A$, $LH_B$, $HL_B$, $HH_B$) coefficients.

i. *Lowpass subband fusion*

The approximation coefficients are averaged, to preserve the features from both the images

$$LL_F = mean(LL_A, LL_B) \quad (1)$$

Where $LLF$ is the approximation band of the fused image.

ii. *Highpass subband fusion*

For each detailed sub bands, the spatial frequency is calculated using Equation SF.

$$LH_F = \begin{cases} LH_A & if \quad SF_{LH_A} > SF_{LH_B} \\ LH_B & otherwise \end{cases} \quad (2)$$

Apply the same technique for: $HL_A$, $HH_A$, $HL_B$, $HH_B$

Finally, the inverse lifting transform based on CDF9/7 wavelet is applied to generate the fused medical image.

## III. Results and Discussion

In this section to evaluate the performance of the proposed algorithm. We fuse normal brain MRI-T1 image with MRI-T2 image from [24] data base. Similarly Brain tumor hypo intense T1 image with hyper intense T2 image are downloaded from [25].

To show the effectiveness of the multi scale transform the comparisons start with PCA (Principal Component Analysis) method, CDF9/7 DWT (discret wavelet transform) based filter banc and CDF9/7 LWT (Lifting wavelet transform) with the first level. To quantitatively compare the performance with existing fusion algorithms, entropy EN, mutual information MI, spatial frequency SF, standard deviation SD and time calcul T , are defined and calculated in the following section. These images were tested on Intel *Core* (I3) 2.13 GHz PC with 2GB of RAM using Matlab 2010a.



© 2015 Global Journals Inc. (US)

*a) Entropy*

Entropy is the quantity of information contained in a series of events. Entropy is a criterion that measures the level of information in the image, more entropy is large, more diffuse the image information.

$$EN = -\sum_{i=0}^{L-1} p_f(i) log_2 p_f(i) \quad (3)$$

*b) Mutual Information (MI)*

Compares the image source and the fused image, more value is small, the relationship between the two images is non-existent.

$$MI = \sum_{af} P_{AF}(a,f) log \frac{P_{AF}(a,f)}{P_A(a)P_F(f)} \quad (4)$$

$P_{AF}(a,f)$ The joint histogram between fused image F and the source image A.

*c) Standard Deviation (SD)*

Is the square root of the variance, the variance of an image reflects the degree of dispersion between the grayscale values and the average value of gray levels, more the standard deviation is large, more there are dispersion

$$SD = \sqrt{\frac{\sum_{i=0}^{N-1}\sum_{j=0}^{M-1} F(i,j))}{NM}} \quad (5)$$

*d) Spatial Frequency (SF)*

It measures the total activity and the level of clarity of an image, an important value mean that the result of fusion is good.

$$SF = \sqrt{RF^2 + CF^2} \quad (6)$$

Where *RF* and *CF* are the row frequency and the column frequency respectively.

The fused image output based on proposed method illustrate in Figure 3 and 4.





*Figure 3 :* Fusion results for Normal brain, (a) MRI-T1 image, (b) MRI-T2 image, (c) PCA fusion, (d) CDF9/7 DWT based fusion, (e) proposed fusion method





*Figure 4 :* Fusion results for Brain tumor, (a) brain image with T1 weighting, (c) PCA fusion, (d) CDF9/7 DWT based fusion, (e) proposed fusion method

This visual representation is not sufficient for analysis of the obtained fusion results. Therefore we have compared the proposed method with other ones on quantitative measures which are entropy, standard deviation, spatial frequency, mutual information.

*Table 1 :* Performance comparison for first set of Brain Web normal database

| Methods | EN | SF | SD | MI | T(s) |
|---|---|---|---|---|---|
| PCA | 5.56 | 24.89 | 49.77 | 3.92 | 1.11 |
| CDF9/7DWT | 6.23 | 38.04 | 51.10 | 5.43 | 1.78 |
| CDF9/7LWT | 7.33 | 46.92 | 97.84 | 5.80 | 1.46 |


*Table 2 :* Performance comparison for second set of Brain tumor


| Methods | EN | SF | SD | MI | T(s) |
|---|---|---|---|---|---|
| PCA | 6.93 | 23.83 | 49.43 | 3.20 | 0.02 |
| CDF9/7DWT | 7.01 | 35.27 | 49.36 | 5.85 | 1.24 |
| CDF9/7LWT | 7.12 | 44.42 | 89.23 | 5.98 | 1.52 |


For Table1 and 2, The graphical representation of all performance evaluation parameters are showns in figure 5.



© 2015 Global Journals Inc. (US)








*Figure 5 :* The graphical of different methods using evaluation parameters

## IV. Conclusion

In the present work, we have introduce a new multimodal medical image fusion using CDF9/7 wavelet based on lifting scheme and its essential objective as to retrieve complementary information from different modality to aid doctors for a good diagnosis. The algorithm is evaluated on the Brain Web database and brain tumor. The experimental results showed that the proposed algorithm conserves important edge and spectral information without much of spatial distortion. Finally, these results are compared with different methods (PCA and CDF9/7 DWT based filter banc). In perspective, we plan to extend our study (proposed algorithm) by incorporating learning techniques.

## References Références Referencias

1. Beladgham, M., A. Bessaid, A. Moulay Lakhdar and A. Taleb Ahmed, (2011). 'Improving quality of medical image compression using biorthogonal CDF wavelet based on lifting scheme and SPIHT coding', *Serbian Journal of Electrical Engineering*, Vol 8, Issue 2, ISSN 1451-4869 , pp 163-179, doi:10.2298/SJEE1102163B, 2011.
2. Z .Wang, Yu X and Zhang L.B.(2008). 'A Remote Sensing Image Fusion Algorithm Based on Integer Wavelet Transform'. *Journal of Optoelectronics Laser*, Vol.19, No.11, (November 2008), pp. 1542-1545.
3. I. Daubechies. *Ten lectures on wavelets*. Society for Industrial and Applied Mathematics, 1992.
4. R. Singh, R. Srivastava, O. Prakash, A. Khare, Mixed scheme based multimodal medical image fusion using Daubechies Complex Wavelet Transform, IEEE/OSA/IAPR International Conference on Infonnatics, Electronics & Vision (ICIEV 2012), pp. 303-309, 2012



© 2015 Global Journals Inc. (US)

5. V.P.S. Naidu, J.R. Rao, Pixel-level Image Fusion using Wavelets and Principal Component Analysis, Defence Science Journal, vol. 58, No. 3, pp. 338-352,2008.
6. J. G. P. W. Clevers, R. Zurita-Mill a, Multisensor and multiresolution image fusion using the linear mixing model, Image Fusion: Algorithms and Applications, pp. 67-84, 2008.
7. H. Li, B. S. Manjunath, S. K. Mitra, Multisensor image fusion using the wavelet transform, Graphical Models and Image Processing, vol. 57, no. 3,pp. 235-245,1995.
8. P. J. Burt, R. J. Kolczynski, Enhanced image capture through fusion, in Proceedings of the 4th IEEE International Conference on Computer Vision (ICCV '93), pp. 173-182, 1993.
9. A. Toet, J. J. Van Ruyven, J. M. Valeton, Merging thermal and visual images by a contrast pyramid, Optical Engineering, vol. 28, no. 7, pp. 789-792, 1989.
10. G. Pajares, 1. M. Cruz, A wavelet-based image fusion tutorial, Pattern Recognition, vol. 37, no. 9, pp. 1855-1872, 2004.
11. K. Amolins, Y. Zhang, P. Dare, Wavelet based image fusion techniques-an introduction, review and comparison, ISPRS Journal of Photogrammetry & Remote Sensing, vol. 62, no. 4, pp. 249-263, 2007.
12. A. Khare, U. S. Tiwary, M. Jeon, Daubechies complex wavelet transform based multilevel shrinkage for deblurring of medical images in presence of noise, International Journal on Wavelets, Multiresolution and Information Processing, vol. 7, no. 5, pp. 587-604, 2009.
13. A. Khare, M. Khare, Y. Y. Jeong, H. Kim, M. leon, Despeckling of medical ultrasound images using complex wavelet transform based Bayesian shrinkage, Signal Processing, vol. 90, no. 2, pp. 428-439, 2010.
14. Kundu, M. K., & Das, S. (2012). NSCT-based multimodal medical image fusion using pulse-coupled neural network and modified spatial frequency. International Federation for Medical and Biological Engineering, 50, 1105–1114.
15. I. Boukli H, Bessaid. A. (2014). 'Hybrid colour medical image compression by CDF wavelet and cosine transforms' *Int. J. Biomedical Engineering and Technology,* Vol. 16, No. 1, 2014
16. W. Sweldens, (1996). 'The lifting scheme: A custom-design construction of biorthogonal wavelets', *Appl Comput Harmon Anal*, pp. 186-200, 1996.
17. Calderbank, A.R., I. Daubechies, W. Sweldens, and BL. Yeo, (1998). 'Wavelet transforms that map integers to integers', *Appl Comput Harmon Anal*, 5, pp. 332-369, 1998.
18. Antonini, M., Barlaud, M., Mathieu, P. and Daubechies, I. (1992). 'Image coding using wavelet transform'. *IEEE Trans. Image Processing*, Vol. 1, April 1992, pp. 205–220.
19. http://www.bic.mni.mcgill.ca/Brain Web
20. http://www.scielo.br




© 2015 Global Journals Inc. (US)

# Global Journals Inc. (US) Guidelines Handbook 2015

www.GlobalJournals.org

# FELLOWS

## FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSM" title to individuals.The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSM" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall,Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

*The following benefits can be availed by you only for next three years from the date of certification:*

FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.





© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSM, you will be given a renowned, secure and free professional email addres with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.




The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.




The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of you credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.




The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.



The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.

## MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSM" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.

MARSM accrediting is an honor. It authenticates your research activities. Afterbecoming MARSM,you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

*The following benefitscan be availed by you only for next three years from the date of certification.*




MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you willbe given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook




We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.






Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

# AUXILIARY MEMBERSHIPS

## Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

*The Institute will be entitled to following benefits:*

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.




The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.




After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

**The following entitlements are applicable to individual Fellows:**




Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

**Other:**

**The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:**

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

**Note :**

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

# Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

**(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.**

**(II) Choose corresponding Journal.**

**(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.**

**(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.**

**(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.**

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

# Preferred Author Guidelines

**MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)**

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

**You can use your own standard format also.**
**Author Guidelines:**

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

**1. GENERAL**

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

**Scope**

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

**2. ETHICAL GUIDELINES**

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

**Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission**

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

**Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.**

**Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.**

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

**3. SUBMISSION OF MANUSCRIPTS**

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

**4. MANUSCRIPT'S CATEGORY**

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

**5.STRUCTURE AND FORMAT OF MANUSCRIPT**

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

**Papers**: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

**Format**

*Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.*

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

**Structure**

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

*Abstract, used in Original Papers and Reviews:*

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

*Acknowledgements: Please make these as concise as possible.*

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

*Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.*

*Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.*

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

*Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.*

**6. AFTER ACCEPTANCE**

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

**6.1 Proof Corrections**

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

**6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)**

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

**6.3 Author Services**

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

**6.4 Author Material Archive Policy**

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

**6.5 Offprint and Extra Copies**

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

## TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

**1. Choosing the topic:** In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

**2. Evaluators are human:** First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

**3. Think Like Evaluators:** If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

**4. Make blueprints of paper:** The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

**5. Ask your Guides:** If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

**6. Use of computer is recommended:** As you are doing research in the field of Computer Science, then this point is quite obvious.

**7. Use right software:** Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

**8. Use the Internet for help:** An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

**9. Use and get big pictures:** Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

**10. Bookmarks are useful:** When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

**11. Revise what you wrote:** When you write anything, always read it, summarize it and then finalize it.

**12. Make all efforts:** Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

**13. Have backups:** When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

**14. Produce good diagrams of your own:** Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

**15. Use of direct quotes:** When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

**16. Use proper verb tense:** Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

**17. Never use online paper:** If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

**18. Pick a good study spot:** To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

**19. Know what you know:** Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

**20. Use good quality grammar:** Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

**21. Arrangement of information:** Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

**22. Never start in last minute:** Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

**23. Multitasking in research is not good:** Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

**24. Never copy others' work:** Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

**25. Take proper rest and food:** No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

**26. Go for seminars:** Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

**27. Refresh your mind after intervals:** Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

**28. Make colleagues:** Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

**29. Think technically:** Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

**30. Think and then print:** When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

**31. Adding unnecessary information:** Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

**32. Never oversimplify everything:** To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

**33. Report concluded results:** Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

**34. After conclusion:** Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

## Informal Guidelines of Research Paper Writing

**Key points to remember:**

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

**Final Points:**

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

**General style:**

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

**Title Page:**

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



**Abstract:**

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

**Introduction:**

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

**Procedures (Methods and Materials):**

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

**Results:**

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

**Discussion:**

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

**Segment Draft and Final Research Paper:** You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

## Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

**Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).**

| Topics | Grades | | |
|---|---|---|---|
| | **A-B** | **C-D** | **E-F** |
| ***Abstract*** | Clear and concise with appropriate content, Correct format. 200 words or below | Unclear summary and no specific data, Incorrect form<br><br>Above 200 words | No specific data with ambiguous information<br><br>Above 250 words |
| ***Introduction*** | Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited | Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter | Out of place depth and content, hazy format |
| ***Methods and Procedures*** | Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads | Difficult to comprehend with embarrassed text, too much explanation but completed | Incorrect and unorganized structure with hazy meaning |
| ***Result*** | Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake | Complete and embarrassed text, difficult to comprehend | Irregular format with wrong facts and figures |
| ***Discussion*** | Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited | Wordy, unclear conclusion, spurious | Conclusion is not cited, unorganized, difficult to comprehend |
| ***References*** | Complete and correct format, well organized | Beside the point, Incomplete | Wrong format and structuring |

# INDEX


## A

Adiponectin · 29
Atherogenic · 28, 29
Atrioventricular · 19

## C

Cardiopulm · 26

## E

Electromyogram · 1, 15

## G

Giallauria · 23, 25

## H

Haddadzadeh · 21, 25

## K

Kolmogorov · 3
Kuribayashi · 3, 14

## L

Lymphedema · 31

## P

Percutaneous · 17, 25

## R

Rheumatological · 1

## S

Soleimannejad · 21, 25
Sultornsaneea · 15

## T

Tilkemeier · 26
Triglycerides · 27, 33, 36, 37, 38, 39, 40



save our planet



# Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.MedicalResearchJournal.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals