

GLOBAL JOURNAL

OF MEDICAL RESEARCH: K

Interdisciplinary

Cerebellar Hemangioblastoma

Infection Prevalence in a Tertiary

Highlights

Health Informatics Integrated

Maternal Health Care Utilization

Discovering Thoughts, Inventing Future

VOLUME 14

ISSUE 3

VERSION 1.0



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

VOLUME 15 ISSUE 3 (VER. 1.0)

OPEN ASSOCIATION OF RESEARCH SOCIETY

© Global Journal of Medical Research . 2015.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website <http://globaljournals.us/terms-and-condition/menu-id-1463/>

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)

Sponsors: *Open Association of Research Society*
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please
email us at local@globaljournals.org

eContacts

Press Inquiries: press@globaljournals.org
Investor Inquiries: investors@globaljournals.org
Technical Support: technology@globaljournals.org
Media & Releases: media@globaljournals.org

Pricing (Including by Air Parcel Charges):

For Authors:

22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

INTEGRATED EDITORIAL BOARD
(COMPUTER SCIENCE, ENGINEERING, MEDICAL, MANAGEMENT, NATURAL
SCIENCE, SOCIAL SCIENCE)

John A. Hamilton, "Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software
Engineering
Director, Information Assurance
Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at
Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D. and M.S. Syracuse University,
Syracuse, New York
M.S. and B.S. Bogazici University,
Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A. Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range
Nutritionist
Ph.D. Edinburgh University - Animal
Nutrition
M.S. Aberdeen University - Animal
Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing &
Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems
Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinipiac
BS, Jilin Institute of Technology; MA, MS,
PhD, (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley
& Istanbul University

Dr. Bart Lambrecht

Director of Research in Accounting and Finance
Professor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD
(Cambridge)

Dr. Carlos García Pont

Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management),
Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE,
University of Navarra
Degree in Industrial Engineering,
Universitat Politècnica de Catalunya

Dr. Fotini Labropulu

Mathematics - Luther College
University of Regina
Ph.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windsor

Dr. Lynn Lim

Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD,
FHEA

Dr. Mihaly Mezei

ASSOCIATE PROFESSOR
Department of Structural and Chemical
Biology, Mount Sinai School of Medical
Center
Ph.D., Eötvös Loránd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram

Department of Accounting and Finance
Lancaster University Management School
Ph.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño

Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso

Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.

Director, EP Laboratories, Philadelphia VA
Medical Center
Cardiovascular Medicine - Cardiac
Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng

MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular
Medicine
David R. Davies Department of Neurology and Clinical
Neuroscience
Northwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli

Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic
Radiology
M.D., State University of New York at
Buffalo, School of Medicine and
Biomedical Sciences

Dr. Roberto Sanchez

Associate Professor
Department of Structural and Chemical
Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun

Professor of Earth and Atmospheric
SciencesPurdue University Director
National Center for Typhoon and
Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences,
National Central University, Chung-Li,
TaiwanUniversity Chair Professor
Institute of Environmental Engineering,
National Chiao Tung University, Hsin-
chu, Taiwan.Ph.D., MS The University of
Chicago, Geophysical Sciences
BS National Taiwan University,
Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick

M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and
Hypertension Division (PMC)
Penn Medicine, University of
Pennsylvania
Presbyterian Medical Center,
Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of
Internal Medicine

Dr. Bassey Benjamin Esu

B.Sc. Marketing; MBA Marketing; Ph.D
Marketing
Lecturer, Department of Marketing,
University of Calabar
Tourism Consultant, Cross River State
Tourism Development Department
Co-ordinator , Sustainable Tourism
Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.

IEEE Senior Member
Chairperson, Department of Computer
Science
AUST - American University of Science &
Technology
Alfred Naccash Avenue – Ashrafieh

PRESIDENT EDITOR (HON.)

Dr. George Perry, (Neuroscientist)

Dean and Professor, College of Sciences

Denham Harman Research Award (American Aging Association)

ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization

AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences

University of Texas at San Antonio

Postdoctoral Fellow (Department of Cell Biology)

Baylor College of Medicine

Houston, Texas, United States

CHIEF AUTHOR (HON.)

Dr. R.K. Dixit

M.Sc., Ph.D., FICCT

Chief Author, India

Email: authorind@computerresearch.org

DEAN & EDITOR-IN-CHIEF (HON.)

Vivek Dubey(HON.)

MS (Industrial Engineering),

MS (Mechanical Engineering)

University of Wisconsin, FICCT

Editor-in-Chief, USA

editorusa@computerresearch.org

Sangita Dixit

M.Sc., FICCT

Dean & Chancellor (Asia Pacific)

deanind@computerresearch.org

Suyash Dixit

(B.E., Computer Science Engineering), FICCTT

President, Web Administration and

Development , CEO at IOSRD

COO at GAOR & OSS

Er. Suyog Dixit

(M. Tech), BE (HONS. in CSE), FICCT

SAP Certified Consultant

CEO at IOSRD, GAOR & OSS

Technical Dean, Global Journals Inc. (US)

Website: www.suyogdixit.com

Email: suyog@suyogdixit.com

Pritesh Rajvaidya

(MS) Computer Science Department

California State University

BE (Computer Science), FICCT

Technical Dean, USA

Email: pritesh@computerresearch.org

Luis Galárraga

J!Research Project Leader

Saarbrücken, Germany

CONTENTS OF THE ISSUE

- i. Copyright Notice
 - ii. Editorial Board Members
 - iii. Chief Author and Dean
 - iv. Contents of the Issue
-
- 1. Health Informatics Integrated System Post-Implementation Evaluation. *1-37*
 - 2. Polycythemia with Cerebellar Hemangioblastoma. *39-45*
 - 3. My Pain is 10 out of 10. Patients Vs. Actors in Clinical Settings. *47-52*
 - 4. Community Acquired Urinary Tract Infection Prevalence in a Tertiary Institution Based in Evbuobanosa, Edo State, Nigeria. *53-64*
 - 5. Maternal Health Care Utilization in the Eastern Cape, South Africa: A Qualitative Investigation. *65-70*
 - 6. A Case of a Non-Verbal Child with Autism Gaining Faculty of Speech with Applied Energy Medicine. *71-73*
-
- v. Fellows and Auxiliary Memberships
 - vi. Process of Submission of Research Paper
 - vii. Preferred Author Guidelines
 - viii. Index



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Health Informatics Integrated System Post- Implementation Evaluation

By Sajeesh Kumar PhD, Anne L. Pedigo, RN MSN

University of Tennessee Health Science Center, United States

Abstract- Part of a well-designed health informatics implementation process includes the mechanisms put in place to help the day-to-day operators of the systems. Continual appraisal of these methods necessitates up-to-date investigations. Understanding critical elements which support a positive transition of health information technology (HIT) within healthcare facilities is the objective of the following research. To help develop these findings, a prospective post-implementation and use assessment survey was conducted on two hospitals in Central Texas. The population studied included RN case managers, social workers and supportive staff in the Continuum of Care departments at two Scott & White Healthcare acute care facilities. The implementation process appeared to provide a mostly encouraging transition with a small number of components noted of concern to the staff. Areas of enhancement were revealed included improving training specific to job roles and supplying more fitting integration of processes and workflows.

Keywords: health informatics; evaluation, postimplementation.

GJMR-K Classification: NLMC Code: W 26.5



HEALTHINFORMATICSINTEGRATEDSYSTEMPOSTIMPLEMENTATIONEVALUATION

Strictly as per the compliance and regulations of:



Health Informatics Integrated System Post-Implementation Evaluation

Sajeesh Kumar PhD ^α, Anne L. Pedigo ^σ RN MSN ^ρ

Abstract- Part of a well-designed health informatics simple implementation process includes the mechanisms put in place to help the day-to-day operators of the systems. Continual appraisal of these methods necessitates up-to-date investigations. Understanding critical elements which support a positive transition of health information technology (HIT) within healthcare facilities is the objective of the following research. To help develop these findings, a prospective post-implementation and use assessment survey was conducted on two hospitals in Central Texas. The population studied included RN case managers, social workers and supportive staff in the Continuum of Care departments at two Scott & White Healthcare acute care facilities. The implementation process appeared to provide a mostly encouraging transition with a small number of components noted of concern to the staff. Areas of enhancement were revealed including improving training specific to job roles and supplying more fitting integration of processes and workflows.

Keywords: health informatics; evaluation, post-implementation.

I. HEALTH INFORMATICS INTEGRATED SYSTEM POST-IMPLEMENTATION EVALUATION

When one speaks of a successful health information technology (HIT) system implementation, there are several dimensions that go into determining that success. While the satisfaction of the workforce is very important, it is only one dependent factor tied to how well the healthcare process has succeeded. How well current practices are redesigned to take advantage of the technology is a factor. Quality of the data is another influence. Confidence in the documentation and the information it contains is an important aspect. How a system will work through barriers and enable facilitators are other dimensions of a successful implementation. Measurement of improvement to patient care is another facet. So many characteristics go into determining a successful implementation. Finding the right instruments to put into position before, during and after an HIT system implementation is an ongoing task that continually needs to be evaluated. As with the integrated systems, implementation standards need to be studied and

enhanced to strive for even better success. A well designed process should allow for success that is on par or surpasses the importance of the former.

II. BACKGROUND

A little over a decade ago, the Institute of Medicine put forward that improved patient safety, efficiency of health care delivery competences and quality of care would be realized by make use of an effective integrated HIT (*Crossing the Quality Chasm: A New Health System for the 21st Century*, 2001). More recently, government incentives and mandates have been placed on healthcare institutions advocating for their adoption of HIT systems (DHS, 2010). While there are legislative whys and wherefores that go into the need for an HIT system, the drive to have a system that helps the patient and staff needs to be the driving force in the desire to find mechanisms which encourage a positive and effective application.

The purpose of the research topic of interest is to identify elements necessary for a successful HIT system implementation at acute care hospital sites. The research study will help determine what critical elements are necessary to have in place in order for healthcare facilities to have a successful transition from an older medical record system to a new electronic medical record (EMR) system.

The research study will evaluate what operations should be set in place by healthcare facilities before transitioning to an HIT system. Moreover, the research will focus on possible ways to prevent issues that may develop during and from the implementation of the new electronic system. The study will survey employees of healthcare facilities who have already transitioned to an HIT system and examine how they believe the implementation process could be improved. Furthermore, barriers to a successful HIT system implementation will be attempted to be identified. As a final point, information found in the study will be used to synthesize material and identification of possible gaps in research.

Information found in the literature review was employed to integrate data and identify gaps in present research such as the need for greater variety of positions giving feedback. While several of the recommendations for successful implementation were

Author ^{α ρ} : Department of Health Informatics & Information Management, University of Tennessee Health Science Center, 920 Madison Avenue Suite 518, Memphis, Tennessee 38163, USA.
e-mail: skumar10@uthsc.edu

similar, some studies had opposing views of nurses' attitudes after implementation. The type of support by the healthcare facility before and after implementation may have been a factor in these findings. Moreover, in the majority of studies, nurses were the population studied and findings were based on these responses.

Although in all five articles the implementation of a comprehensive HIT system was being evaluated, rarely was health care personnel who work outside of direct patient care evaluated. No staff within areas such as admissions or billing was interviewed (Table 1).

Table 1

Summary of Literature Reviews

Author, Year Published	Research Objective	Study Design, Method, Time Frame, Sample and Response Rate	Instrument Used in Study	Analytical Technique	Key Findings and Limitations
(Kirkendall, Goldenhar, Simon, Wheeler, & Andrew Spooner, 2013)	Examination of perceptions, expectations and experiences in regards to the transition from a CPOE system to a fully integrated HIT SYSTEM by healthcare employees within an inpatient setting.	Design & method: One pre-implementation and one post-implementation online surveys Time frame: January 5-9, 2010 (5-day pre-implementation survey; Open for 5 days) and January 10-February 10, 2011 (1-year post-implementation; Open for 1 month) Sample: 751 5-day pre-implementation survey; 1,954 1-year post-implementation survey (Nurses, prescribers, staff positions and other inpatient staff personnel) Response rate: 5-day pre-implementation survey (5.2%); 1-year post-implementation survey (13.6%)	7-factor structure Information Systems Expectations and Experiences (I-SEE) survey which assessed 1) Provider-patient communication, 2) Inter-provider communication 3) Inter-organizational communication 4) Work-life changes 5) Improved care 6) Support & resources 7) Patient care processes Administered online via REDCap.	Construct validity and reliability was assessed with current & previous results. Exploratory factor analysis resulted in a 7-factor structure giving better reliability & validity. SAS statistical software was utilized.	Key findings: 1) Nurses had less positive attitudes about the transition than non-nursing respondents. 2) Differences diminished after implementation. 3) Nursing scores increased significantly for job satisfaction, quality & safety of patient care, organizational support for transition and the rights of patient care but did not increase significantly for communication at 1 year post survey. Limitations: 1) Survey was administered only 5 days prior to rollout which could have influenced motivation to complete survey. 2) Response rate was fairly low. 3) Possibility of some staff having prior HIT SYSTEM experience in outpatient setting. 4)
(Spetz, Burgess Jr, & Phibbs, 2012)	Identification of influences and tactics associated with successful implementation of hospital-	Design & method: Qualitative retrospective-mixed-methods of semi-structured	A semi-structured interview guide was developed from a review of the literature of technology implementation	A thematic analysis was performed with initial cods drawn from the content of the interview	Key findings: Five broad themes stemmed from interviews that affected the success 1) Organizational stability and implementation team leadership

	based information technology systems by patient-care providers and IT staff within an inpatient setting.	interviews Time frame: June 2006-September 2007 (15-month period) Sample: 118 interviews (Nurses, pharmacists, physicians, IT staff and senior management) Response Rate: Not discussed in article if anyone refused interview.	and the effects of IT systems and suggestions from an Advisory Committee consisting of VA medical, pharmacy, nursing leaders and representatives of the VA headquarters.	guides.	2) Implementation timelines 3) Equipment availability and reliability 4) Staff training 5) Changes in work flow Limitations: 1) A retrospective analysis is limited to the memories which may be inaccurate or biased. 2) Furthermore, some staff are no longer available to interview. 3) In addition, the analysis was conducted by only one investigator which may decrease reliability. 4) Lastly, the VA is unique and experiences may differ from that of a freestanding hospital. 5)
(Laramée, Bosek, Shaner-McRae, & Powers-Phaneuf, 2012)	Comparison of attitudes before implementation and 6 & 18 months after implementation of a comprehensive HIT SYSTEM of nurses within an inpatient setting.	Design & method: One pre-implementation and two post-implementation online surveys Time frame: December 2008 (6-months pre-survey; Open for 4 weeks); December 2009 (6-months post-survey; Open for 4 week); December 2010 (1- months post-survey; Open for 4 week) Sample: 312 6-month pre- survey, 410 6-month post-survey & 262 18-month post-implementation survey (RNs, LPNs, APRNs and Management) Response rate: 6-month pre-survey (18%). 6-	Modified Nurses' Attitude Toward Computerization Questionnaire which reflected the HIT SYSTEM rather than the computer with an open-ended question added for the 6-month post-implementation survey and one multiple choice question & an open-ended question added for the 18-month post-implementation survey. All administered online via REDCap.	Data were analyzed using STATA 10.1 software. Descriptive analysis and χ^2 were used to analyze demographic variables. Two-tailed t tests were used to compare differences between 3 time periods. A modified Colaizzi's method was used for qualitative analysis.	Key findings: 1) Attitudes became less positive after implementation. Pre-implementation (74.2%), 6 months post-implementation (65.9%) & 18 months post-implementation (67.7%). 2) Nurse age & years of experience affect attitude negatively. 3) Documentation improved despite workload impact. 4) Implementation process was a challenging and dramatic change. Limitations: 1) Description of experiences of nurses at one medical facility, generalization to other HIT SYSTEM implementations is limited. 2) Internal validity may be compromised due to the low respond rate & potential selection bias associated with those who did complete survey.

		month post-survey (24%); 18-post-implem survey (15%)			
(A. S. Laramée, Bosek, Kasprisin, & Powers-Phaneuf, 2012)	Exploration of factors and strategies believed to be effective in creating positive attitudes and overcoming barriers leading to previous successful application of HIT SYSTEM in preparation of upcoming new implementation at a rural academic medical center.	Design & method: Descriptive exploratory qualitative research design using semi-structured focus groups interviews Time frame: December 2008 (6-months pre-implementation survey; Open for 4 weeks); December 2009 (6-months post-implementation survey; Open for 4 weeks); December 2010 (1- months post-implementation survey; Open for 4 week) Sample: 40 self-selected members in 11 focus groups (RNs, MDs, managers, nurse educators, unit secretaries, techs, dieticians)	Focus group interviews were conducted using semi-structured questions. A seven-item questionnaire was developed & distributed to staff to validate themes identified in focus groups.	Audiotapes were analyzed utilizing the intuit, analyze & describe method. Triangulation of interdisciplinary team and two clinical departments increased breadth of data. At least two researchers analyzed data from each group.	Key findings: Four major themes found to be fundamental to successful implementation of HIT SYSTEM 1) Reduce unrealistic expectations & fears related to individual competency with initial work with HIT SYSTEM. 2) Allow staff time for individual pursuit of learning about the HIT SYSTEM& their skills in using the system. 3) Clear processes for using the HIT SYSTEM are needed. 4) Make the HIT SYSTEM support individuals accessible 24/7 and make it customer-focused. Limitations: Limitations were not discussed in article. Assurance was given regarding the reliability and validity of the qualitative data analysis.
(Ward, Vartak, Schwichtenberg, & Wakefield, 2011)	Assessment of impact of workflow and patient care from the employment of an HIT SYSTEM on nurses within a rural referral hospital.	Design & method: Two pre-implementation paper surveys and one post-implementation online survey Time frame: No specific date is given; Day one of training expectations	7-factor structure Information Systems Expectations and Experiences (I-SEE) survey which assessed 5) Provider-patient communication, 6) Inter-provider communication 7) Inter-	Cronbach α was greater than .70. Confirmatory factor analysis was steady with a priori expectations. Descriptive analyses were used to examine characteristics	Key findings: 1) Eight of the 47 survey items decreased significantly from the first survey to the last. 2) 37 survey items decreased significantly from the second survey to the last. 3) Nurses with previous HIT SYSTEM experience expressed more positive responses than nurses with no previous HIT

		survey & last day of training survey 3-month pre-implementation; 6-months post-implementation survey Sample: 1,395 anonymous staff, mostly RNs & LPNs over all 3 survey admins. Response rate: Although it was stated that there was a possible 2,700 employees, the break-down per survey was not stated.	organizational communication 8) Work-life changes 9) Improved care 10) Support & resources 11) Patient care processes Administered online via REDCap.	of job categories, work units & survey responses.	SYSTEM experience. 4) Nurses with more years' experience were less positive of HIT SYSTEM perceptions. Limitations: 1) Study focused mainly on feedback of nurses at a single hospital. 2) Due to use of survey of perceptions, response biases may have been demonstrated.
--	--	--	---	--	--

III. METHODOLOGY

The methodology of the research study is parted into its research design, population, and data collection procedures. Additionally, the suitable data collection instrument is determined based on the research design and population. Applied to the study will be the appropriate data analysis.

A prospective post-implementation survey was used as the research method on the comprehensive HIT system within the facility healthcare system. The intent of the design was to help describe the current views of the healthcare staff in relation to the quality of the system, the implementation and its current operation.

Research was conducted at two acute care hospitals that recently rolled out the EMR system within the last year. The study population was end users of the integrated system within the Continuum of Care departments of acute care hospital sites in Temple, Texas. The first facility is a 64-bed pediatric specialty care and teaching hospital. The second is a 636-bed specialty care and teaching hospital. The health information technology employed was the commercial software system, Epic. The execution of the research study used the direction laid out in Health Informatics Research Methods: Principles and Practice (Layman, 2009).

a) Data Collection Procedures

Data collection was performed by anonymous submission online via REDCap (REDCap, 2009). Notification was given through the employer email system with permission from management. A cover letter was included stating participation was voluntary

and not part of an institutional initiative (Figure 1). After one week, a reminder email was provided to the same staff. At the end of fourteen days, the link to survey was ended.

Figure 1 : Cover Letter introducing Epic System Post-Implementation and Use Assessment Survey

The following survey is intended to help understand staff perceptions and attitudes regarding the quality of the new electronic medical record system, Epic. In addition, the study will conduct a benefits evaluation to appreciate the quality of the information provided by the system, as well as, the level of satisfaction amongst end-users.

****Only respond to this survey if you use the Epic system as part of your usual job responsibilities AND work mainly in a hospital setting.****

Participation in survey is completely voluntary. No Protected Health Information is asked. The survey is being conducted for research purposes only and it is not a Baylor Scott & White institutional initiative. All submissions are anonymous and will be maintained on a secure autonomous website.

The survey will take approximately take 15-20 minutes to complete.
Thank you in advance for your participation.

Please follow the link to access online survey.

<https://pedsredcap.uthsc.edu/redcap/surveys/?s=4q4p5ZVb9S>

Any questions regarding the survey, please send your inquiries to apedigo@sw.org.

b) Data Collection Instrument

Several articles found during the literature review presented instruments that were further evaluated in formulation of a suitable questionnaire for the research study. The data collection instruction employed was shaped from the merging two public surveys: the Health Information Technology Reference-Based Evaluation Framework and the Canada Health Infoway System and Use Assessment Survey (Sockolow, Weiner, Bowles, & Lehmann, 2011) (Canada Health Infoway, 2007) (Figures 2 and 3). Both surveys were available for public use. Neither survey required permission to use in forthcoming studies. The combined survey measured structural quality, quality of information logistics, effects on quality of processes, effects on outcomes and quality of care, unintended consequences or benefits and barriers or facilitators to clinician's adoption (Figure 4).

Figure 2 : Health Information Technology Reference-Based Evaluation Framework

Participant Code: _____

Employee / Staff Perceptions Electronic Health Record System Survey

Instructions to Participants

The following survey is intended to help researchers from Drexel University better understand staff perceptions and attitudes about the quality of clinical documentation. Specifically we are interested in learning about your experience using an electronic health record and understanding what impact an electronic health record has on patient care, and how it affects you.

Please check one (1) response for each question. Thank you for completing this survey.

Page 1 of 3

Participant Code: _____

Your Job Title _____ Date _____

Have you had prior experience outside of your facility with any electronic health records or computerized provider order entry systems? No ____ Yes ____ If yes, about how many years of experience _____

Years working in healthcare _____

Would you rate your computer knowledge as below average; average; above average; advanced? _____

Your Age _____ Gender _____

Please indicate the extent to which you agree with the following statements regarding the electronic health record.

Please check only one (1) response per item.

	Strongly Disagree	Moderately Disagree	Mildly Disagree	Mildly Agree	Moderately Agree	Strongly Agree
1. The <u>electronic health record</u> is consistently available	[]	[]	[]	[]	[]	[]
2. The <u>electronic health record</u> is subject to frequent system problems	[]	[]	[]	[]	[]	[]
3. The <u>electronic health record</u> is user-friendly	[]	[]	[]	[]	[]	[]
4. Sufficient support is available to use the <u>electronic health record</u>	[]	[]	[]	[]	[]	[]
5. The <u>electronic health record</u> features enable me to perform my work well	[]	[]	[]	[]	[]	[]
6. Patient care data being recorded is accurate and valid	[]	[]	[]	[]	[]	[]
7. Patients have concerns about the <u>electronic health record</u> security or confidentiality	[]	[]	[]	[]	[]	[]
8. Patient care services are provided in a timely manner	[]	[]	[]	[]	[]	[]
9. Patient care orders in the <u>electronic health record</u> are appropriate	[]	[]	[]	[]	[]	[]
10. The <u>electronic health record</u> contributes to the safety of patients	[]	[]	[]	[]	[]	[]
11. The <u>electronic health record</u> supports effective communication between most team members about patient care	[]	[]	[]	[]	[]	[]
12. The <u>electronic health record</u> contributes to patient outcomes	[]	[]	[]	[]	[]	[]

Page 2 of 3

Participant Code: _____

	Strongly Disagree	Moderately Disagree	Mildly Disagree	Mildly Agree	Moderately Agree	Strongly Agree
13. The <u>electronic health record</u> contributes to patient's knowledge of their health condition	[]	[]	[]	[]	[]	[]
14. The <u>electronic health record</u> is worth the time and effort required to use it	[]	[]	[]	[]	[]	[]
15. Overall, I am satisfied with the <u>electronic health record</u>	[]	[]	[]	[]	[]	[]
16. I think patients are satisfied with clinicians using the <u>electronic health record</u>	[]	[]	[]	[]	[]	[]
17. My department had a role in introducing the <u>electronic health record</u> at my facility	[]	[]	[]	[]	[]	[]
18. People who use the <u>electronic health record</u> should have had more to say about its design	[]	[]	[]	[]	[]	[]
19. I have first hand knowledge that problems with the <u>electronic health record</u> interfere with patient care	[]	[]	[]	[]	[]	[]
20. A reason for my facility's adoption of the <u>electronic health record</u> was the system's ability to exchange patient information with nursing homes and hospitals	[]	[]	[]	[]	[]	[]
21. Sufficient resources are provided for me to learn to use the <u>electronic health record</u>	[]	[]	[]	[]	[]	[]
22. Part of the increase in costs of healthcare is because of computers	[]	[]	[]	[]	[]	[]
23. What worked well or what are your concerns related to the system:						

Thank you for completing this survey.

Page 3 of 3

Figure 3 : Canada Health Infoway System and Use Assessment Survey

Canada Health Infoway SYSTEM AND USE ASSESSMENT SURVEY	
LOCATION	
DATE	
To Whom It May Concern:	
The Ministry of Health & Long-Term Care (MHLTC) and Canada Health Infoway (CHI) are conducting a benefits evaluation study in order to improve the quality of the information provided by the health information systems, as well as, the level of satisfaction amongst end-users. Your feedback and assistance with this survey will help MHLTC and CHI to develop better systems and deliver better services. The following survey consists of specific questions on: the ease and functionality, information quality, service quality related to CHI health information system implemented at your Hospital or Centre. The survey will take approximately 10-15 minutes to complete. Please circle the response that best represents your opinion. Information that is collected during this survey will be kept anonymous and confidential. Please return the completed survey using the enclosed postage paid self-addressed envelope. If you have any questions about the survey, please contact _____ Thank you in advance for your participation. Sincerely yours, Canada Health Infoway / SPONSOR	
Version Date: March 2007	1

**Canada Health Infoway
SYSTEM AND USE ASSESSMENT SURVEY**

SECTION 1. OVERALL USER SATISFACTION

1. In general, how satisfied are you overall with the system you are currently working with? By "system" we mean, the ease and functionality of the system itself, the quality of the information given and the quality of the services provided for the system.

Highly satisfied ☐ Moderately satisfied ☐ Neither satisfied nor dissatisfied ☐ Moderately dissatisfied ☐ Not at all satisfied ☐

2. Please indicate your level of agreement or disagreement with each of the following statements below.

	Strongly Agree	Moderately Agree	Moderately Disagree	Strongly Disagree	Not Sure	Not Applicable
a.) The system improves my productivity	☐	☐	☐	☐	☐	
b.) The system improves the quality of care I can provide	☐	☐	☐	☐	☐	
c.) The system makes my job easier	☐	☐	☐	☐	☐	
d.) The system enhances our ability to coordinate the continuity of care	☐	☐	☐	☐	☐	
e.) The system improves our sharing of patient information amongst providers	☐	☐	☐	☐	☐	
f.) The system enhances the efficiency of ordering lab tests, X-rays, prescriptions, etc.	☐	☐	☐	☐	☐	☐
g.) The alerts, reminders and order set features (i.e. support tools) improve the quality of my decision-making	☐	☐	☐	☐	☐	☐

3. Are there aspects of the system that you would change, and if so, which ones would they be? Please describe your comments.

4. Do you have any experiences with the system where it has supported the provision of care? Please describe your comments.

Version Date: March 2007

2

**Canada Health Infoway
SYSTEM AND USE ASSESSMENT SURVEY**

SECTION 2. SYSTEM QUALITY

5. Based on your experiences to date with the system, how acceptable is the quality of the system itself (as described by the specific characteristics listed below)? Would you say it is:

Highly acceptable Moderately acceptable Neither acceptable nor unacceptable Moderately unacceptable Not at all acceptable
☐ ☐ ☐ ☐ ☐

6. Please indicate your level of agreement or disagreement with each of the following statements below.

	Strongly Agree	Moderately Agree	Moderately Disagree	Strongly Disagree	Not Sure
a.) The system is easy to use	☐	☐	☐	☐	☐
b.) The response time is acceptable	☐	☐	☐	☐	☐
c.) The system is integrated with my workflow	☐	☐	☐	☐	☐
d.) The system security is acceptable	☐	☐	☐	☐	☐
e.) The system features enable me to perform my work well	☐	☐	☐	☐	☐
f.) The system is reliable in its performance	☐	☐	☐	☐	☐
g.) Overall, the quality of the system is excellent	☐	☐	☐	☐	☐

SECTION 3. INFORMATION QUALITY

7. In general, when thinking about the quality of the information provided by the system, do you find the quality of the information to be:

Highly acceptable Moderately acceptable Neither acceptable nor unacceptable Moderately unacceptable Not at all acceptable
☐ ☐ ☐ ☐ ☐

8. Please indicate your level of agreement or disagreement with each of the following statements below.

	Strongly Agree	Moderately Agree	Moderately Disagree	Strongly Disagree	Not Sure
a.) The information is complete	☐	☐	☐	☐	☐
b.) The information is quickly provided	☐	☐	☐	☐	☐
c.) The information provided is accurate	☐	☐	☐	☐	☐
d.) The information provided is relevant	☐	☐	☐	☐	☐
e.) The information is available when I need it	☐	☐	☐	☐	☐
f.) The format and layout of the information is acceptable	☐	☐	☐	☐	☐

Version Date: March 2007

3

**Canada Health Infoway
SYSTEM AND USE ASSESSMENT SURVEY**

SECTION 4. SERVICE QUALITY

9. In general, when thinking about the quality of the services (i.e. technical support and training services) provided for the system, do you find the quality of these services to be:

Highly acceptable Moderately acceptable Neither acceptable nor unacceptable Moderately unacceptable Not at all acceptable
☐ ☐ ☐ ☐ ☐

10. Please indicate your level of agreement or disagreement with each of the following statements below.

	Strongly Agree	Moderately Agree	Moderately Disagree	Strongly Disagree	Not Sure
a.) The implementation process at this Hospital or Centre was acceptable	☐	☐	☐	☐	☐
b.) The current level of training is acceptable	☐	☐	☐	☐	☐
c.) The level of on-going support provided is acceptable	☐	☐	☐	☐	☐

SECTION 5. PUBLIC HEALTH SURVEILLANCE SPECIFIC

-TO BE COMPLETED BY PUBLIC HEALTH SURVEILLANCE PERSONNEL ONLY -

11. Please indicate your level of agreement or disagreement for each of the following statements below.

	Strongly Agree	Moderately Agree	Moderately Disagree	Strongly Disagree	Not Sure
a.) The system improves the detection and management of reportable diseases	☐	☐	☐	☐	☐
b.) The system improves the management of immunization process	☐	☐	☐	☐	☐

Version Date: March 2007

4

**Canada Health Infoway
SYSTEM AND USE ASSESSMENT SURVEY**

SECTION 6. SYSTEM USAGE

12. In a typical day, how many times do you 'use' the system?

_____ Number of times, a day

Always ☐ Rarely ☐

13. In a typical week, please indicate the number of days in which you use the system.

_____ Number of days, a week

14. Please estimate what percent of your patients do you use the system?

_____ % patients (FILL IN)

Don't know ☐

15. How likely are you to recommend the system to other healthcare providers at other Hospitals or Centres?

Definitely ☐ Probably ☐ May or may not ☐ Probably Not ☐ Definitely not ☐

16. Given a choice, would you like to increase or decrease your future use of the system that you are currently working with? Would that be a significant or moderate increase / decrease, or would you like your future use to stay the same?

Significant Increase	Moderately Increase	Moderately Decrease	Significant Decrease	<u>REMAIN THE SAME</u>
☐	☐	☐	☐	☐

SECTION 7. OTHER COMMENTS

17. Do you have any other comments you would like to make regarding the system ?



**Canada Health Infoway
SYSTEM AND USE ASSESSMENT SURVEY**

SECTION 8. DEMOGRAPHIC INFORMATION

18. What is your profession?

- | | | | |
|------------------------------------|--------------------------|---|--------------------------|
| Administrative support staff | ☐ | Family physician | ☐ |
| Imaging technologist | ☐ | Specialist physician (please specify below) | ☐ |
| Laboratory technician | ☐ | <input type="text"/> | |
| Nurse | ☐ | Other (please specify below) | ☐ |
| Pharmacist | ☐ | <input type="text"/> | |

19. How would you describe your "use" of the system? (Check all that apply)

- | | | | |
|--|--------------------------|---|--------------------------|
| I use the system for clinical decision making | ☐ | I use the system to both access patient information and in clinical decision making | ☐ |
| I use the system to access patient information and support the clinical decision maker | ☐ | | |

20. How long have you been using the system?

- | | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Less than a month | 1-3 months | 4-6 months | 7-12 months | 1-2 years | 3-5 years |
| ☐ | ☐ | ☐ | ☐ | ☐ | ☐ |

21. Currently, how do you receive your patient results?

_____ % FAX _____ % SYSTEM _____ % OTHER (please specify / write below)

22. How would you rate your computer proficiency?

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Basic | Average | Advanced | Expert |
| ☐ | ☐ | ☐ | ☐ | ☐ |

23. Please check the response(s) that best describe the settings where you work.

- | | | | | |
|---|--------------------------|---|-----------|--------------------------|
| Academic / Teaching Hospital | ☐ | a. Do you work within the emergency department? | Yes | ☐ |
| Community Clinic / Health Center | ☐ | | No | ☐ |
| Community Hospital | ☐ | | | |
| Nursing Home / Long Term Care Facility | ☐ | | | |
| Private Office / Clinic | ☐ | | | |
| Other (please specify/write answer below) | ☐ | | | |

24. Where are you located?

- | | | | |
|-----------------------------|--------------------------|----------------------------|--------------------------|
| Alberta | ☐ | Nunavut | ☐ |
| British Columbia | ☐ | Ontario | ☐ |
| Manitoba | ☐ | Prince Edward Island | ☐ |
| New Brunswick | ☐ | Quebec | ☐ |
| Newfoundland | ☐ | Saskatchewan | ☐ |
| Northwest Territories | ☐ | Yukon | ☐ |
| Nova Scotia | ☐ | | |

**THANK YOU FOR YOUR HELP IN IMPROVING THE INFORMATION SYSTEM.
PLEASE RETURN YOUR COMPLETED QUESTIONNAIRE USING THE ENCLOSED, POSTAGE
PAID ENVELOPE.**

Version Date: March 2007

6

Figure 4

Page 1 of 7

Epic System Post-Implementation and Use Assessment Survey

Basic Information

- 1) Select your primary work location:
 - ☐ Baylor Scott & White Brenham Hospital
 - ☐ Baylor Scott & White College Station Hospital
 - ☐ Baylor Scott & White McLane Children's Hospital
 - ☐ Baylor Scott & White Round Rock Hospital
 - ☐ Baylor Scott & White Taylor Hospital
 - ☐ Baylor Scott & White Temple Continuing Care Hospital
 - ☐ Baylor Scott & White Temple Hospital
- 2) Select your profession:
 - ☐ Advanced Practice Staff (i.e. Physician Assistant, Nurse Practitioner, CRNA)
 - ☐ Allied Health Staff (i.e. PT, OT, SLP, Respiratory, Technician, Technologist)
 - ☐ Administrative Support Staff (i.e. Administrative Assistants, Receptionists, Case Management Assistants)
 - ☐ Case Management Staff (i.e. Nurse Case Manager, Social Worker)
 - ☐ Clinical Support Staff (i.e. CNA, HUC)
 - ☐ HIM/Coding Staff (i.e. Claim Adjustment Coordinator, Coding Specialist)
 - ☐ IT Staff (i.e. Application Analyst, Server Engineer)
 - ☐ Nursing Staff (i.e. LVN, RN)
 - ☐ Pharmacist
 - ☐ Physicians (Resident)
 - ☐ Physicians (Fellow)
 - ☐ Physicians (Less than 3 years post-residency)
 - ☐ Physicians (Greater than 3 years post-residency)
 - ☐ Other
- 3) If job title not provided in previous question, please provide: _____
- 4) Select your age range:
 - ☐ 25 or younger
 - ☐ 26 to 35
 - ☐ 36 to 45
 - ☐ 46 to 55
 - ☐ 56 to 65
 - ☐ 66 or older
- 5) How would you rate your computer proficiency?
 - ☐ None
 - ☐ Basic
 - ☐ Average
 - ☐ Advanced
 - ☐ Expert
- 6) Have you had prior experience outside of your facility with any electronic medical record system?
 - ☐ Yes
 - ☐ No

- 7) If answer to previous question is "Yes", how many years experience do you have working with an electronic medical record system?

- ☐ Less than 2 years
☐ 2-5 years
☐ More than 5 years

- 8) How long have you been using the current Baylor Scott & White Epic system?

- ☐ Less than a month
☐ 1-3 months
☐ 4-6 months
☐ 7-11 months
☐ 1-2 years

EPIC SYSTEM QUALITY - Please read each statement and indicate the response that is closest to your belief.

	Strongly Agree	Moderately Agree	Mildly Agree	Mildly Disagree	Moderately Disagree	Strongly Disagree	Not Sure
9) The system is easy to use.	☐	☐	☐	☐	☐	☐	☐
10) The system is reliable in its performance.	☐	☐	☐	☐	☐	☐	☐
11) The system is consistently available.	☐	☐	☐	☐	☐	☐	☐
12) The system's response time is acceptable.	☐	☐	☐	☐	☐	☐	☐
13) The system supports effective communication between team members.	☐	☐	☐	☐	☐	☐	☐
14) The system's ability to exchange patient information with other systems is acceptable.	☐	☐	☐	☐	☐	☐	☐
15) The system has been integrated appropriately with my previous workflows.	☐	☐	☐	☐	☐	☐	☐
16) The system features enable me to perform my work well.	☐	☐	☐	☐	☐	☐	☐
17) The system security is acceptable.	☐	☐	☐	☐	☐	☐	☐
18) Based on your experiences to date with the Epic system, how acceptable is the quality of the system itself as described by the specific characteristics listed above?							☐ Highly Acceptable ☐ Moderately Acceptable ☐ Neither Acceptable nor Unacceptable ☐ Moderately Unacceptable ☐ Not at all Acceptable
19) Comments related to Epic's System Quality (If possible, please provide which question number the comment is related to.)							

EPIC INFORMATION QUALITY - Please read each statement and indicate the response that is closest to your belief.

	Strongly Agree	Moderately Agree	Mildly Agree	Mildly Disagree	Moderately Disagree	Strongly Disagree	Not Sure	
20) The information provided is relevant.	☐	☐	☐	☐	☐	☐	☐	
21) The information provided is accurate.	☐	☐	☐	☐	☐	☐	☐	
22) The information is complete.	☐	☐	☐	☐	☐	☐	☐	
23) The format and layout of the information is acceptable.	☐	☐	☐	☐	☐	☐	☐	
24) The information is available when I need it.	☐	☐	☐	☐	☐	☐	☐	
25) When thinking about the quality of the information provided by Epic in general, how do you find the quality of the information to be?	☐ Highly Acceptable ☐ Moderately Acceptable ☐ Neither Acceptable nor Unacceptable ☐ Moderately Unacceptable ☐ Not at all Acceptable						☐	
26) Comments related to Epic's Information Quality (If possible, please provide which question number the comment is related to.)								☐

EPIC SERVICE QUALITY - Please read each statement and indicate the response that is closest to your belief.

	Strongly Agree	Moderately Agree	Mildly Agree	Mildly Disagree	Moderately Disagree	Strongly Disagree	Not Sure	
27) The implementation process at my facility was acceptable.	☐	☐	☐	☐	☐	☐	☐	
28) The current level of training at my facility is acceptable.	☐	☐	☐	☐	☐	☐	☐	
29) The level of on-going support provided at my facility is acceptable.	☐	☐	☐	☐	☐	☐	☐	
30) When thinking about the quality of the services (i.e. technical support and training services) provided for Epic in general, how do you find the quality of these services to be?	☐ Highly Acceptable ☐ Moderately Acceptable ☐ Neither Acceptable nor Unacceptable ☐ Moderately Unacceptable ☐ Not at all Acceptable						☐	
31) Comments related to Epic's Service Quality (If possible, please provide which question number the comment is related to.)								☐

EPIC CLINICAL QUALITY - Please read each statement and indicate the response that is closest to your belief.

	Strongly Agree	Moderately Agree	Mildly Agree	Mildly Disagree	Moderately Disagree	Strongly Disagree	Not Sure
32) The system contributes to improved patient outcomes.	☐	☐	☐	☐	☐	☐	☐
33) The system contributes to the safety of the patients.	☐	☐	☐	☐	☐	☐	☐
34) The system contributes to the patient's knowledge of their health condition.	☐	☐	☐	☐	☐	☐	☐
35) The patients are satisfied with the clinicians using the system.	☐	☐	☐	☐	☐	☐	☐
36) The patients have concerns about the system's security and confidentiality.	☐	☐	☐	☐	☐	☐	☐
37) The patient care data being recorded is accurate and valid.	☐	☐	☐	☐	☐	☐	☐
38) The patient care services are able to be provided in a <u>more</u> <u>timely</u> manner.	☐	☐	☐	☐	☐	☐	☐
39) The selection of patient care orders in system is appropriate.	☐	☐	☐	☐	☐	☐	☐
40) The system contributes to improved clinical documentation.	☐	☐	☐	☐	☐	☐	☐
41) Based on your experiences to date with the Epic system, how acceptable is the clinical data of the system itself as described by the specific characteristics listed above?	☐ Highly Acceptable ☐ Moderately Acceptable ☐ Neither Acceptable nor Unacceptable ☐ Moderately Unacceptable ☐ Not at all Acceptable						
42) Comments related to Epic's Clinical Quality (If possible, please provide which question number the comment is related to.)							

GENERAL COMMENTS

- 43) What specific features of Epic are especially appreciated?
- 44) What specific aspects of Epic could be improved on by the vendor?
- 45) Do you have any lessons learned since the Epic system implementation?
- 46) Do you have additional goals related to Epic that you or your department have not yet completed?
- 47) Have there been any unexpected benefits gained for your department or the organization since implementing Epic?

c) Data Analysis

Statistical software, SPSS, was utilized to create various types of statistical analyses, including descriptive statistics such as the standard deviation to responses. Furthermore, descriptive analysis was used to examine characteristics of survey responses (IBM SPSS Statistics, 2013).

IV. RESULTS

The following results describe the response rate and break down the demographics of the respondents.

a) Response Rate of Population

The response rate was determined to be 37.78%. One hundred seven possible respondents were

emailed a cover letter and link to the autonomous website. Again, one week later the same cover letter and link were emailed to the same one hundred and seven staff members. The link was terminated one week later. In total, thirty-four valid surveys were completed.

b) Representativeness of Population

The staff ranged in age from younger than twenty-five to greater than sixty-six. The largest number of respondents was present in the fifty-six to sixty-five year age range (32.4%). The majority stated their computer proficiency as average (61.8%) and had prior EMR experience (55.9%). (Figure 5, 6 & 7; Table 3).

Figure 5 : Pie Chart of Age Range





Figure 6 : Computer Proficiency Frequency






Figure 7 : Prior EMR Experience Frequency

Table 3 : Staff Demographics

Profession

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Administrative Support Staff	2	5.9	5.9	5.9
	Case Management Staff	29	85.3	85.3	91.2
	Other	3	8.8	8.8	100.0
	Total	34	100.0	100.0	

Age Range

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	25 or younger	3	8.8	8.8	8.8
	26 to 35	4	11.8	11.8	20.6
	36 to 45	9	26.5	26.5	47.1
	46 to 55	4	11.8	11.8	58.8
	56 to 65	11	32.4	32.4	91.2
	66 or older	3	8.8	8.8	100.0
	Total	34	100.0	100.0	

Computer Proficiency

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Basic	4	11.8	11.8	11.8
	Average	21	61.8	61.8	73.5
	Advanced	8	23.5	23.5	97.1
	Expert	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

Prior EMR Experience

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	19	55.9	55.9	55.9
	No	15	44.1	44.1	100.0
	Total	34	100.0	100.0	

Years w/ EMR Experience

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Less than 2 years	5	14.7	23.8	23.8
	2-5 years	7	20.6	33.3	57.1
	More than 5 years	9	26.5	42.9	100.0
	Total	21	61.8	100.0	
Missing	System	13	38.2		
	Total	34	100.0		

Current Baylor Scott & White Epic Experience

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Less than a month	1	2.9	2.9	2.9
	1-3 months	1	2.9	2.9	5.9
	4-6 months	1	2.9	2.9	8.8
	7-11 months	26	76.5	76.5	85.3
	1-2 years	5	14.7	14.7	100.0
	Total	34	100.0	100.0	

c) Research Questions

In developing an understanding of the attitude of the staff, the quality of the system, its information and the service provided regarding the HIT system were measured. Additionally, the particular aspects of the clinical data were analyzed. A five-level Likert scale was utilized to measure the employee's stance on the quality of the HIT system, the information within the HIT system, the service provided to support the HIT system and particular aspects related to the clinical information of the HIT system.

In regard to the quality of the system, a majority of the staff strongly agree that the system is consistently available (47.1%) and has acceptable security (50%). As for the system appropriately integrating with previous workflows, the employees were mostly divided between mildly agree (26.5%), moderately agree (29.4%) and strongly agree (29.4%). None of workers disagreed in a majority to any of the aspects measured related the quality of the system. The remainder moderately agreed that the system was easy to use (70%), its performance was reliable (44.1%), had acceptable response time (47.1%), provided effective communication between team members (41.2%), had acceptable exchange of information with other systems (38.2%) and enabled staff to perform work well (38.2%). (Figure 8; Table 3)

Figure 8




Table 4 : Epic System Quality

System - Easy to Use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	6	17.6	17.6	17.6
	Moderately Agree	24	70.6	70.6	88.2
	Mildly Agree	2	5.9	5.9	94.1
	Mildly Disagree	2	5.9	5.9	100.0
	Total	34	100.0	100.0	

System - Reliable Performance

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	13	38.2	38.2	38.2
	Moderately Agree	15	44.1	44.1	82.4
	Mildly Agree	5	14.7	14.7	97.1
	Mildly Disagree	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

System - Consistently Available

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	16	47.1	47.1	47.1
	Moderately Agree	14	41.2	41.2	88.2
	Mildly Agree	3	8.8	8.8	97.1
	Mildly Disagree	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

System - Acceptable Response Time

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	14	41.2	41.2	41.2
	Moderately Agree	16	47.1	47.1	88.2
	Mildly Agree	3	8.8	8.8	97.1
	Moderately Disagree	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

System - Effective Communication b/t Team Members

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	11	32.4	32.4	32.4
	Moderately Agree	14	41.2	41.2	73.5
	Mildly Agree	7	20.6	20.6	94.1
	Mildly Disagree	2	5.9	5.9	100.0
	Total	34	100.0	100.0	

System - Acceptable Exchange Information w/ Other Systems

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	8	23.5	23.5	23.5
	Moderately Agree	13	38.2	38.2	61.8
	Mildly Agree	6	17.6	17.6	79.4
	Mildly Disagree	2	5.9	5.9	85.3
	Moderately Disagree	2	5.9	5.9	91.2
	Strongly Disagree	1	2.9	2.9	94.1
	Not Sure	2	5.9	5.9	100.0
	Total	34	100.0	100.0	

System - Integrated Appropriately w/ Previous Workflows

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	10	29.4	29.4	29.4
	Moderately Agree	10	29.4	29.4	58.8
	Mildly Agree	9	26.5	26.5	85.3
	Mildly Disagree	1	2.9	2.9	88.2
	Moderately Disagree	2	5.9	5.9	94.1
	Not Sure	2	5.9	5.9	100.0
	Total	34	100.0	100.0	

System - Enables Staff to Perform Work Well


		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	12	35.3	35.3	35.3
	Moderately Agree	13	38.2	38.2	73.5
	Mildly Agree	8	23.5	23.5	97.1
	Not Sure	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

System - Acceptable System Security

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	17	50.0	50.0	50.0
	Moderately Agree	10	29.4	29.4	79.4
	Mildly Agree	4	11.8	11.8	91.2
	Not Sure	3	8.8	8.8	100.0
	Total	34	100.0	100.0	

The criteria measured related to the system's information was mostly seen as moderately agreeable. A majority of the staff moderately agree that the information is accurate (52.9%), relevant (47.1%), complete (47.1%)

and has an acceptable layout (41.2%). An even number moderately agrees (41.2%) as strongly agree (41.2%) that the information is available when needed. (Figure 9; Table 4)

Figure 9*Table 4 : Epic Information Quality**Information - Relevant*

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	14	41.2	41.2	41.2
	Moderately Agree	16	47.1	47.1	88.2
	Mildly Agree	4	11.8	11.8	100.0
	Total	34	100.0	100.0	

Information - Accurate

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	12	35.3	35.3	35.3
Moderately Agree	18	52.9	52.9	88.2
Mildly Agree	3	8.8	8.8	97.1
Not Sure	1	2.9	2.9	100.0
Total	34	100.0	100.0	

Information - Complete

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	11	32.4	32.4	32.4
Moderately Agree	16	47.1	47.1	79.4
Mildly Agree	5	14.7	14.7	94.1
Mildly Disagree	1	2.9	2.9	97.1
Not Sure	1	2.9	2.9	100.0
Total	34	100.0	100.0	

Information - Acceptable Layout

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	13	38.2	38.2	38.2
Moderately Agree	14	41.2	41.2	79.4
Mildly Agree	7	20.6	20.6	100.0
Total	34	100.0	100.0	

Information - Available When Needed

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	14	41.2	41.2	41.2
Moderately Agree	14	41.2	41.2	82.4
Mildly Agree	4	11.8	11.8	94.1
Mildly Disagree	2	5.9	5.9	100.0
Total	34	100.0	100.0	

In the three characteristics of service measured, a majority of staff moderately agreed that the implementation process (55.9%), level of training (47.1%) and on-going support (47.1%) is acceptable. (Figure 10; Table 5)

Figure 10




Table 5 : Epic Service Quality

Service - Acceptable Implementation Process

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	7	20.6	20.6	20.6
	Moderately Agree	19	55.9	55.9	76.5
	Mildly Agree	3	8.8	8.8	85.3
	Mildly Disagree	2	5.9	5.9	91.2
	Moderately Disagree	1	2.9	2.9	94.1
	Strongly Disagree	1	2.9	2.9	97.1
	Not Sure	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

Service - Acceptable Level of Training

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	6	17.6	17.6	17.6
	Moderately Agree	16	47.1	47.1	64.7
	Mildly Agree	8	23.5	23.5	88.2
	Mildly Disagree	2	5.9	5.9	94.1
	Moderately Disagree	1	2.9	2.9	97.1
	Strongly Disagree	1	2.9	2.9	100.0
	Total	34	100.0	100.0	

Service - Acceptable On-Going Support

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Strongly Agree	10	29.4	29.4	29.4
	Moderately Agree	16	47.1	47.1	76.5
	Mildly Agree	8	23.5	23.5	100.0
	Total	34	100.0	100.0	

Because most of the respondents do not work directly with the patients, the majority answered that they were not sure of the patient's satisfaction with clinicians' use of system (35.3%) or patient's concerns with system security and confidentiality (41.2%). A majority strongly believe that the clinical data has improved patient outcomes (41.2%), improved patient safety (41.2%),

improved patient's knowledge of their health (38.2%) and improved clinical documentation (38.2%). A majority moderately believe the clinical data of the patient is accurate and valid (44.1%), the timely manner of the patient care services has increased (35.3%) and that there is an appropriate selection of patient care orders (35.3%). (Figure 11; Table 6)

Figure 11




Table 6 : Epic Clinical Quality

Improved Patient Outcomes

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	14	41.2	41.2	41.2
Moderately Agree	11	32.4	32.4	73.5
Mildly Agree	6	17.6	17.6	91.2
Not Sure	3	8.8	8.8	100.0
Total	34	100.0	100.0	

Improved Safety of Patient

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	14	41.2	41.2	41.2
Moderately Agree	12	35.3	35.3	76.5
Mildly Agree	6	17.6	17.6	94.1
Not Sure	2	5.9	5.9	100.0
Total	34	100.0	100.0	

Improved Knowledge of Health by Patients

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	13	38.2	38.2	38.2
Moderately Agree	10	29.4	29.4	67.6
Mildly Agree	4	11.8	11.8	79.4
Mildly Disagree	2	5.9	5.9	85.3
Not Sure	5	14.7	14.7	100.0
Total	34	100.0	100.0	

Patient Satisfied w/ Clinicians Use of System

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	7	20.6	20.6	20.6
Moderately Agree	8	23.5	23.5	44.1
Mildly Agree	6	17.6	17.6	61.8
Mildly Disagree	1	2.9	2.9	64.7
Not Sure	12	35.3	35.3	100.0
Total	34	100.0	100.0	

Concerns w/ Security & Confidentiality by Patients

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	4	11.8	11.8	11.8
Moderately Agree	2	5.9	5.9	17.6
Mildly Agree	3	8.8	8.8	26.5
Mildly Disagree	6	17.6	17.6	44.1
Moderately Disagree	3	8.8	8.8	52.9
Strongly Disagree	2	5.9	5.9	58.8
Not Sure	14	41.2	41.2	100.0
Total	34	100.0	100.0	

Patient Care Data is Accurate and Valid

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	9	26.5	26.5	26.5
Moderately Agree	15	44.1	44.1	70.6
Mildly Agree	6	17.6	17.6	88.2
Not Sure	4	11.8	11.8	100.0
Total	34	100.0	100.0	

Timely Manner of Patient Care Services Increased

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	9	26.5	26.5	26.5
Moderately Agree	12	35.3	35.3	61.8
Mildly Agree	5	14.7	14.7	76.5
Mildly Disagree	1	2.9	2.9	79.4
Not Sure	7	20.6	20.6	100.0
Total	34	100.0	100.0	

Appropriate Selection of Patient Care Orders

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	9	26.5	26.5	26.5
Moderately Agree	12	35.3	35.3	61.8
Mildly Agree	7	20.6	20.6	82.4
Not Sure	6	17.6	17.6	100.0
Total	34	100.0	100.0	

Improved Clinical Documentation

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Strongly Agree	13	38.2	38.2	38.2
Moderately Agree	12	35.3	35.3	73.5
Mildly Agree	4	11.8	11.8	85.3
Mildly Disagree	1	2.9	2.9	88.2
Not Sure	4	11.8	11.8	100.0
Total	34	100.0	100.0	

The standard deviation of the criteria within the four quality themes were calculated and presented within Table 7. Within Table 8, cross tabulations are provided based on prior EMR experience vs. each of the acceptability of quality of the system, information, service and clinical data. The number of staff with prior EMR experience (N=19) slightly outnumbered the staff with no prior experience (N=15). Having experience with an EMR system or having no experience did not appear to affect the acceptability. In the four measures, the respondents in both groups found the quality of the system, its information, its service and specifically the clinical area all moderately acceptable.

Table 7 : Mean & Standard Deviations of System, Information, Service and Clinical Quality Measurements
Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
System - Easy to Use	34	1	4	2.00	.696
System - Reliable Performance	34	1	4	1.82	.797
System - Consistently Available	34	1	4	1.68	.768
System - Acceptable Response Time	34	1	5	1.76	.855
System - Effective Communication b/t Team Members	34	1	4	2.00	.888
System - Acceptable Exchange Information w/ Other Systems	34	1	7	2.65	1.668
System - Integrated Appropriately w/ Previous Workflows	34	1	7	2.50	1.581
System - Enables Staff to Perform Work Well	34	1	7	2.03	1.167
System - Acceptable System Security	34	1	7	2.06	1.705
Information - Relevant	34	1	3	1.71	.676
Information - Accurate	34	1	7	1.88	1.094
Information - Complete	34	1	7	2.03	1.167
Information - Acceptable Layout	34	1	3	1.82	.758
Information - Available When Needed	34	1	4	1.82	.869
Service - Acceptable Implementation Process	34	1	7	2.35	1.390
Service - Acceptable Level of Training	34	1	6	2.38	1.129
Service - Acceptable On-Going Support	34	1	3	1.94	.736
<i>Descriptive Statistics</i>					
	N	Minimum	Maximum	Mean	Std. Deviation
Clinical - Improved Knowledge of Health by Patients	34	1	7	2.59	2.047
Clinical - Patient Satisfied w/ Clinicians Use of System	34	1	7	3.79	2.508
Clinical - Concerns w/ Security & Confidentiality by Patients	34	1	7	4.88	2.185
Clinical - Patient Care Date is Accurate and Valid	34	1	7	2.50	1.796
Clinical - Timely Manner of Patient Care Services Increased	34	1	7	2.97	2.209
Clinical - Appropriate Selection of Patient Care Orders	34	1	7	2.82	2.081
Clinical - Improved Clinical Documentation	34	1	7	2.38	1.875
Valid N (listwise)	34				

Table 8 : Cross Tabulations

*Acceptability of the Quality of the Epic System * Prior EMR Experience*

Count

		Prior EMR Experience		Total
		Yes	No	
Acceptability of the Quality of the Epic System	Highly Acceptable	7	7	14
	Moderately Acceptable	9	7	16
	Neither Acceptable nor Unacceptable	2	1	3
	Moderately Unacceptable	1	0	1
Total		19	15	34

*Acceptability of the Quality of the Information Provided in Epic * Prior EMR Experience*

Count

		Prior EMR Experience		Total
		Yes	No	
Acceptability of the Quality of the Information Provided in Epic	Highly Acceptable	8	8	16
	Moderately Acceptable	10	6	16
	Neither Acceptable nor Unacceptable	1	1	2
	Total	19	15	34

*Acceptability of the Quality of the Services Provided for Epic * Prior EMR Experience*

Count

		Prior EMR Experience		Total
		Yes	No	
Acceptability of the Quality of the Services Provided for Epic	Highly Acceptable	7	4	11
	Moderately Acceptable	7	7	14
	Neither Acceptable nor Unacceptable	3	2	5
	Moderately Unacceptable	2	2	4
Total		19	15	34

*Acceptability of the Clinical Data within Epic * Prior EMR Experience*

Count

		Prior EMR Experience		Total
		Yes	No	
Acceptability of the Clinical Data within Epic	Highly Acceptable	7	5	12
	Moderately Acceptable	8	8	16
	Neither Acceptable nor Unacceptable	4	2	6
	Total	19	15	34

From the four core categories, each quality set's comment section was reviewed for common themes applicable to productive transition of HIT systems. Within the system quality focus, interoperability between modules within the system and to other systems is a noted concern of staff. As one respondent stated, "communication in the system is available but isn't utilized as well as possible." Another staffer mentions that the system "doesn't consistently interface properly with Midas." (Figure 12) For the information quality, an

issue raised was the inability to access information. The view of a Case Manager is different than that of a nurse which brings concern that information is not being interpreted in the same manner (Figure 13).

Figure 12

Comments related to Epic System Quality

- ❖ Signed and held orders for patient status are being released by physician and nursing personnel changing the patient status to an incorrect status.
- ❖ Doesn't consistently interface properly with Midas. 2. Communication in the system is available but isn't utilized as well as possible. 3. The finance billing system is flawed. It can't register the payroll deduction payment plan that is in effect. The billing also can't automatically roll different cases charges into the main guarantor account so the accounts can register as payment in progress.
- ❖ Would like a better way to print out MAR without explanation of how to administer meds. Would like a more compact MAR. Would like a better way to print several days' worth of vital signs. Under CM snapshot- adult vitals last day is perfect except that it only shows the last day instead of several days.
- ❖ I sit at a desk in front of the computer all day long in a key code locked office so there's no traffic. No patients that come through or anything. It would be nice if a warning box popped up that was big and clear that Epic was going to shut down in 60 seconds so that I could click on it and keep it open. It is frustrating to be working on a case in Midas and have Epic go down all the time and have to keep logging in when I've been sitting in front of the computer the whole time. The other thing is that it's not very clear what dates you're looking at for labs under the overview tab. You really have to concentrate where you're at for the dates. It puts it in a 24 hour period but having the lab values put in rows under a particular date would be more helpful.
- ❖ The EPIC system has streamlined our work time and has been very easy to use.

Figure 13

Comments related to Epic Information Quality

- ❖ The medical record is compartmentalized and groups of people have access to limited information. This can be a communication issue between units such as Case Management or UR and the RCO for billing purposes or the communication between Nursing and Case Management in the discharge process.
- ❖ This is largely dependent upon the quality of documentation by health care providers-not an EPIC issue per se.
- ❖ Some information not reflected at times- delayed.

The largest numbers of concern are in relation to the service quality. One of the concerns is the training was not specific enough for particular job titles. An example given was a class attended by a Case Manager but included staff from the Admissions department. The class was taught using a task list for the Admissions department which was a "different view and way to enter" the system's authorization module. The Case Management felt the "class was not tailored enough" for their department. The same concern was noted by a staff member who not employed during the

implementation but came after. She felt the training was inappropriate for her job description. Along the lines of training, it was mentioned for "more training services on over all process of Epic flow of documentation of a patient." (Figure 14) The staff seems to be unsure of how the system's modules are interconnected. Lastly, concerns were stated in regard to the timeliness of resolving issues. "IT is slow to respond and resolve issues when they arise" was the comment of one employee. For the final quality measure, the statements

indicated that the staff was unsure because they did not deal with patients directly (Figure 15).

Figure 14

Comments related to Epic Service Quality

- ❖ Now that Central TX Region is all on EPIC I anticipate the quality of service will increase and be timelier.
- ❖ Need classes addressing case management and utilization process. Need more training services on over all process of Epic flow of documentation of a patient. I did have a tech come over and he was very helpful. Many classes were set up for certain departments. While other departments did not have but very little training or understanding of how to work in Epic. The Tip sheets were very helpful and the one on one tech support was very supportive.
- ❖ When first taking EPIC training, it did not relate to what we do here. Many questions and frustrations expressed in classes and for a few months after. Today it seems to run ok
- ❖ IT is slow to respond and resolve issues when they arise.
- ❖ The training received was too general. Multiple job titles in the same class and we all have a different view of EPIC. For example, the class I attended on authorization/pre-certification included admissions department. They have a different view and way to enter the auth/cert screen than I do but the class was taught using their worklist. UR does not use that worklist; therefore, the class was not tailored enough for us.
- ❖ I wasn't here for the implementation of Epic. The training for me was for what a case manager or social worker does. It didn't apply to my job whatsoever. Not even a little bit. Everything I learned for Epic is what my co-workers taught me. And all the computer help I get is mainly from one nurse in my office who helps us -- she's very bright and can navigate very well around the Epic system.

Figure 15

Comments related to Epic Clinical Quality

- ❖ I work in an office. I don't work with patients anymore so I am not able to contribute to these questions.
- ❖ I don't know how the patient's feel about the EPIC system
- ❖ Not all patients have access or knowledge to access EPIC in the home environment. Appointments without f/u telephone calls or written notice are frequently missed.

The survey concluded with more general questions related to the implementation process. The topics mentioned by the staff tended to reflect the appropriate training of staff with statements such as "training should have been more specific to my job" and "educate staff thoroughly to obtain the best results. Benefits stated my respondents were more in relation to the system such as "work flow is improved" and "f aster easier access to information." (Figure 16)

Figure 16

Survey General Comments

What specific features of Epic are especially appreciated?

- ❖ Able to complete documentation more efficiently. Documentation is readily available to be viewed by all disciplines. Finding physician orders, labs, demographic information, medical notes, etc. is much easier to access.
- ❖ The fact that you can review clinic notes and in-hospital notes to follow a patient.
- ❖ Everything is in one system. No longer do I have to go into various systems to find notes from various professionals.
- ❖ Electronic is great
- ❖ Clinical documentation is all inclusive within the Epic system.
- ❖ Timeliness of reports being available
- ❖ if notes are in the computer- they are available and do not have to hunt chart Labs and imaging results are faster to view
- ❖ icons that populate the patient list to indicate orders, consults , etc.
- ❖ all data and patient stays in one location - ability to use filters to only see what I need
- ❖ easy to chart and read documents
- ❖ Documentation of all areas in one place. Easy access to previous admissions.
- ❖ Scheduling
- ❖ Note Documents MAR manage orders

What specific aspects of Epic could be improved by the vendor?

- ❖ The access to all clinical notes in the same place.
- ❖ The work queues require more adequate routing rules or setting the rules correctly
- ❖ when need to print MARS or vitals- would be helpful to have a more concise form to print
- ❖ make faxing to skilled nursing facilities available through the system
- ❖ better way to print MAR and Vitals
- ❖ Bringing in notes from previous EMR
- ❖ It would be helpful to have the capability of keeping EPIC up for longer periods of time for those of us working 12 hour shifts in front of a computer. When it times out every ten minutes or so-it creates a significant waste of time from a manpower perspective.
- ❖ Have a warning box pop up in the middle of the screen that gives you 60 seconds before shutting down. Have the labs in one date order at a time under the overview tab-not a 24 hour block of time from yesterday at 0700 to today.
- ❖ Sticky Notes
- ❖ How can user identify salient points related to the work concentration to use effectively in her work on her day of review? I.e. not accomplished and needs to be done.

Do you have any lessons learned since the Epic system implementation?

- ❖ Educate staff thoroughly to obtain the best results.
- ❖ More hands-on training and less classroom lectures would be helpful. I really did not learn much until I actually began charting in the system.
- ❖ training should have been more specific to my job
- ❖ Before EPIC teaches classes-- they need to be prepared for Q&A; who do go to and f/u person for us to contact or that email will be sent out
- ❖ Yes! Auto search is not always the most helpful when scheduling appointments
- ❖ Learned to Navigate thru the system effectively

Do you have additional goals related to Epic that you or your department have not yet completed?

- ❖ I am still working on report writing to establish departmental metrics for CM, UR and ACS/ER.
- ❖ to better navigate the pre hospital encounters
- ❖ Still working with EPIC staff to optimize usage.
- ❖ Identify patients on Facesheet that patient needs items completed

Have there been any unexpected benefits gained for your department of the organization since implementing Epic?

- ❖ Increased documentation and more thorough information documented.
- ❖ My supervisor is able to track the number of consults put into epic.
- ❖ From a financial standpoint able to follow the billing process more adequately.
- ❖ finding info quicker and can view from anywhere-- not just where the chart had been located < or Not>
- ❖ Being able to have an discharge assistants consults queue
- ❖ Work flow is improved.
- ❖ Faster easier access to information

V. DISCUSSION

The significance of the results continues to help develop critical elements necessary for a successful transition to a new comprehensive system. The study focused on the end users' beliefs regarding the quality of the system and particularly, its information and service. Areas of enhancement were revealed included improving training specific to job roles and supplying more fitting integration of processes and workflows. Likewise, confirmatory aspects of current procedures were observed throughout the study. After the implementation, a greater part of the respondents appreciated many of the aspects of having the new technology such as the ease of use, the ability to access to documents within one system and timeliness of information.

Key limitations of the study should to be underscored. The study was conducted at two

associated healthcare facilities located in one city in central Texas. Moreover, the questionnaire was limited to responses from same type of department within the two hospitals. The responses were limited to staff that do not have access to patient care as a routine part of their job responsibilities. Lastly, the fear of participating in survey may have limited the response. Disbelief in true anonymity may have limited or swayed respondents in their scoring or comments.

The resulting recommendations are focused on fostering staff engagement. Taking guidance from a lecture presented by Rod Brace (2014), "The Science of Engagement", engagement is correlated to making progress. As part of progress, there needs to be clarity of goals, a feasible challenge and feedback on actions. But

to make progress, staff will need motivation. Motivation is provided by allowing choices, knowledge and connection to the progress.

As an illustration, the barrier of providing job-specific training could be tackled. Addressing the goal of job-specific training would acknowledge the staff concerns. Providing acknowledgement and recognizing the concerns will engage the personnel. Respond quickly with a plan of action will continue the commitment. Finally, provided feedback will continue the support of a positive transition.

In close, understand the critical elements to support positive HIT transitions are essential but the continued engagement of end users is also vital. Before, during and after implementation, healthcare personnel need to feel competent and related to the transition. Two future studies are recommended. First, a study could be developed to correlate staff engagement to positive HIT changeovers. The second would still be covering the gap in present research which continues to be the need for greater variety of positions giving feedback.

VI. CONCLUSIONS

The subsequent conclusions and recommendations will provide a summary of findings. Along with the findings, conclusions related to the implications to a positive implementation process related to the study and previous studies are provided.

The participants were employed within the Continuum of Care departments of two acute care inpatient facilities. The majority of respondents declared themselves to be Case Management staff. This group includes RN Case Managers and Social Workers. The remaining staff was administrative support staff or management staff of the Continuum of Care departments.

The quality of the four areas of focus all was seen in a largely positive light. Over eighty percent of the respondents moderately or strongly agreed that the system was easy to use, had reliable performance, was consistently available and had an acceptable response time. While acceptable response time did have a ninety-seven percent positive response, one staffer did moderately disagree. Two other areas did contain responses that ranged from strongly agree to moderately or strongly disagree which were the acceptability of information exchange with other systems and the appropriate integration of previous workflows.

As the system information as a whole and the clinical information surveyed individually, the workers replied a mostly affirmative response or stated that they were unsure. Most felt the information was relevant, accurate and had an acceptable layout. A small minority mildly disagreed the information was complete (2.9%) or

available when needed (6.9%). Within the clinical quality survey questions, the response of "Not Sure" was selected than any of the four quality specific areas. From the comments given by the respondents, this was due to the staff not working directly with the patients. Still, a majority strongly believed that the system had provided improved patient outcomes, patient safety, patient knowledge of their health and improved clinical documentation.

While the quality of service still received mostly agreeable responses, it provided the large number of comments of concern by the respondents. Although the majority of survey takers moderately agree the implementation process, level of training and on-going supports were acceptable, the three questions also had responses that included mild, moderate and strong disagreement. The primary issue noted appeared to be centered on job-specific training. Whereas the remarks did convey a desire to better understand the overall process of Epic, the many staff members mentioned the need for training related to "addressing case management." One employee mentioned that there were "many questions and frustrations expressed in classes and for a few months after" because "when (the staff) first took Epic training, it did not relate to what they did." (Figures 5 and 6).

Similar to previous studies, some of the same topics were observed in this study. As with other studies, the implementation process appeared to provide a mostly encouraging transition with a small number of components noted of concern to the staff. Similar to the study in "Transitioning from a computerized provider order entry and paper documentation system to an electronic health record: Expectations and experiences of hospital staff", positive characteristics observed included the quality and safety of patient care. Readily available all-inclusive clinical documentation and the ability to locate patient demographic information quickly were additional benefits of transitioning.

Moreover, conceivable enhancements for future implementations were illustrated with the recent study. One feature of greater apprehension was highlighted by staff with two other concerns of smaller notation. As mentioned in the article "Learning from Within to Ensure a Successful Implementation of an Electronic Health Record", the few of the staff within the current study expressed the similar need for further attention to processes and workflows within the new HIT system. Another minor concern was improving the exchange of information with other systems. More than an ability that can be imparted to the staff during the transition process, the implementation of this the element may be a requirement on the quality of the system itself. The

greatest concern appears to be appropriate staff training. While an understanding of the overall structure of Epic is wanted, a focus on more job-specific training was repeatedly articulated. In summary, the critical elements essential for a successful transition emerging from the study appear to include appropriate training, attention to incorporating processes and workflows, swift feedback to questions and concerns and attention to the staff impression and opinion regarding the HIT system and its implementation.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Brace, Rod (2014). The Science of Engagement. Present at the Memorial Hermann Case Management Retreat. April 11, 2014.
2. Canada Health Infoway (2007). Canada Health Infoway System And Use Assessment Survey. Retrieved from <http://healthit.ahrq.gov/health-it-tools-and-resources/health-it-survey-compendium/canada-health-infoway-system-and-use>.
3. *Crossing the Quality Chasm: A New Health System for the 21st Century*. (2001). The National Academies Press.
4. Department of Health and Human Services: Health Information Technology: Initial Set. (2010). Federal Register, 75(144), 44590-44654. Retrieved from <http://www.gpo.gov/fdsys/pkg/FR-2010-07-28/pdf/2010-17210.pdf>
5. IBM SPSS Statistics (2013). Statistics Coach. Retrieved from Help section of IBM SPss Statistics.
6. Kirkendall, E. S., Goldenhar, L. M., Simon, J. L., Wheeler, D. S., & Andrew Spooner, S. (2013). Transitioning from a computerized provider order entry and paper documentation system to an electronic health record: Expectations and experiences of hospital staff. *International Journal of Medical Informatics*.
7. Laramee, A. S., Bosek, M., Kasprisin, C. A., & Powers-Phaneuf, T. (2012). Learning From Within to Ensure a Successful Implementation of an Electronic Health Record. *CIN: Computers, Informatics, Nursing*, 30, TC181-190.
8. Laramee, A. S. M. A.-B., Bosek, M. D. R. N., Shaner-McRae, H. D. R. F., & Powers-Phaneuf, T. M. R. (2012). A Comparison of Nurse Attitudes Before Implementation and 6 and 18 Months After Implementation of an Electronic Health Record. *CIN: Computers, Informatics, Nursing*, 30(10), 521-530.
9. Layman, E., & Watzlaf, V. (Eds.). (2009). *Health informatics research methods: Principles and practice*. Chicago, IL: American Health Information Management Association.
10. Paul A. Harris, Robert Taylor, Robert Thielke, Jonathon Payne, Nathaniel Gonzalez, Jose G. Conde, Research electronic data capture (REDCap) - A metadata-driven methodology and workflow process for providing translational research informatics support, *J Biomed Inform.* 2009 Apr;42(2):377-81.
11. Sockolow, P. S., Weiner, J. P., Bowles, K. H., & Lehmann, H. P. (2011). A new instrument for measuring clinician satisfaction with electronic health records. *Comput Inform Nurs*, 29(10), 574-585. doi: 10.1097/NCN.0b013e31821a1568
12. Spetz, J., Burgess Jr, J. F., & Phibbs, C. S. (2012). What determines successful implementation of inpatient information technology systems? *American Journal of Managed Care*, 18(3), 157-162.
13. Ward, M. M., Vartak, S., Schwichtenberg, T., & Wakefield, D. S. (2011). Nurses' Perceptions of How Clinical Information System Implementation Affects Workflow and Patient Care. *Computers Informatics Nursing*, 29(9), 502-511 10.1097/NCN. 1090b1013e31822b38798.



This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Polycythemia with Cerebellar Hemangioblastoma

By Elizabeth JeyaVardhini Samuel, Nagarajan Natarajan, Ramesh & Latha M

Pondicherry Institute of Medical Sciences, India

Abstract- 56 year old male presented with one month history of vomiting with abdominal pain; hemogram suggested polycythemia with hemoglobin of 16.5-18.1gm%, gastroscopy showed pan gastritis, bone marrow smear, biopsy showed erythroid, megakaryocytic hyperplasia;ultra sonogram abdomen, pelvis did not show any significant abnormality; patient improved with parenteral hydration, proton pump inhibitors; after ~7 months he returned with weakness, weight loss; hemoglobin alone had increased to 19 to 20gm%, with a hematocrit of 56.1%, other cell lines were not involved, suggesting secondary polycythemia; CT scan brain revealed Cerebellar Hemangioblastoma with obstructive hydrocephalus; emergency ventriculo peritoneal shunt was performed, followed by excision of the tumor; patient recovered and hemoglobin normalized.

Keywords: polycythemia, cerebellar hemangioblastoma.

GJMR-K Classification: NLMC Code: WL 320



Strictly as per the compliance and regulations of:



Polycythemia with Cerebellar Hemangioblastoma

Elizabeth JeyaVardhini Samuel ^α, Nagarajan Natarajan ^σ, Ramesh ^ρ & Latha M ^ω

Abstract- 56 year old male presented with one month history of vomiting with abdominal pain; hemogram suggested polycythemia with hemoglobin of 16.5-18.1gm%, gastroscopy showed pan gastritis, bone marrow smear, biopsy showed erythroid, megakaryocytic hyperplasia; ultra sonogram abdomen, pelvis did not show any significant abnormality; patient improved with parenteral hydration, proton pump inhibitors; after ~7 months he returned with weakness, weight loss; hemoglobin alone had increased to 19 to 20gm%, with a hematocrit of 56.1%, other cell lines were not involved, suggesting secondary polycythemia; CT scan brain revealed Cerebellar Hemangioblastoma with obstructive hydrocephalus; emergency ventriculo peritoneal shunt was performed, followed by excision of the tumor; patient recovered and hemoglobin normalized.

Keywords: polycythemia, cerebellar hemangioblastoma.

I. CASE DETAILS

5 6 year old male presented with vomiting, upper abdominal pain of ~30 days duration, associated with mild weight loss; he was a known hypertensive on irregular treatment but not a diabetic; he was a non smoker, non ethanol consumer, he used snuff, his wife had undergone permanent sterilization. General examination showed medium built middle aged male, with moderate dehydration, but no pallor, icterus, lymphadenopathy or pedal edema; his blood pressure measured 150/100 mm Hg; cardiovascular system, respiratory systems showed no abnormality; he was conscious oriented, with no neurological deficit; epigastrium showed tenderness, without guarding or rigidity; there was no hepatosplenomegaly.

His initial hemoglobin was 18gm%, with hydration the hemoglobin reduced to 16.5 to 17 gm%; liver function tests showed mild elevation of transaminase to 42 mg/dl, renal parameters were normal; peripheral smear showed reactive lymphocytosis, normochromic, normocytic red blood cells, with normal white blood cells, adequate platelets. Ultra sonogram of the abdomen showed simple renal cortical cysts, fatty change liver. Oesphago gastro

duodenoscopy showed pan gastritis; bone marrow biopsy done to evaluate for reactive lymphocytosis, with suspicious borderline polycythemia, showed hyper cellular marrow with erythroid, megakaryocytic hyperplasia.

He received treatment with proton pump inhibitors, parenteral hydration; patient's vomiting stopped, his abdominal pain improved and he requested discharge due to personal reasons.

6-7 months later patient returned with weakness, debility, for the past >30 days duration with obvious weight loss; examination suggested subtle Cerebellar signs, with suspicious, mild nystagmus on looking to the left side, suspicious impairment of tandem walking, there was no other neurological deficit; investigations revealed hemoglobin of 19.5 to 20 gm%, hematocrit of 56.1%, other cell lines were normal; the weight loss was very significant ~ 50%; since true secondary polycythemia was suggested by pure erythrocytosis without involving the other cell lines, ultra sonogram abdomen was normal, with significant weight loss, computerized axial tomography scan of brain was planned, to assess for Cerebellar Hemangioblastoma, probably having resulted in secondary polycythemia.

Computerized axial Tomography scan of the brain showed a 3.5cm × 3.8 cm mass lesion, with cystic areas, suggesting a possibility of Hemangioblastoma, with complete effacement of 4th ventricle, seen to cause obstructive hydrocephalus of the 3rd and both lateral ventricles; retinal examination showed papilloedema, but no hemangiomas; Magnetic resonance imaging of the brain showed solid, cystic mass in vermis, left Cerebellar hemisphere, to cause obstructive hydrocephalus Figure 1.

Author α : Consultant in Medicine. e-mail: elizabethsamuel@gmail.com

Author σ : Associate Professor

Author ρ : Post graduate Resident Senior Medical Officer Department of General Medicine.

Author ω : Pondicherry Institute of Medical Sciences, Pondicherry, India.



Figure 1: Tumor in cerebellum




Figure 1: Tumor in cerebellum




Figure 1 : Obstructive hydrocephalus caused by the tumor.

Emergency ventriculo peritoneal shunt was done followed by tumor excision by sub occipital craniectomy; post operative period was uneventful;

Histopathological examination of the excised tumor showed two components of the tumor-vascular and stromal; vascular component is in the form of network of capillaries and large thin walled vascular spaces; stromal component showed vacuolated large stromal cells with round to oval bland nuclei; occasional enlarged nuclei were seen; sparse mitoses were seen; foci of hyalinized vascular stroma were noted; tumor parenchyma interface was discrete in many foci, while creeping tumor was seen in occasional foci; capillary Hemangioblastoma-WHO grade 1 was reported Figure 2.




Figure 2 : HPE-Cerebellar Hemangioblastoma




Figure 2 : HPE- Cerebellar Hemangioblastoma

Post operatively hemoglobin returned to 15gm%, patient was discharged with anti epileptics to review in 3 months.

II. DISCUSSION

Polycythemia refers to high hemoglobin greater than upper limit of normal, in adult females >16.5gm% or hematocrit >0.48, in adult males hemoglobin of

18.8gm% or hematocrit >0.52 due to increase number of red blood cells [true polycythemia] or reduction in blood volume [apparent, relative polycythemia].

Causes of true polycythemia are ¹primary polycythemia—a myeloproliferative disorder e.g. Polycythemia Rubra Vera; secondary true polycythemia due to increased secretion of erythropoietin secondary to tissue hypoxia as in high altitude, lung disease,

cyanotic heart disease, high affinity hemoglobin, or secondary polycythemia due to inappropriately increased erythropoietin secretion by renal diseases such as hydronephrosis, cysts, carcinoma, other tumors like bronchogenic carcinoma, uterine fibroids, hepatoma, pheochromocytoma, Cerebellar Hemangioblastoma.

Polycythemia Rubra Vera is a ² clonal disorder involving a multipotent hematopoietic progenitor cell in which phenotypically normal red blood cells, granulocytes and platelets accumulate in the absence of a recognizable stimulus.

Since the patient had significant weight loss with true polycythemia in the second visit, not involving the other cells of granulocytes, and platelets, secondary polycythemia was considered; since ultra sonogram of the abdomen did not show any abnormality, search for tumors including Cerebellar Hemangioblastoma was undertaken.



Hemangioblastoma are rare benign tumors that occupy central ³ nervous system and other regions like kidney, liver, pancreas; the incidence among posterior fossa tumors averages 11%; highest incidence is in 3rd to 6th decades of life; family history is significant with autosomal dominant inheritance, related to deletion of tumor suppressor gene, on the short arm of chromosome 3.

Hemangioblastoma are benign vascular tumors that arise from embryonic remnants of mesoderm origin, trapped in nervous system, during first trimester ⁴ of life. Cerebellar Hemangioblastoma a rare cause of ⁵ erythrocytosis, consequent to ectopic production of erythropoietin by the tumor cells.

2 cases of Hemangioblastoma followed up long term showed malignant ⁶ behavior of transition to renal cell carcinoma from renal cysts

Hemangioblastoma is a tumor of the central nervous system, arising from vascular system, and can be present in retina ^{7 8 9 10} and spinal cord; can be associated with polycythemia, pancreatic cysts and Von Hippel Lindau syndrome.

As per the corresponding Authors earlier publications, couples observing contraception have 4-7 fold ¹¹ increase in tumors, cancers, among 35- >50 years of age, by fragmentation of the germ cells with associated decreased surveillance by estrogen, androgen ¹² resulting in uncontrolled proliferation of any cell, preceded by no differentiation, resulting in neoplasms. If only the wife's permanent sterilization can be reversed by tubal recanalization by awareness, financial aid, surgical expertise, the germ cells destruction will stop and the androgen, estrogen ¹³ will return to 79.9% of age normal, restoring genomic repertoire, cell cycle of differentiation, followed by controlled multiplication, so that recurrence and malignant transformation of the tumor will be curtailed.



III. CONCLUSION

56 year old male presented with spurious or relative polycythemia suspect, later on progressed to true secondary polycythemia with significant weight loss, prompting a search for neoplasms; clinically he had subtle Cerebellar signs and imaging confirmed the diagnosis of Cerebellar Hemangioblastoma, with associated obstructive hydrocephalus; emergency ventriculo peritoneal shunt followed by excision of the tumor was performed and the histopathological evaluation confirmed WHO grade 1 Cerebellar Hemangioblastoma; hemoglobin normalized with excision of the tumor.

IV. ACKNOWLEDGEMENTS

The needful expert services of Neurosurgery, General pathology, Radiology colleagues is deeply acknowledged with gratitude.

RÉFÉRENCES

1. JO Craig, DBL McClelland, H.G. Watson: Polycythemia; Davidson`s Principles and practice of Medicine, 21st edition, 2010, 998-999
2. Jerry L Spivak: Polycythemia Vera, Harrison`s Principles of Internal Medicine; 18th edition, 2012, volume 1, 898
3. HMFarrukh: Cerebellar Hemangioblastoma presenting as secondary erythrocytosis; West J Med. Feb. 1996; 164 (2) :169-171
4. L Ferranti, P Celli, B Fraioliand, A Santoro: Hemangioblastoma of the posterior cranial fossa; Acta Neurochirurgica 71, 283-294 1984
5. Chichiu SO, Lu en-Cheong Ho: Polycythemia secondary to Cerebellar Hemangioblastoma; American Journal of Hematology 71: 346-347
6. Seung Hwan Lee, Bong Jin Park, tae Sung Kim, Young Jin Lim: Journal of Korean Neurosurgical society , 2010, September 48 (3)
7. <http://neurosurgery.mgh.harvard.edu/newwhobt.htm>
8. Louis, David N (1991) WHO classification of tumors of the CNS. IARC. ISBN 92-832-2430-2
9. Lindsay, Kenneth W, Ian Bone, Robin Callander, Van Gijn (1991) Neurology and Neurosurgery illustrated United States Churchill Livingstone
10. Kaelin, WilliamG (2005): Von Hippel Lindau associated malignancies; Drug discovery today, Disease mechanism 2 (2): 225-231, doi: 10.1016/j.ddmec.2005.04.003
11. Elizabeth JS : Cancer and Germ cells; OJPM. 2014,4, 606-615; <http://www.scirp.org/journal/ojpm> <http://dx.doi.org/10.4236/ojpm.2014.47070>
12. Elizabeth JS: Increased prevalence of Renal Diseases-Metabolic syndrome and fragmented germ cells with reduced Endogenous Estrogen: Androgen *International Journal of Innovative Research and Studies*; Sep 2014 vol. 3 Issue 9 pp 71-83
13. Elizabeth JS: Abortion, Contraception Reversal- Medical Miracle- Autologous Germ cells replant- Decline in global degenerative, autoimmune, neoplastic, infectious diseases; decline in rising environmental estrogen [equating with aborted blood pollution] and decline in depletion of ozone. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* e-ISSN: 2279-0853, p-ISSN: 2279-0861. Volume 13, Issue 9 Ver. I (Sep. 2014) pp94-100



This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

My Pain is 10 out of 10. Patients Vs. Actors in Clinical Settings

By Mariyah Hussain, MD, Saeed Ahmed, MD, Hena Jawaid, MBBS, Mustafa Qureshi, MD,
Hooria Manzoor, MD, Tazeen Azfar, MD, Swati Sood, MD, Rizwan Ahmed, MBBS & Safiullah, MD

Nassau University Medical Center, New York

Abstract- Over the past two decades, opioid medication abuse among the U.S. population has expanded to a scourge extent. While the U.S. only accommodates 4% of the world's population, Americans consume 86% of the world's opioids, 99% of the worldwide hydrocodone supply, and 66% of the world's illegal drugs. Prescription opioids, which are currently the second most misused class of medication following cannabis. Results of the 2010 National Survey on Drug Use and Health (NSDUH) showed that a predicted 22.6 million people aged 12 or older were current or past month illicit drug users. Nearly 7 million individuals among these used marijuana and 5.1 million used painkillers. Only 17.3% of users of non-therapeutic opioids indicated that they received the drugs through a prescription from a physician. The widely growing use of therapeutic opioids shows hydrocodone topping all prescriptions with 136.7 million prescriptions in 2011, with all narcotic analgesics reaching more than 238 million prescriptions. Opioid analgesics are now accountable for a higher mortality rate than suicide and motor vehicle accidents. The majority deaths (60%) occur in patients who received prescriptions based on prescribing guidelines by medical boards, irrespective of the doses. In comparison, 40% of deaths occur in individuals abusing the drugs obtained through illegal means.

Keywords: drug seeking behaviors, opioid legitimate use, misuse, abuse and pain.

GJMR-K Classification: NLMC Code: WL 704



MY PAIN IS 10 OUT OF 10 PATIENTS VS ACTORS IN CLINICAL SETTINGS

Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

My Pain is 10 Out of 10. Patients Vs. Actors in a Clinical Setting

Mariyah Hussain, MD ^α, Saeed Ahmed, MD ^σ, Hena Jawaid, MBBS ^ρ, Mustafa Qureshi, MD ^ω, Hooria Manzoor, MD ^ξ, Tazeen Azfar, MD ^ς, Swati Sood, MD ^χ, Rizwan Ahmed, MBBS ^ν & Safiullah, MD ^θ

Abstract- Over the past two decades, opioid medication abuse among the U.S. population has expanded to a scourge extent. While the U.S. only accommodates 4% of the world's population, Americans consume 86% of the world's opioids, 99% of the worldwide hydrocodone supply, and 66% of the world's illegal drugs. Prescription opioids are currently the second most misused class of medication following cannabis. Results of the 2010 National Survey on Drug Use and Health (NSDUH) showed that a predicted 22.6 million people aged 12 or older were current or past month illicit drug users. Nearly 7 million individuals among these used marijuana and 5.1 million used painkillers. Only 17.3% of the users of non-therapeutic opioids indicated that they received the drugs through a prescription from a physician. The widely growing use of therapeutic opioids shows hydrocodone topping all prescriptions with 136.7 million prescriptions in 2011, with all narcotic analgesics reaching more than 238 million prescriptions. Opioid analgesics are now accountable for a higher mortality rate than suicide and motor vehicle accidents. The majority deaths (60%) occur in patients who received prescriptions based on prescribing guidelines by medical boards, irrespective of the doses. In comparison, 40% of deaths occur in individuals abusing the drugs obtained through illegal means. The objective of this article is to review the available evidence concerning misuse and differentiate a legitimate user from a drug-seeking addict. We would also discuss various aspects of drug abuse in clinical practice, mainly dealing with these patients in Emergency and Psychiatric settings.

Keywords: drug seeking behaviors, opioid legitimate use, misuse, abuse and pain.

I. INTRODUCTION

Sensory perception and its ability to bring external stimuli into awareness or consciousness have a significant role in the life of a human being. Pain is the major and most discussed, explored, and investigated part of a sensory system. Its severity correlates with the measure of touch, pressure, and vibration senses.

Author α: Foundation University Medical College, Pakistan.

Author σ: Nassau University Medical Center, NY.

e-mail: Ahmedsaeedmd@gmail.com

Author ρ: Aga Khan University, Pakistan.

Author ω: Brookdale University Hospitals and Medical Center (BUHMC).

Author ξ: Saint Vincent's Medical Center, Transplant Research Institute, California.

Author ς: Baqai Medical University, Pakistan.

Author χ: Dr. Rajendra Prasad Govt. Medical College, India.

Author ν: Liaquat Medical and Dental College, Karachi.

Author θ: Chandka Medical College, Pakistan.

The concept of pain has received significant focus because of its intensity and the motivation, which provokes one to seek a remedy through pharmacological, psychological, or social outlets. Regulation of pain-induced-behavior operates on the basis of two approaches: pain avoidance through anticipation or pleasure approach (1).

We try to understand most of the concepts in the domain of a Bio-Psychosocial model. Another (evolving) extension of this evaluation model is spiritual. This area, however, is undergoing current studies that might give us further insight in the future.

II. BIOLOGY OF PAIN

The biological underpinnings of pain deal with endorphins, enkephalin, and substance P(2-4). The saturation of GABA, cannabinoids, and nicotinic receptors underlie the maladaptive human behaviors to obtain pleasure. In other words, it is done to suppress nociception to the point of euphoria. The thalamus is a relay station for all sensory information that tries to correspond by activating response team for an action. The two ramifications of information processes go to the cerebral cortex (rational response) and the amygdala (emotional response), respectively. The final decision depends on connectivity and better transmission. Better the integration of synaptic connectivity and dendritic branching of an area, better or dominant will be the response.

III. PSYCHOSOCIAL ASPECTS OF PAIN

Most of the unrelenting and excruciating nature of pain does not respond to medical measures despite optimum dosages. This behavior reflects the associated parameters of pain. Mostly, this involves the emotional factor. Associated emotional memory enhances the consequential effect of pain despite adequate pain reduction. Emotional pain does not respond to pharmacological measures and the patient continues to stay in pain whilst showing persistent drug utilization behavior. It is an unfulfilled emotional need that shapes up drug seeking attitudes and establishes dependency.

In psychology, conversion is a known defense mechanism described by Freud in 1895. It is a conversion of psychological distress into a physical symptom. The measure of conflict reflects pain intensity,

severity, and nature. As soon as it gets resolved, the pain subsides on its own. It is an unconscious attention-seeking attitude which invites people for an attempt towards conflict resolution, the wish that cannot be verbalized otherwise.

The structure of personality, temperament, childhood experiences, history of environmental stimuli, parenting, interaction, life events, loss and social resources - all these factors determine our coping strategies and adaptation skills. The environmental stimuli can be good (authoritative parenting, supportive structure, academic achievements, ample resources, proper attachment, and encouragement) and can be bad (physical, emotional, sexual abuse, neglect, malnutrition, parental conflicts or separation weak attachment, authoritarian parenting, bullying, discouragement, and poor academic scores). Early experiences set up the basis of survival rules for children. These rules are an arbitrary reflection of their imagination regarding the world and defining their perception (5-7). The fearful personality architecture would have greater pain anticipation, which can stimulate early drug-seeking behavior (8).

The patients with a traumatic onset, injurious account are five times more at risk of obtaining opioid as a pain relieving strategy as compared to others who suffered the same extent of injuries giving non-traumatic perspective (2). The presence of self-efficacy belief is a belief in one's ability to succeed in a specific situation. If a person thinks, "symptom cannot be reduced, no matter whatever he does", then it is likely that symptoms may not be diminished by adequate measures. However, more optimistic expectation influences pain sensation (9).

The somatosensory amplification can be a communication mode for expressing distress (mainly psychological in nature). It depicts reduced threshold of tolerance, inculcates fear avoidance (10), and retrieval of (associated negative) episodic memory or pain related emotion (anxiety) (10, 11). This example can be understood by the presence of chronic pelvic pain sufferers with a history of sexual abuse. It has been demonstrated in many observational studies, maximum saturation does not heal one from pain neither does it reduce demands of pharmacological supplements. This behavior has invoked an idea to reflect the processing of pain in human brain with its associated emotional, memory, and temporal aspects.

IV. LEGITIMATE PATIENT VS A DRUG ABUSER

Medical user and non-medical drug users can be differentiated on the basis of symptom explanations and premorbid adjustment. With a history of poor work record, occupational and social changes, suspicion of non-medical drug use should be raised. The state of premorbid and comorbid substance use is crucial (12).

Over the past two decades, opioid drug therapy has soared up for non-cancer pain and hence opioid misuse has skyrocketed too (13). The United States Substance Abuse and Mental Health Services Administration defines the non-medical use of prescription opioids as a prevailing issue, which is increasing rather rapidly. According to their sources in 2002 and 2005; this point was further elaborated as taking someone else's medication or taking drugs only for the mere purpose of experiencing its effects (14, 15). Prescription drug abuse can additionally entail taking more than the optimum dosage, increasing frequency of drug administration, or even altering the drug formulation (eg. crushing) - all for the purpose of obtaining higher amount of the desired medication. In 2008, 73.8% of the prescription drug overdose deaths were a result of an overdose due to opioid pain relievers, thus highlighting its significant impact on mortality (16).

A study by Cicero TJ et al. showed the initial (genuine) prescription of opioid drug therapy in 79% of males and 85% of females on basis of pain led to 60-70% of them to misuse it in order to get "high" (17). According to Passik et al., the growing problem of the non-medical use of opioids requires mores refinement in order to be cured. Hence, consultations with specialists in mental health and other related fields should be encouraged (18).

V. SUSPICIOUS BEHAVIOR TO INVESTIGATE

Better screening and validated tools can help in making the process of selection more objective and authentic. The assessing instruments should be able to identify problems, predict risk for diversion, and choice for treatment (13). The consideration should be given to three componential categories 1) genetics 2) environmental influence 3) drug-related elements (19). That said, these factors do not affect independently but the effect is dimensional. Without the presence of psychosocial effects and family history, the potential of dependence reduces significantly despite drug-related factors like frequency, dose, etc.

Risk factor stratification should be done to help determine the intensity and frequency of monitoring the patients based on their risk of drug abuse. However, we should not use it to deny pain management in patients with high risk of abuse. All patients should be monitored to a minimal extent, which could then be more detailed based on the category in which they fit (low, medium or high-risk group) (13). A patient with a high risk of misuse tends to exaggerate in order to amplify the consequential effect of pain on a daily life routine, as compared to the user (20). Low pain tolerance is another hallmark for misuse (21). Younger individuals and a history of psychiatric illness are both predisposing factors for abusing prescribed drugs (22). According to a result of a multivariable analysis, there is

a significant association linked with past year non-medical use of prescription opioids with younger age (especially 18-21), Hispanic ethnicity, unemployment status, anxiety problems (panic, depression, agoraphobic, social phobic symptoms), history of alcohol dependence, cigarette smoking, history of other prescription drug misuse, other illicit substance use, and young age of initiating substance use (23). Based on previous studies, Conway and Dowling have found a correlation existing between anxiety and non-medical drug use, abuse, and dependence (24-29). This correlation raises the possibility that some opioid misuse could be better recognized and understood as self-medication. As in 1999, Fishman has devised excellent measures to deal with these stressful situations where differentiation is difficult to investigate (30):

- A. One pharmacy /one doctor visit
- B. If prescription is lost or prescription is utilized earlier, then it cannot be replaced
- C. Serials of urine drug screening tests.

VI. DOES THE PHYSICAL EVIDENCE FIT THE STORY?

Visser et al, describes in his management approach that a drug-seeking patient may be referred

Table 1: Prototypical Drug Seeking Behavior (Vukmir et al., 2004)

1. Multiple visits for same complaint
2. Unable to focus on anything other than the medicine
3. Lost prescription
4. Doctor unavailability
5. Allergic to new narcotic alternatives
6. Desire narcotics
 - a. Oral-codeine, oxycodone
 - b. IV-Demerol, morphine
7. Substitute benzodiazepines
8. Typical conditions that cannot be measured:
 - a. Headache
 - b. Urethral Colic
 - c. Toothache
 - d. Abdominal pain
9. Unbearable pains
10. Overly creative requests
11. Appearance change or alias

VII. DO'S AND DON'T WHEN A SUSPECTED DRUG ABUSER CONFRONTS YOU

a) DO'S

1. According to Health Insurance Portability and Accountability Act (HIPAA), for the purpose of treatment it is legal to share information with other healthcare professionals without the patients consent. Therefore, data should be obtained regarding health care (outpatient) visits.
2. Frequency of prescription utilization and checks on early use behavior.

as one "who is determined at all and any costs to support their narcotic dependency" (31). If a patient presents with an unreasonable and disproportionate focus on obtaining a desired specific pharmaceutical agent, without understanding its more appropriate other medical uses; he can be distinguished as a prototypical patient (32).

Signs of drug seeking should be differentiated from signs of addiction. An initial prescription to a medical user can merit subsequent dependence on prolonged use and can cause drug-seeking behavior. The strategy should be aimed to substitute stronger depending agent slowly with one of the weaker affinity drugs (33). Another angle of assessing addiction or getting clue of it is by 1) frequently "missing" prescription slips 2) "lost medications" 3) frequent narration of unpredictable circumstances leading towards more injuries (requiring more medicines) (34). Females are prone to more substance misuse due to affective instability and emotional problems while males' inclination is mostly due to legal purposes (35).

3. One pharmacy/one doctor rule should be implemented in case of suspicion.
4. Enumerate other medicine's name of the same class to the patient without using medical jargon.
5. Evaluate and confirm suspicious historical and exam findings (36).
6. Other illicit drug usages should be screened out.
7. Patient-provider relationship should be maintained in a practical manner.
8. Patient's autonomy should be maintained within legal boundaries.

b) *DON'T*

1. Patients should not be forced to reveal their identities and show diagnosis to pharmacists.
2. The possibility of under treatment should be examined (e.g. under-treating a patient with genuine health care needs).
3. Do not show discriminatory behavior towards potential drug user. *The practice should be discouraged not the person.*

VIII. HOW DO I IDENTIFY A DRUG IMPAIRED CO-WORKER?

There are some explicit signs which can be readily picked up in recent drug users. Frequent Monday absenteeism, progressive impaired concentration, wearing long sleeves when inappropriate, unreliability in keeping appointments and meeting deadlines, increasing personal and professional isolation, change in attitude and behavior with colleagues, insistence on personal administration of narcotic injections to patients, deteriorating functionality, and diminished physical health can serve as helpful indicators. Frequent drug testing cannot be the absolute answer (37). Some psychological responses can point out dependent behavior. The associated behaviors can be exaggerated pain avoidance, anger, depression, fear, mood swings, irritability, and the cognitive pattern of catastrophization (38, 39). The effect can be appreciated on the social role, coping strategies, physical and health-related behavior (40-42).

IX. OUR RESPONSIBILITIES AS A PHYSICIAN

With the great deal of impact prescription drug use is having on mortality, it becomes essential for physicians to accept responsibility and adopt practices which will help them identify potential drug addicts. Clinicians need to be well informed that acute pain responds well to opioids but not all pain is responsive to this class of drugs. Clinicians have become hesitant to prescribe opioids due to the fear of abuse. On the other hand, they might also underestimate risk of abuse among their patients. Therefore, it is vital to educate clinicians about the 4-As of opioid treatment - providing sufficient Analgesia to patients so that they may be able to engage in Activities of daily living while avoiding any Adverse effects and Aberrant medication-related behaviors (43). It is also highly important to initiate opioid therapy with addressing expectations of the patient regarding the treatment. Clinicians need to be open about benefits and risks of prescription opioids and should ensure that the patient has realistic expectations about the therapy.

External information outlets should be utilized properly such as drug screening tests, spouse interviews, interviews from a close relative or friend, premorbid occupational record or functioning should be

inspected. Treatment agreements should be made with patients. These agreements can require more detailed and periodic surveillance. Regular monitoring must require assessment and proper documentation of pain severity, functional competence, psychological health, and advancement toward obtaining treatment goals, treatment compliance, and presence of adverse effects, strange drug-related behaviors, and substance use. The other supporting instruction from external sources such as testing of biologic screening (e.g., urine), interviews with family or caregivers, review of medical records, payer opioid prescription data keeper or input from prescription monitoring programs can be helpful and should be utilized as required (13).

Models for risk stratification can be developed to keep a checklist on early refills, increasing quantity, frequent outpatient visits, and deteriorating social and personal aspects of life. Prescription monitoring programs should be held in different states and physician practices should be assessed on the basis of evidence-based guidelines and current recommendations should be used to supplement the information to caretakers and families (44). The educating physician can help them improve patient assessment and management too.

The safety principles for pharmaco-vigilance must be implemented to supervise the wider effects of drugs after its marketing (Post-market surveillance). Drug categories should be identified which contain addiction potential. As the reward circuitry is present in the brain (mesolimbic pathway), thus most of the drugs crossing the blood-brain barrier hold a dependency potential, mainly stimulants, sedative-hypnotics, relaxants, opioid analgesics, anesthetic agents, and anticonvulsants. Regulatory measures should include advertisement of disadvantages associated with abuse, misuse, and addiction. Behavioral strategies should be shaped up to deal addiction and dependence (45). The practitioner must be vigilant in prescribing opioids and should be able to enhance his own endeavors to halt substance abuse, but at the same time caution should be exercised not to jeopardize pain management in patients who are likely to benefit from it.

X. DISCUSSION

There is still little known about how many people take opioid for a legitimate medical condition before they later transition into the phase of non-medical opioid usage. There is also insufficient data describing the illicit market for prescription opioid (46, 47). Therefore, Zacny et al, suggest that there is a need for active monitoring in these areas as non-medical drug use is emerging as a growing public health problem at a rapid pace. Likewise Vukmir et al believe the key to treating this problem is to target the psychic, physical, and functional components individually. (32).

The retrospective analysis is not the only medium of understanding the persistent pain phenomenon, but the prospective examination can also help in revealing the maladaptive behavioral mechanisms. The potential of abuse is increasing due to poor adherence of healthcare professionals with prescription guidelines, impaired checkpoints in primary, secondary and tertiary healthcare systems and low-quality post-market surveillance of drugs. We need to be aware that the best and most effective strategies to prevent and treat the non-medical use of prescription opioids remain undiscovered. According to Vukmir et al, the case has been simplified since we have no disputes regarding the pain treatment in cancer patients and patients with acute pain syndromes. The debate emerges when we deal with patients who present with a "25 out of 10" pain and specifically seek narcotic regimes (32). Furthermore, there is a critical need to introduce awareness campaigns, select clinical criteria and develop weighted scores for identifying aberrant behavior, incorporate prescription drug monitoring programs, and implement health-promoting strategies that will contribute to reducing drug-seeking behaviors. Clinicians need to be reminded that when people do not receive adequate pain control, it makes them desperate. It is imperative that clinicians address these concerns in a nonjudgmental manner, such that does not negatively affect the therapeutic alliance. Therefore, physicians should be educated to consider multiple etiologies for atypical behavior while being compassionate in responding well to the patients' complaints of pain (18). It is quite evident that there is still not enough empirical data available to clinicians in this expertise. Hence, a need to conduct more trials with a formalized and accurate diagnostic criterion will enable clinicians to differentiate adequately between legitimate drug users and drug addicts.

REFERENCES RÉFÉRENCES REFERENCIAS

- Higgins ET. Beyond pleasure and pain. *American psychologist*. 1997;52(12):1280.
- Turk DC, Okifuji A. Psychological factors in chronic pain: evolution and revolution. *Journal of consulting and clinical psychology*. 2002;70(3):678.
- Gatchel RJ, Turk DC. *Psychosocial factors in pain: Critical perspectives*: Guilford Press; 1999.
- Turk DC, Flor H. *Chronic Pain: A Biobehavioral Perspective*. *Psychosocial factors in pain: Critical perspectives*. 1999;1:18.
- Keogh E, Cochrane M. Anxiety sensitivity, cognitive biases, and the experience of pain. *The Journal of Pain*. 2002;3(4):320-9.
- Asmundson GJ. Commentary: Anxiety, sensitivity and the pain experience. *European Journal of Pain*. 2001;5(1):23-5.
- Dahl J, Wilson KG, Nilsson A. Acceptance and commitment therapy and the treatment of persons at risk for long-term disability resulting from stress and pain symptoms: A preliminary randomized trial. *Behavior therapy*. 2004;35(4):785-801.
- Geiderman JM. Sympathetic dystrophy. *Annals of emergency medicine*. 2001;37(4):412-4.
- Doleys DM, Dolce JJ, Doleys AL, Crocker M, Wolfe SE. Evaluation, narcotics and behavioral treatment influences on pain ratings in chronic pain patients. *Archives of physical medicine and rehabilitation*. 1986;67(7):456-8.
- McCracken LM, Gross RT, Sorg PJ, Edmonds TA. Prediction of pain in patients with chronic low back pain: effects of inaccurate prediction and pain-related anxiety. *Behaviour research and therapy*. 1993;31(7):647-52.
- Crombez G, Vlaeyen JW, Heuts PH, Lysens R. Pain-related fear is more disabling than pain itself: evidence on the role of pain-related fear in chronic back pain disability. *Pain*. 1999;80(1):329-39.
- Liebschutz JM, Saitz R, Weiss RD, Averbuch T, Schwartz S, Meltzer EC, et al. Clinical factors associated with prescription drug use disorder in urban primary care patients with chronic pain. *The Journal of Pain*. 2010;11(11):1047-55.
- Sehgal N, Manchikanti L, Smith HS. Prescription opioid abuse in chronic pain: a review of opioid abuse predictors and strategies to curb opioid abuse. *Pain Physician*. 2012;15(3 Suppl):ES67-ES92.
- Use FAT. *Substance Abuse & Mental Health Services Administration*. 2010.
- Cohen DA, Reporting A. *Substance Abuse and Mental Health Services Administration* 6. 2002.
- Paulozzi LJ, Mack KA, Hockenberry JM. Variation among states in prescribing of opioid pain relievers and benzodiazepines—United States, 2012. *Journal of safety research*. 2014;51:125-9.
- Cicero TJ, Lynskey M, Todorov A, Inciardi JA, Surratt HL. Co-morbid pain and psychopathology in males and females admitted to treatment for opioid analgesic abuse. *Pain*. 2008;139(1):127-35.
- Passik SD, Kirsh KL, Webster L. Pseudoaddiction revisited: a commentary on clinical and historical considerations. *Pain management*. 2011; 1 (3): 239-48.
- Ballantyne JC, LaForge KS. Opioid dependence and addiction during opioid treatment of chronic pain. *Pain*. 2007;129(3):235-55.
- Sullivan MD, Edlund MJ, Fan M-Y, DeVries A, Braden JB, Martin BC. Risks for possible and probable opioid misuse among recipients of chronic opioid therapy in commercial and medicaid insurance plans: The TROUP Study. *Pain*. 2010; 150 (2): 332-9.

21. Compton P, Charuvastra VC, Kintaudi K, Ling W. Pain responses in methadone-maintained opioid abusers. *Journal of pain and symptom management*. 2000; 20(4):237-45.
22. Boscarino JA, Rukstalis M, Hoffman SN, Han JJ, Erlich PM, Gerhard GS, et al. Risk factors for drug dependence among out-patients on opioid therapy in a large US health-care system. *Addiction*. 2010;105(10):1776-82.
23. Becker WC, Sullivan LE, Tetrault JM, Desai RA, Fiellin DA. Non-medical use, abuse and dependence on prescription opioids among US adults: psychiatric, medical and substance use correlates. *Drug and alcohol dependence*. 2008; 94 (1):38-47.
24. Goodwin RD, Hasin DS. Sedative use and misuse in the United States. *Addiction*. 2002;97(5):555-62.
25. Haddox J, Smith M. Characteristics of non-medical users of hydromorphone: results from the National Survey on Drug Use and Health. College of Problems of Drug Dependence, Orlando, FL. 2005.
26. McCabe SE, Teter CJ, Boyd CJ. Illicit use of prescription pain medication among college students. *Drug and alcohol dependence*. 2005; 77 (1):37-47.
27. Simoni-Wastila L, Ritter G, Strickler G. Gender and other factors associated with the nonmedical use of abusable prescription drugs. *Substance Use & Misuse*. 2004;39(1):1-23.
28. Conway KP, Compton W, Stinson FS, Grant BF. Lifetime comorbidity of DSM-IV mood and anxiety disorders and specific drug use disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry*. 2006.
29. Dowling K, Storr CL, Chilcoat HD. Potential influences on initiation and persistence of extramedical prescription pain reliever use in the US population. *The Clinical journal of pain*. 2006; 22 (9):776-83.
30. Fishman SM, Bandman TB, Edwards A, Borsook D. The opioid contract in the management of chronic pain. *Journal of pain and symptom management*. 1999;18(1):27-37.
31. Vissers R. Drug seekers offer opportunity to practice humane medicine. *ACEP News, Post Assembly Edition*. 2002.
32. Vukmir RB. Drug seeking behavior. *The American journal of drug and alcohol abuse*. 2004;30(3):551-75.
33. Weissman DE, Haddox JD. Opioid pseudo-addiction—an iatrogenic syndrome. *Pain*. 1989; 36 (3):363-6.
34. Zacny J, Bigelow G, Compton P, Foley K, Iguchi M, Sannerud C. College on Problems of Drug Dependence taskforce on prescription opioid non-medical use and abuse: position statement. *Drug and alcohol dependence*. 2003;69(3):215-32.
35. Jamison RN, Ross EL, Michna E, Chen LQ, Holcomb C, Wasan AD. Substance misuse treatment for high-risk chronic pain patients on opioid therapy: a randomized trial. *Pain*. 2010;150(3):390-400.
36. Solis K. Ethical, legal, and professional challenges posed by “controlled medication seekers” to healthcare providers, part 2. *American Journal of Clinical Medicine*. 2010;7(2):86-92.
37. Hickox SA. Drug Testing of Medical Marijuana Users in the Workplace: An Inaccurate Test of Impairment. *Hofstra Lab & Emp LJ*. 2011;29:273.
38. Eccleston C. Role of psychology in pain management. *Br J Anaesth*. 2001;87(1):144-52.
39. Eccleston C, Crombez G, Aldrich S, Stannard C. Worry and chronic pain patients: a description and analysis of individual differences. *European journal of pain (London, England)*. 2001;5(3):309-18.
40. Morley S. Psychology of pain. *British journal of anaesthesia*. 2008;101(1):25-31.
41. Vlaeyen JW, Morley S. Cognitive-behavioral treatments for chronic pain: what works for whom? *The Clinical journal of pain*. 2005;21(1):1-8.
42. Pincus T, Morley S. Cognitive-processing bias in chronic pain: a review and integration. *Psychological bulletin*. 2001;127(5):599.
43. Passik SD, Kirsh KL. Assessing aberrant drug-taking behaviors in the patient with chronic pain. *Current pain and headache reports*. 2004;8(4):289-94.
44. Control CfD, Prevention. Adult use of prescription opioid pain medications-Utah, 2008. *MMWR Morbidity and mortality weekly report*. 2010; 59 (6):153.
45. Silverman K, Chutuape MAD, Bigelow GE, Stitzer ML. Voucher-based reinforcement of attendance by unemployed methadone patients in a job skills training program. *Drug and Alcohol Dependence*. 1996;41(3):197-207.
46. Compton WM, Volkow ND. Abuse of prescription drugs and the risk of addiction. *Drug Alcohol Depend*. 2006;83 Suppl 1:S4-7.
47. Compton WM, Volkow ND. Major increases in opioid analgesic abuse in the United States: concerns and strategies. *Drug Alcohol Depend*. 2006;81(2):103-7.



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Community Acquired Urinary Tract Infection Prevalence in a Tertiary Institution Based in Evbuobanosa, Edo State, Nigeria

By Otajewwo, F. D & Amedu, S. S

Western Delta University, Nigeria

Abstract- The prevalence rate of asymptomatic UTI among students of a community tertiary institution based in Evbuobanosa near Benin City, Nigeria was the focus of this work. A total of 390 freshly voided midstream urine samples collected into sterile plastic screw capped universal containers containing boric acid as preservative from students aged between 15 – 30 years were used for the study. Samples were screened for significant bacteriuria and puscells (neutrophil) counts. All positive samples with pyuria were aseptically cultured by standard methods on sterile Cystine Lactose Electrolyte Deficient (CLED) agar, MacConkey agar and Sabouraud Dextrose agar plates and incubated appropriately. Antibiotic susceptibility testing was done on isolated and identified uropathogens by Kirby – Bauer agar diffusion disc method. While 231(59.2%) samples yielded bacterial growth, 159(40.8%) yielded no growth.

GJMR-K Classification: NLMC Code: QY 185



Strictly as per the compliance and regulations of:



Community Acquired Urinary Tract Infection Prevalence in a Tertiary Institution Based in Evbuobanosa, Edo State, Nigeria

Otajewwo, F. D^α & Amedu, S. S^σ

Abstract- The prevalence rate of asymptomatic UTI among students of a community tertiary institution based in Evbuobanosa near Benin City, Nigeria was the focus of this work. A total of 390 freshly voided midstream urine samples collected into sterile plastic screw capped universal containers containing boric acid as preservative from students aged between 15 – 30 years were used for the study. Samples were screened for significant bacteriuria and puscells (neutrophil) counts. All positive samples with pyuria were aseptically cultured by standard methods on sterile Cystine Lactose Electrolyte Deficient (CLED) agar, MacConkey agar and Sabouraud Dextrose agar plates and incubated appropriately. Antibiotic susceptibility testing was done on isolated and identified uropathogens by Kirby – Bauer agar diffusion disc method. While 231(59.2%) samples yielded bacterial growth, 159(40.8%) yielded no growth. The uropathogens isolated included *Staphylococcus aureus* (33.1%), *Escherichia coli* (20.8%), coagulase negative Staphylococci (15.6%), *Klebsiella aerogenes* (7.9%), *Coliform* organisms (7.9%), *Candida albicans* (7.9%) and *Proteus* spp (6.8%) of which gram negative bacilli and gram positive bacteria accounted for 43.4% and 48.7% respectively. *Candida albicans*, a fungal uropathogen accounted for 7.9%. Occurrence of UTI in male and female students was 57.1% and 42.9% respectively of which UTI occurred highest in the 21 – 25 age group. More than 50% of isolated bacterial strains were sensitive to gentamicin only with more than 90% resistant to augmentin and nalidixic acid. *Staph aureus*, *Kleb aerogenes*, *Coliform* organisms and *Proteus* spp were multidrug resistant as they resisted 4, 7, 5 and 7 drugs respectively. Findings emphasize the need for regular and routine isolation of uropathogens in health institutions and subjection of same to invitro antibiotic sensitivity testing in order to obtain updated antibiograms of UTI pathogens. This will address the emergence of resistance genes and reduce healthcare cost and long stay in hospitals.

1. INTRODUCTION

Urinary tract infection(UTI) is the most common infection experienced by humans after respiratory and gastro-intestinal infections and also the most common cause of both community-acquired and hospital acquired (nosocomial) infections for patients admitted to the hospitals (Najaret *al.*,2009). UTI can be asymptomatic or symptomatic characterized by a wide range of symptoms from mild voiding irritation to

bacteraemia, sepsis or even death (Ranjbar *et al.*, 2009).

Infection of the urinary tract could manifest differently depending on the site of the infection and length of time involved (Takhar, 2011). Those that affect the lower urinary tract are called cystitis(i.e. involving the bladder alone with symptoms including painful urination, burning sensation, frequent urination or urge to urinate or both while those that affect upper urinary tract are referred to as pyelonephritis(i.e. involve the kidneys and other organs (Sarah,2010). The symptoms of the upper urinary tract infection include fever and flank pain during urination in addition to those of the lower urinary tract (Sarah 2010).

Urinary tract infection occurs more frequently in females than males due to the shortness and width of the female urethra to the vagina which makes it liable to trauma during sexual intercourse as well as bacteria being passed from the urethra into the bladder during pregnancy (Ebie *et al.*, 2001). The moist environment of the female's perineum favours microbial growth and predisposes the female bladder to bacterial contamination (Ebie *et al.*, 2001). In addition, urine of females was found to have more suitable pH and osmotic pressure for the growth of *Escherichia coli* than urine from males (Obiogbolu *et al.*, 2004).

Most UTIs are caused by gram negative bacteria like *Escherichia coli* and *Klebsiella*spp (Omonighoet *al.*, 2001; Ebie *et al.*,2001). Other bacterial pathogens frequently isolated include gram positive bacteria such as *Staph. aureus*, *Staph. epidermidis* and *Enterococcus*spp formerly called *Strept. faecalis* (Kashefet *al.*, 2010; Theodros, 2010;Mulugeta and Bayeh,2014) as well as *Proteus mirabilis*, *Pseudomonas aeruginosa*, coagulase negative staphylococci, *Acinetobacter*spp and *Serratia*spp (Shankel,2007).

Drug resistance among bacteria causing UTI has increased since the introduction of UTI chemotherapy (Nerurkar *et al.*,2012; Sood and Gupta,2012; Bahadin *et al.*,2011; Haideret *al.*,2010). The aetiological agents and their susceptibility patterns vary in various regions and geographical locations (Mulugeta and Bayeh, 2014). Knowledge of the local bacterial aetiology and susceptibility pattern is required to trace any change that might have occurred with time so that

Authors α σ : Department Of Microbiology & Biotechnology Western Delta University, Oghara, Nigeria.
e-mails: fesdotaj@yahoo.com, dafotaj@hotmail.com

updated recommendation for optimal empirical therapy of UTI can be made (Leegard *et al.*, 2000).

A number of studies have been done on the prevalence and antimicrobial resistance patterns of UTIs (Onhet *et al.*, 2006; Mbata, 2007; Okonko *et al.*, 2009). Be that as it may, there is no documented UTI study that has been carried out in Ewobanosa community. Ewobanosa community is a bini speaking community situated about 20km from Benin City, Edo State of Nigeria. The community has a privately owned polytechnic situated tangentially on the express road that connects Benin City to Asaba (in Delta State of Nigeria). It was therefore to extend the frontiers of knowledge of UTI that this work aimed at studying community acquired urinary tract infection prevalence among students in a tertiary educational institution in Evbuobanosa, Nigeria was carried out with the following objectives:

- Determine the frequency distribution of microbial uropathogens in urine samples obtained from the students.
- Determine the sex distribution of uropathogens isolated.
- Determine the age distribution of the uropathogens isolated.
- Determine the antibiotic susceptibility profiles of isolated uropathogens.

II. MATERIALS AND METHODS

a) Sampling

A total number of the three hundred and ninety (390) midstream urine samples were collected from students (both residential and non-residential) of Lighthouse Polytechnic situated in Evbuobanosa, a community in Orhionwon local government area of Edo State of Nigeria. Samples (which were freshly voided), were collected into sterile screw capped plastic universal containers containing a few crystals of boric acid as preservative. Recruited students were instructed on how to collect the samples. All samples were appropriately labeled and processed immediately in the Microbiology laboratory of Western Delta University, Oghara, Delta State. Students recruited were grouped into 15-20, 21-25 and 26-30 age brackets. The study designed was a descriptive cross sectional study. Samples were collected in March, 2014.

b) Ethical Clearance

Ethical clearance was sought and granted by the ethical committee of the Polytechnic. In addition, the recruited students gave their verbal informed consent after thorough explanation of the rationale for the study.

c) Processing of Samples

Samples were tested for significant bacteriuria by use of a modified semi quantitative technique

described by Mbata (2007). A standard bacteriological loopful of each urine sample (0.01ml) was spread over the surface of sterile Cystine Lactose Electrolyte Deficient (CLED) agar plates (LabM, UK). After inoculation, the plates were inverted and incubated at 37°C for 18-24 hrs. The number of bacterial colonies was counted and multiplied by 100 to give an estimate of the number of bacterial organisms per millilitre of urine. A significant bacterial count was taken as any count equal to or in excess of 10⁵ per millilitre.

d) Confirmation of Significant Bacterial Count by Microscopy

All samples that recorded significant bacterial counts were subjected to urine microscopy test to detect presence of five pus cells per high power focus (5PC/HPF) or 10 white blood cells (pus cells) /mm³ in urine sediments or deposits (Smith, 2004) using x40 objective microscopically. All samples that were positive for significant bacterial count and also recorded 5PC/HPF or 10PC/mm³ or more were cultured on suitable laboratory media.

e) Cultural Studies

Urine sample that recorded 5PC/HPF or 10PC/mm³ and were positive for significant bacteriuria test were cultured aseptically on sterile CLED agar (Lab M, UK), MacConkey agar (Lab M, UK) and Saboraud Dextrose agar (Lab M, UK) plates according to standard methods. All inoculated plates were incubated at 37°C for 24hrs. Pure isolates were then obtained and identified according to schemes provided by Cowan and Steel (1993) and Alexopoulos (1996). All identified isolates were subjected to antibiotic sensitivity testing. Fungal isolates were excluded from antibiotic sensitivity testing. All urine samples that yielded no bacterial growth were noted.

f) Antibiotic Sensitivity Testing

Antibiotic susceptibility testing was carried out on confirmed uropathogens according to the agar diffusion disc technique described by Bauer *et al.* (1966). A colony of each pure (axenic) isolate was streaked on sterile Mueller Hinton agar plates aseptically using sterile inoculating wire loop. The relevant multidiscs containing their respective minimum inhibitory concentrations (MICs) were used and included erythromycin (5ug), cefuroxime (30ug), gentamicin (10ug), nalidixic acid (30ug), cefixime (5ug), ceftazidime (30ug), ofloxacin (10ug), augmentin (30ug), nitrofurantoin (200ug), ciprofloxacin (10ug), streptomycin (10ug), chloramphenicol (30ug), tetracycline (10ug) and cotrimoxazole (25ug). Discs were aseptically placed (impregnated) firmly onto the surface of the dried plates using sterile forceps. The plates were left at room temperature for one hour to allow diffusion of the different antibiotics from the disc into the medium (Mbata, 2007).

The plates were then incubated at 37°C for 18hrs. Interpretation of results was done using the length of diameter of zone of inhibition (NCCLS, 2000). Zones of inhibition greater than 10mm were considered sensitive, 5-10mm moderate sensitive and no zone of inhibition, resistant.

g) Statistical Analysis

Simple percentages were used throughout.

III. RESULTS

Table 1 shows the isolated identified uropathogens from the samples that yielded growth after 24hrs incubation period. A total of 390 midstream urine samples were processed of which 159 (40.8%) yielded no growth at the end of 48hrs incubation at 37°C. A total of 231 (59.2%) samples yielded growth of seven microbial uropathogens were identified as follows: Bacterial colonies that were in clusters, gram positive, catalase positive, coagulase positive, DNAase negative, phosphatase positive, mannitol fermenting, raised, round and smooth colonies were identified as *Staphylococcus aureus*. All gram negative raised entire, circular, motile, lactose, glucose fermenting, indole positive, methylred positive, voges praskauer negative, citrate negative and urease negative bacilli strains were identified as *Escherichia coli*.

Bacterial colonies that were positive for all the characteristics of *Staph aureus* colonies but were negative for slide and tube coagulase tests were identified as coagulase negative staphylococci. Gram negative mucoid non-motile, lactose, adonitol, inositol, glucose fermenting, voges praskauer positive, urease positive, citrate positive and indole negative bacilli strains were identified as *Klebsiella aerogenes*. Coliform organisms were mixed cultures made of *Escherichia coli*, *Enterococcus faecalis* and *Clostridium perfringens*. Gram positive yeast cells, pseudohyphae positive, chlamydospores positive, germ tube positive, glucose, maltose galactose and sucrose assimilation positive strains were identified as *Candida albicans*. Gram negative, swarming, fish odour colonies on sodium chloride containing media, indole negative and urease positive strains were identified as *Proteus* spp. The frequency occurrence of the isolates and their strains therefore were as follows: *Staphylococcus aureus* (33.1%), *Escherichia coli* (20.8%), coagulase negative Staphylococci (15.6%), *klebsiella aerogenes* (7.9%), Coliform organism (7.9%), *Candida albicans* (7.9%) and *Proteus* spp (6.8%). On the whole, total gram negative and gram positive bacilli isolated represented 43.4% and 48.7% respectively of which 43.4% and 56.6% belonged to enterobacteraceae and non enterobacteraceae respectively.

Table 1 : Frequency of distribution of microbial uropathogens isolated from Samples

Isolated Uropathogens	No of Strains	Frequency of Strains
<i>Staphylococcus aureus</i>	75	33.1%
<i>Escherichia coli</i>	48	20.8%
Coagulase negative Staph	36	15.6%
<i>Klebsiella aerogenes</i>	18	7.9%
Coliform organisms	18	7.9%
<i>Candida albicans</i>	18	7.9%
<i>Proteus</i> spp	15	6.8%
Total n=7	231	100.0%

Gram negative bacteria: 43.4%
Gram positive bacteria: 48.7%
Enterobacteriaceae: 43.4%
Non-Enterobacteriaceae: 56.6%

Table 2 shows the sex distribution of the microbial strains isolated. Out of the total 231 strains isolated, 132 (57.1%) and 99 (42.9%) were obtained from male and female students respectively. In a

decreasing order, the highest occurring uropathogens in male students were *Staph aureus* (34.1%), *Escherichia coli* (22.7%), coagulase negative staphylococci (18.2%), Coliform organisms (9.1%), *Klebsiella aerogenes* (6.8%),

Candida albicans (4.6%) and *Proteus* spp (4.6%). *Candida albicans* (12.1%), *klebsiella aerogenes* (9.1%),
Similarly *Staph aureus* (30.3%), *Escherichia coli* (18.2%), coagulase negative staphylococci (12.1%), *Proteus* spp (9.1%) and Coliform organisms (6.1%)
uropathogens were isolated from female students.

Table 2 : Sex Distribution of uropathogens Isolated from Processed Samples

Isolated Uropathogens	No of Strains males (%)	No of Strains females (%)	Total %
<i>Staphylococcus aureus</i>	45 (34.1)	30 (30.3)	75(32.5)
<i>Escherichia coli</i>	30 (22.7)	18 (18.2)	48 (20.8)
<i>Coag. Neg. Staph</i>	24 (18.2)	12 (12.1)	36 (15.6)
<i>Klebsiella aerogenes</i>	09(6.8)	09 (9.1)	18 (7.8)
<i>Coliform organisms</i>	12 (9.1)	06 (6.1)	18 (7.8)
<i>Candida albicans</i>	06 (4.6)	12 (12.1)	18 (7.8)
<i>Proteus</i> spp	06 (4.6)	09 (9.1)	15 (6.5)
Total n=7	M 132 (57.1%)	F 99(42.9%)	231(100.0%)

The age distribution of the entire 231 uropathogen strains isolated is shown in Table 3. Recruited students were grouped into 15-20, 21-25 and 26-30 age brackets. Hence, 102 (44.2%), 87 (37.7%) and 42 (18.2%) of the sampled students belonged to 21-25, 15-20 and 26-30 age groups respectively. Seventy two (70.6%), 45(51.7%) and 15(35.7%) students of 21-25, 15-20 and 26-30 age group respectively were males while 30 (29.4%), 42(48.3%) and 27(64.3%) students of the same age groups were females. Consequently, the highest number of uropathogen strains of 72(70.6%) was isolated from male students belonging to 21-25 age group. Fifteen (35.7%) uropathogen strains were the least isolated from male students who occurred in the 26-30 age bracket. The highest and lowest uropathogen strains of 42 (48.3%) and 27 (64.3%) respectively were isolated from female students of 15-20 and 26-30 age group respectively.

Table 3 : Age Distribution of uropathogens Isolated

Uropathogens Isolated	15 -20		21 -25		26 – 30	
	M	F	M	F	M	F
<i>Staph aureus</i>	18 (40.0)	12 (28.6)	21 (29.2)	09 (30.0)	06 (40.0)	09 (33.3)
<i>Escherichia coli</i>	09 (20.0)	12 (28.6)	18 (25.0)	06 (20.0)	03 (20.0)	0 (0.0)
<i>Coag. neg Staph</i>	09 (20.0)	03 (7.2)	12 (16.7)	03 (10.0)	03 (20.0)	06 (22.2)
<i>Klebsiella aerogenes</i>	06 (13.3)	03 (7.2)	03 (4.2)	03 (10.0)	0 (0.0)	03 (11.1)
<i>Coliform organisms</i>	0 (0.0)	06 (14.3)	09 (12.5)	0 (0.0)	0 (0.0)	0 (0.0)
<i>Candida albicans</i>	0 (0.0)	03 (7.2)	03 (4.2)	06 (20.0)	03 (20.0)	06 (22.2)
<i>Proteus spp</i>	0 (0.0)	03 (7.2)	06 (8.3)	03 (10.0)	0 (0.0)	03 (11.1)

In Table 4, the antibiotic susceptibility patterns of all isolated uropathogens (with exception of *Candida albicans* and coagulase negative staphylococci) to some selected antibiotics are shown. The antibiotic sensitivity profiles of the 231 uropathogen strains to ciprofloxacin, cefuroxime, gentamicin, cefixime, ceftazidime, ofloxacin, streptomycin, erythromycin, tetracycline, cotrimoxazole, augmentin, nalidixic acid, nitrofurantoin, chloramphenicol are shown. Twenty four (32.0%), 11 (22.9%), 06 (33.4%) 09(50.0%) and 06(40.0%) strains of *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella aerogenes*, Coliform organisms, and *Proteus spp* respectively were sensitive to all the 14 (fourteen) antibiotics used.

With respect to *Staph aureus*, 12 antibiotics used were twelve because nalidixic acid and nitrofurantoin were not tested on it and whereas erythromycin and streptomycin were used for *Staph aureus*, they were not used for the other uropathogens. On the average, a higher percentage of pathogens resisted all fourteen antibiotics compared to those that were sensitive.

In terms of effectiveness of each of the antibiotics used, 108(62.1%), 60(34.5%), 48(27.6%), 45(25.9%), 39 (22.4%), 24(13.8%), 24 (13.8%), 21 (12.1%), 15(8.6%), 15(8.6%), 12(6.9%), 12(6.9%), 06(3.5%) and 03(1.7%) strains of all isolated pathogens were sensitive to gentamicin, ofloxacin, streptomycin, nitrofurantoin, ciprofloxacin, tetracycline, cotrimoxazole, chloramphenicol, ceftazidime, cefixime, erythromycin, cefuroxime, augmentin and nalidixic acid in that decreasing order. The two most sensitive drugs in this study were therefore gentamicin and ofloxacin whereas augmentin and nalidixic acid were the least sensitive. Hence, augmentin and nalidixic acid were the most

resisted antibiotics with ofloxacin and gentamicin as the least resisted.

Table 4 : Susceptibility Profiles of Isolated Uropathogens to Selected Antibiotics after 24hrs Incubation at 37°C

Isolated uropathogens	SELECTED ANTIBIOTICS USED (%)														MEAN (%)
	ERY	CRX	GEN	NAL	CXM	CAZ	OFL	AUG	NIT	CPR	STR	CHL	TET	COT	
<i>Staph aureus</i> n = 75	12 (16.0)	0 (0.0)	66 (88.0)	NA	0 (0.0)	0 (0.0)	12 (16.0)	0 (0.0)	NA	06 (8.0)	48 (64.0)	18 (24.0)	12 (16.0)	18 (24.0)	24 (32.0)
<i>Escherichia coli</i> n = 48	NA	06 (12.5)	24 (50.0)	03 (6.3)	09 (18.8)	03 (6.3)	30 (62.5)	06 (12.5)	27 (56.3)	12 (25.0)	NA	03 (6.3)	06 (12.5)	03 (3.3)	11 (22.9)
<i>Klebsiella aerogenes</i> n = 18	NA	0 (0.0)	06 (33.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	06 (33.4)	06 (33.4)	NA	0 (0.0)	06 (33.4)	03 (16.7)	06 (33.4)
<i>Coliform organisms</i> n = 18	NA	06 (33.4)	12 (66.7)	0 (0.0)	06 (33.4)	12 (66.7)	12 (66.7)	0 (0.0)	06 (33.3)	09 (50.0)	NA	0 (0.0)	0 (0.0)	0 (0.0)	09 (50.0)
<i>Proteus spp</i> n = 15	NA	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	06 (40.0)	0 (0.0)	06 (40)	06 (40)	NA	0 (0.0)	0 (0.0)	0 (0.0)	06 (40.0)
Total n= 174	12 (6.9)	12 (6.9)	108 (62.1)	03 (1.7)	15 (8.6)	15 (8.6)	60 (34.5)	06 (3.5)	45 (25.9)	39 (22.4)	48 (27.6)	21 (12.1)	24 (13.8)	24 (13.8)	

Table 5 shows the frequency occurrence of multidrug resistance among the isolated bacterial uropathogens. All the bacterial uropathogens apart from *Escherichia coli* were resistant to more than three antibiotics at a time. For clarity and for the purpose of this study, a pathogen is described as multidrug resistant to any of the selected antibiotics if more than 50% of its strains are resistant to it and hence, a

pathogen is multidrug resistant if it resists up to three drugs at a time. Therefore, only four out of the five bacterial uropathogens isolated were multidrug resistant. None was resistant to 3 drugs. *Staph aureus* strains were resistant to 4 drugs while *Coliform* organisms were resistant to 5 drugs. *Klebsiella aerogenes* and *Proteus* spp strains were resistant to 6 drugs each.

Table 5 : Occurrence of Multidrug Resistance among bacterial Uropathogens Isolated

Bacterial Uropathogens	Resistance to				
	3 drugs	4 drugs	5 drugs	6 drugs	>6 drugs
<i>Staph aureus</i> <i>n</i> = 75	-	+	-	-	-
<i>Escherichia coli</i> <i>n</i> = 48	-	-	-	-	-
<i>Coliform orgs</i> <i>n</i> = 18	-	-	+	-	-
<i>Proteusspp</i>	-	-	-	-	+

IV. DISCUSSION

In this study, out of 390 urine sample processed, 159(40.8%) samples yielded no growth and no significant microbial growth combined. Hence, 231(59.2%) samples yielded growth and significant growth all together. The absence of bacterial growth in 40.8% of processed samples was established as such when microscopically, a non-significant pus cell count (less than 5) was observed (Otajewwo, 2014). That 231 (59.2%) processed samples yielded significant microbial growth suggested a UTI prevalence rate of 59.2% in this study. This finding is consistent with 55%, 54.58%, 60% and 66% prevalence rates reported by earlier authors (Onuoha and Fatokun, 2014; Tolkoff and Rubin, 1986; Naylor, 1984; Warren, 1987; Kunin 1985) and inconsistent with higher prevalence rates of 77.3% 71.3%, 71.6%, 77.9%, and 86.0% as reported by some authors (Otajewwo, 2014; Jellheden and Norrby, 1996; Mbata 2007; Alabi *et al.*, 2014). It is on record that lower rates of 30.0%, 35.5%, 38.5%, 38.6%, 39.0%, 39.0% and 39.7% have been reported by some authors (Anochie *et al.*, 2001; Ebie *et al.*, 2001; Anigilaje and Bitto 2013; Akinyemi *et al.*, 2000; Ogomaka *et al.*, 2014; Otajewwo 2013 and Oladeinde *et al.*, 2011). Other previous authors recorded much lower UTI prevalence rates of 11.9%

26.7, 28.2% and 28.9% (Aiyegero *et al.*, 2007; Sobazak 2010; Harvna *et al.*, 2014; Okon *et al.*, 2014). The high or low incidence rates may be attributed to poor environmental conditions where the subjects reside in terms of lack of proper personal and environmental hygiene, low socio – economic status, sexual intercourse or sexual promiscuity, pregnancy *e.t.c* among Nigerian men and women (Akinyemi *et al.*, 2000; Kolawole *et al.*, 2009; Andriole, 1985).

Seven uropathogens were isolated in this study (Table 1) of which gram negative bacilli constituted 43.4% while gram positive bacteria accounted for 48.7%. Isolates that belonged to enterobacteriaceae and non-enterobacteriaceae families were 43.4% and 56.6% respectively (Table 1). Whereas *Staphylococcus aureus* and coagulase negative staphylococci were the only gram positive bacteria, *Candida albicans* represented the only member of the non-enterobacteriaceae. Finding is consistent with the report of a previous author who isolated 86.1% gram negative bacilli and 13.9% gram positive bacteria of which enterobacteriaceae accounted for 49.9% (Otajewwo, 2013). The report of this study does not agree with a report of Oluremi *et al.* (2011) which stated 85.0% gram negative bacilli and 15.0% gram negative of which enterobacteriaceae organisms constituted

66.7%. The report of Otajevwo (2014) stating 65.2% and 34.8% gram negative and gram positive bacteria respectively as well as 55.7% and 44.3% enterobactereaceae and non – enterobactereaceae respectively as isolates is also not consistent with the findings in this study.

In this study, the most and second most occurring uropathogens were *Staphylococcus aureus* (33.1%) and *Escherichia coli* (20.8%). Other uropathogens isolated were coagulase negative staphylococcus (15.6%), *Klebsiella aerogenes* (7.9%), *Coliform* organisms (7.9%), *Candida albicans* (7.9%) and *Proteus* spp. This result suggests that the two least isolated uropathogens in the study area are *Candia albicans* and *Proteus* spp. The occurrence of *Staphaureus* and *E.coli* as the most commonly occurring uropathogens is consistent with the report of a previous study (Otajevwo, 2014) as well as with reports (though in reverse order) of some earlier authors which stated *E.coli* and *Staphaureus* as the most and second most implicated organisms in UTI (Onuoha and Fatokun, 2014; Okon *et al.*, 2014; Akobi *et al.*, 2014; Ogomaka *et al.*, 2014; Okonko *et al.*, 2010; Nwanze *et al.*, 2007).

The present report however, does not agree with results of other studies (Omigie *et al.*, 2009; Abubakar, 2009; Otajevwo, 2013; Oluremi *et al.*, 2011; Daza *et al.*, 2001; Dimitrov *et al.*, 2004; Inabo and Obanibi, 2006; Dash *et al.*, 2008; Nwadioha *et al.*, 2010). Report in this study does not also agree with some other previous reports which stated *Klebsiella* spp and *Proteus* spp (Alabi *et al.*, 2014), *E.coli* and *Streptococcus faecalis* (Anigilaje and Bitto, 2014), *E.coli* and *Proteus* spp (Haruna *et al.*, 2014), *E.coli* and *Pseudomonas aeruginosa* (Habibu, 2014) as the most and second most occurring uropathogens in UTI. In this study, *Proteus* spp, *Klebsiella* spp and *Staph aureus* were isolated and these organisms have been incriminated in hospital acquired infections often following catheterization or gynaecological surgery (Cheesbrough 2000; Tapsak *et al.*, 1995). *Proteus* infection is also associated with renal stones (Cheesbrough, 2000). In this study and previous similar studies, it is yet again confirmed the prominent involvement of *Staphaureus* and *E.coli* in UTI cases (Logoria and Gonzalez, 1987; Obiogbolu *et al.*, 2009; Otajevwo and Eriagbor, 2014; Okon *et al.*, 2014; Onuoha and Fatokun, 2014; Ogomaka *et al.*, 2014; Akobi *et al.*, 2014).

Overall, UTI prevalence rate was higher in males (57.1%) than in the female students (42.9%) of which *Staphaureus*, *E.coli* and *Coliform* organisms occurred more in the male students than in the female students (Table 2). The earlier report of Otajevwo (2014) which recorded higher UTI prevalence rate does not agree with the finding of this report. The reason for a higher UTI prevalence rate in males than females in this study though not clear, may be due to lack of

circumcision, receptive anal intercourse (as in homosexuals) and HIV infection (Orret and Davis, 2006). Also in this study, *Candida albicans* occurrence was higher in female students compared to the males. According to Ochei and Kolhatkar (2008), yeast cells appear in urine as a result of contamination from women with vaginal candidiasis (occasionally seen in the urine of men) or may be seen in the urine of diabetic patients due to presence of sugar in the urine. Yeasts may also cause recurrent infections in debilitated and immune-compromised patients (Cheesbrough, 2000; Ochei and Kolhatkar, 2008). The occurrence of *Coliform* organisms is crucial in pregnant women although its occurrence in females is lower compared to males. The isolation of *Candida albicans* and *Coliform* organisms in this study is consistent with the report of a previous work (Otajevwo and Eriagbor, 2014). It has been documented that *Coliforms* and *Enterococcus* spp can cause UTI when present in high numbers on the perineum (Behzardi and Behzardi, 2010; Moore *et al.*, 2002).

This study, strangely, did not confirm reports of other studies which stated that UTI occurs more in females than in males except at the extremes of life (Ebie *et al.*, 2001; Kolawole *et al.*, 2009). In terms of age bracket, UTI occurred highest (44.2%) in the 21–25 age group followed by 37.7% occurrence rate of students in the 15 – 20 age bracket. This finding does not agree with the reports of some previous authors which stated that UTI is more frequent in females than in males (Mbata, 2007; Ibeawuchi and Mbata, 2002; Asinobi, 2002; Olaita, 2006). Findings in this work however, agree with the reports of these same authors which stated that UTI is most prevalent during youth and adulthood as indicated by the 21 – 25 age group occurring as the most implicated in UTI in this study. This age group as well as the 15 – 20 group consist of teenagers, adolescents and young people who are characteristically vulnerable to increased sexual activities that predispose them to UTI (Oladeinde *et al.*, 2011; Oluremi *et al.*, 2011).

There is considerable evidence of practice variation in the use of diagnostic tests, interpretation of signs or symptoms and initiation of antibiotic treatment such as drug selection, dose, duration and route of administration (Jamieson, 2006). For patients with symptoms of UTI and bacteriuria, the main aim of treatment is to get rid of bacteria causing the symptoms. Besides, there is need to obtain sensitivity reports before the start of antibiotic treatment to checkmate emergence of resistance strains and thus help in proper patient management. The decision to use a particular antibiotic however depends on its toxicity, cost and attainable level (Ibeawuchi and Mbata, 2002).

Therefore, antibiotic sensitivity testing was done on all seven isolates (minus *Candida albicans* – a fungal uropathogen in this study) and the resulting sensitivity

profile (antibiogram) showed susceptibility reactions of 108 (62.1%), 60(34.5%), 48(27.6%), 45(25.9%), 39(22.4%), 24(13.8%), 24(13.8%), 21(12.1%), 15(8.6%), 12(6.9%), 06(3.5%) and 03(1.7%) for gentamicin, ofloxacin, streptomycin, nitrofurantoin, ciprofloxacin, tetracycline, cotrimoxazole, chloramphenicol, cefixime, ceftazidime, erythromycin, cefuroxime, augmentin and nalidixic acid respectively. This suggests that more than 50% of the bacterial uropathogens implicated in UTI in this study were sensitive to gentamicin only. This finding with respect to gentamicin is consistent with the reports of some previous studies (Okon *et al.*, 2014; Akobi *et al.*, 2014; Onuoha and Fatokun, 2014; Otajewwo and Eriagbor, 2014). Less than 50% of the isolated uropathogens were sensitive to ofloxacin and streptomycin

It is cheering and surprising that gentamicin recorded more than 50% sensitivity. Cheering because, it is cheap and readily available and hence, patients who go down with UTI can afford it if prescribed. The use and administration of gentamycin however, should be closely monitored in order to avoid any possible abuse owing to its accessibility, availability and cheapness. It is surprising because, gentamicin ought to be prone to abuse (due to its cheapness) with the attendant development of resistance genes by pathogens to it. It can be reasoned however that gentamycin though cheap, is not easily accessible for abuse because it is available in injection form only. The fluoroquinolone–ofloxacin recorded below moderate sensitivity (i.e. 27.6%). Although this finding agrees with that of a previous study which recorded 30.2% (Otajewwo and Eriagbor, 2014), it is quite low when compared with 100% ofloxacin sensitivity recorded by Anigilaje and Bitto (2014) and more than 50% ofloxacin sensitivity reported by Okon *et al.* (2014). Sensitivity of streptomycin which was less than 30% was also very low when compared to 100% streptomycin sensitivity reported recently by an author (Habibu, 2014). Nitrofurantoin (22.4%) and ciprofloxacin (13.8%) sensitivities were quite low. The low sensitivities recorded for the fluoroquinolones (ofloxacin and ciprofloxacin) in this study was worrisome because these drugs are expensive and therefore are not readily accessible for abuse. In a similar study, Nakhjavani *et al.* (2007) reported that the widespread use of fluoroquinolones in medical centres is a possible cause of high level resistance to fluoroquinolones in UTI patients. Nwadioha *et al.* (2010) surprisingly, recorded a high sensitivity (80.0%) and above of UTI bacterial agents with ciprofloxacin. This is however open to re–evaluation through further studies by other workers.

The low sensitivity recorded for nitrofurantoin in this study is at variance with 100%, 97.6% and more than 50% sensitivities recorded by previous authors (Oluremi *et al.*, 2011; Haruna *et al.*, 2014; Alabi *et al.*,

2014). The least sensitive antibiotics were augmentin and nalidixic acid. The low sensitivity of nalidixic acid is surprising because it is a drug that is not commonly and routinely used in medical practice. The less than 4% sensitivity of augmentin is very low and disturbing in view of its usefulness in the treatment of UTI's and other diseases. The almost total resistance of augmentin is alarming as it may have lost its value in the treatment of UTI (Oluremi *et al.*, 2011).

Four out of the seven isolated uropathogens were resistant to more than three drugs and hence they were all multi–drug resistant (MDR). In this study, *Staph aureus*, *kleb.aerogenes*, *Coliform* organisms and *Proteus* spp were resistant to four drugs, seven drugs, five drugs and seven drugs respectively (Table 5). Multi drug resistance of *Staph aureus* may be due to beta–lactamase (penicillinase) encoding genes it carries on its plasmids as well as other extracellular and intracellular factors the organism elaborates. Besides, multi–drug resistant *Staph aureus* strains have been widely reported in some studies (Abubakar, 2009; Aiyegoro *et al.*, 2007; Gales *et al.*, 2000). Multi–drug resistance of *Kleb. aerogenes* and *Proteus* spp are documented. The prevalence of multiple antibiotic resistant strains in this study is a possible indication that very large population of bacterial isolates has been exposed to several antibiotics (Oluremi *et al.*, 2011).

V. CONCLUSION

A prevalence UTI rate of 59.2% in this study (though not very high), indicates that UTI may be a health problem among students (who are mainly residential) of Lighthouse Polytechnic, Evbuobanosa, near Benin City. The findings suggest that most (if not all) the uropathogens isolated in this study are excretable through urine to the environment of the study area, Ewobanosa and its environs. Besides, the students should be advised to step up their personal hygiene in their hostels and immediate environment. Where signs and symptoms of UTI are noticed notwithstanding the above, healthcare providers of the Polytechnic may administer doses of gentamicin only or ofloxacin or perhaps streptomycin for therapy. A prompt therapeutic intervention in this regard, will prevent asymptomatic UTI cases becoming symptomatic with the accompanying renal damage.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Abubakar, E.M. (2009). Antimicrobial agents in urinary tract infections at the specialist hospital, Yola, Adamawa state, Nigeria. *Journ Clin. Med. Res.*1 (1):001-008.
2. Akobi, E.C., Inyinbor, H.E., Akobi, O.A., Emumwen E.G., Ogedengbe S.O., Uzoigwe, E.O., Abayomi, R.O and Okorie, I.E.(2014).Incidence of urinary tract

- infection among pregnant women attending antenatal clinic at federal Medical Centre, Bida, Niger State, north central Nigeria. *Amer. Journ of Infectious Disease and Microbiology*. 2:34-38.
3. Akinyemi, K.O., Alabi, S.A., Taiwo, M.A and Omonigbehin, E.O. (2000). Antimicrobial sensitivity Pattern and plasmid profiles of pathogenic bacterial isolated from urinary tract subjects In Lagos. *Quarterly Journ Hosp. Med*. 1:7-11.
4. Alabi, O.S., Onyenwe, N.E., Satoye, K.A and Adeleke, O.E. (2014). Prevalence of extended beta-lactamase producing isolates from asymptomatic bacteriuria among students in a tertiary institution in Ibadan, Nigeria. *Nature of Science*. 12 (4):111-114.
5. Alexopolous, C.S; Mims, C.W and Blackwell, M.(1996). *Introductory Mycology*. Fourth edition. John Wiley and sons, Inc.; New York, p 868.
6. Aiyegoro, O.A., Igbinosa, O.O., Ogowonyi, I.N., Odadjare, E.E., Igbinosa, O.E and Okoh, A.L. (2007). Incidence of urinary tract infections among children and adolescents in Ile Ife, Nigeria. *Afr. Journal Microbiol. Res*. 1(2):013-019.
7. Andriole, V.T. (1985). The role of tamm-horsfall protein in the pathogenesis of reflux nephropathy and chronic pyelonephritis. *Yale Journ. Biol. Med*. 58:91-100.
8. Anigilaye, E.A and Bitto, T.T.(2013). Prevalence and predictors of urinary tract infections among Children with cerebral palsy in Makurdi, Nigeria. *Open Journal of Pediatrics*. 3:350-357.
9. Anochie, I.C., Nkanginieme, K.E.O and Eke, F.U.(2001). The influence of instruction about the method of urine collection and storage on the prevalence of urine tract infection. *Niger. Journ. Paediatrics*. 28(2):39-42.
10. Bauer, A.W., Kirby, W.M, Sherris, J.C and Juck, M.(1996). Antibiotic susceptibility testing by a Standard disc method. *Am. Sourn. Clen. Path*. 45:493-496.
11. Cheesbrough, M. (2000). *District Laboratory Practice in Tropical Countries*. Cambridge International Press. 434p.
12. Cowan, S.T and Steel, K.J.(1993). *Manual for the identification of medical bacteria*. 3rd edn. Cambridge University press, London, New York, Rockville and Sydney. 150p.
13. Dash, R., Samara, Z and Dinari, G.(2008). Uropathogens of various populations and their antibiotic susceptibilities. 10(10):724-726.
14. Dimitrov, T. S., Udo, E.E and Emara, A.F.(2004). Antibiotic susceptibility pattern of community acquired urinary tract infections in Kuwait hospital. *Med. Princ. Pract*. 13:334-339.
15. Ebie, M.Y., Ayanbadayo, J and Tanyiana, K.B. (2001). Urinary tract infection in a Nigeria military hospital. *Nigeria Journal of Microbiol*. 3: 25-29
16. Habibu, A. U.(2014). Prevalence of *Proteus mirabills* and *Pseudomonas aeruginosa* among female patients with suspected urinary tract infections attending Muhammad Abdullahi Wase Specialist Hospital, Kano, Nigeria. *Intern. Journal of Engineering and Science*. 3:28-31.
17. Haruna, M.S., Magu, J., Idume, J., Nosiri, C and Garba, M.A. (2014). Antibiotic susceptibility of some uropathogenic bacterial isolates from Ahmadu Bello University Teaching Hospital Zaria, Nigeria. *IOSR Journal of Pharmacy and Biological Sciences*. 9:20-23.
18. Ibeawuchi, R; and Mbata, T.I. (2002). Partional and Irrational use of *Antibiotics*. *Afri. Health*, 24(2):16-18.
19. Inabo, H.I and Obani, H.B.T.(2006). Antimicrobial susceptibility of some urinary tract clinical isolate to commonly used antibiotics *Afr. Journ. Biotechnol*. 5(5):487-489.
20. Jamieson DJ, Thesis RN, Rasmussen S.A. (2006). Emerging infections and Pregnancy. *Emerging infectious Dis*. 12(7):491-43.
21. Jellheden, B and Norrby, R.S.(1996). Symptomatic urinary tract infection in women in primary health care. Bacteriological, clinical and diagnostic aspects in relation host response to infection. *Scand. Prim. Health Care*. 14(2):122-128.
22. Kolawole, A. Kolawole, O. M, Kandaki-Olukemi, Y. T. Babatunde, S. K., Durowade, K. A. and Kolawole, C. F. (2009). Prevalence of urinary tract infections (UTI) among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State, Nigeria. *International Journal of Medicine and Medical Sciences* 1:163-167.
23. Kunin, C.(1985). Use of antimicrobial agents in urinary tract infections. *Adv. Nepir. Hosp*. 14:39-45.
24. Mbata, T. I.(2007). Prevalence and antibiogram of urinary tract infection among prison inmates in Nigeria. *Int. Jour. Microbiol*. 3(2):34-39.
25. Nakhjavani, F.A., Mirsalehian, A., Hamilan, M., Kazeem, B., Mirafshar, M and Jaba, I.F. (2007). Antimicrobial susceptibility testing for *Escherichia coli* strains to fluoroquinolones in urinary tract infections. *Iran Journal Publ. Health*. 36(1):89-92.
26. Nwadiana, S.I., Nwokedi, E.E., Jombo, G.T.A., Kasnibu, E and Alao, O.O.(2010). Antibiotic susceptibility pattern of uropathogenic bacterial isolates from community and hospital acquired urinary tract infections in a Nigeria tertiary hospital. *Internet Journal of Infect. Dis*. 8(1):001-008.
27. Obiogbolu, C.H., Okonkwo, L and Anyammere, C.O.(2009). Incidence of urinary tract infections among pregnant women in Awka metropolis, southeastern Nigeria. *Sci. Res. Essays*. 4:820-824.
28. Ochei, J and Kolhatkar, A. (2008). *A medical Laboratory Science Theory and Practice*.

New Delhi:Tata McGraw hill publishing company limited.255p.

29. Oladeinde, B.H., Omoregie, R.,Olley, M and Anunibe, J.A.(2011).Urinary tract infection in a rural community of Nigeria. *North Aer: Journ. Med. Sci*, 3(2): 75-77.
30. Olaita J.O. (2006). Asymptomatic bacteriuria in female student population of a Nigeria University.*The Intern.Microbial*.4:2-2.
31. Ogomaka, I.A., Dike, J.N., Ogbulie, T.E., Dike, O., Nwokeji, M.C and Egbuobi, R.C.(2014) Urinary tract infection among teenagers in Umudioka in Orlu local government of Imo State, Nigeria. *Scientific Research Journ.* 2:43-48.
32. Omigie, O, Okoror, L. Umolu, P and Ikuuh, G. (2009).Increasing resistance to quinolones. A four year prospective study of urinary infection pathogens.**2**: 171-175.
33. Otajevwo, F.D. (2013). Urinary tract infection among symptomatic outpatients visiting a Tertiary Hospital Based in Midwest Nigeria.*Global Journ. Health Sci*.
34. Otajevwo, F.D and Eriagbor, C. (2014). Asymptomatic urinary tract infection occurrence among students of a private university in western Delta, Nigeria. *World Journal of Medicine and Medical Science*. 2: 455-463.
35. Sobozak, B. A.(2010). Urinary tract infection in females. *Journ. ofMed.* 4:45-50





This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Maternal Health Care Utilization in the Eastern Cape, South Africa: A Qualitative Investigation

By Mr. Mluleki Tsawe & A. Sathiya Susuman

University of the Western Cape, South Africa

Abstract- Background: This study presents qualitative investigations showed in Mdantsane a township in the Eastern Cape Province of South Africa. The aim of the study is to examine the socio-demographic determinants of maternal health care utilization in this province.

Conclusion: The study has indicated that most health professionals and patients (women) are not aware of the available maternal health services, and this lack of awareness leads to a minimal utilization of such services.

Methods The qualitative investigation revealed some information with regard to the determinants of maternal health care utilization in the study area. **Results:** The investigation suggests that lack of awareness about the maternal health services offered within the public health system is an important determinant of the frequency in which maternal health services are used.

Keywords: utilization; antenatal care; postnatal care; public health; maternal health; private health; primary health care; millennium development goals.

GJMR-K Classification: NLMC Code: QW 553



Strictly as per the compliance and regulations of:



Maternal Health Care Utilization in the Eastern Cape, South Africa: A Qualitative Investigation

Mr. Mluleki Tsawe^α & A. Sathiya Susuman^σ

Abstract- Background: This study presents qualitative investigations showed in Mdantsane a township in the Eastern Cape Province of South Africa. The aim of the study is to examine the socio-demographic determinants of maternal health care utilization in this province.

MethodsThe qualitative investigation revealed some information with regard to the determinants of maternal health care utilization in the study area.

Results: The investigation suggests that lack of awareness about the maternal health services offered within the public health system is an important determinant of the frequency in which maternal health services are used.

Conclusion: The study has indicated that most health professionals and patients (women) are not aware of the available maternal health services, and this lack of awareness leads to a minimal utilization of such services.

Keywords: utilization; antenatal care; postnatal care; public health; maternal health; private health; primary health care; millennium development goals.

I. BACKGROUND

South Africa is working very hard to reduce maternal mortality. The South African Department of Health has adopted several initiatives to reduce the maternal mortality rate, and thus move closer to the 2015 Millennium Development Goal Five (of reducing maternal mortality by at least 75%). One such strategy is the adoption of the UNFPA's Campaign for Accelerated Reduction of Maternal Mortality in Africa (CARMMA) by the Department of Health. The department of health plans to use this campaign to fast-track the reduction of the maternal mortality rate, which is estimated at 300 maternal deaths per 100 000 live births, in order to meet the 2015 targets¹. Another strategy that the department of health has adopted is the Strategic Plan for Maternal, New-born, Child and Women's Health (MNCWH). The objective of the MNCWH is to reduce maternal mortality rates by at least 10% by 2016, and to ensure that all women have access to reproductive health services¹.

Author α: Research Scholar Dept. of Statistics & Population Studies
University of the Western Cape Cape Town, South Africa.
e-mail: 2861022@uwc.ac.za

Author σ: Professor (Associate) Dept. of Statistics & Population Studies
University of the Western Cape Cape Town, South Africa.
e-mail: sappunni@uwc.ac.za

In South Africa, Primary Health Care (PHC) is an initiative implemented by the government to provide basic health care within the public health sector. Primary health care has also been initiated in many other developing countries to deliver health care needs for their populations. This health care system is aimed at providing health care services to many people who cannot afford them from the private health sector^{2, 3, 4}. Within South Africa, this system faces many challenges, including management, health coverage, access to health care and health care utilization^{3, 4, 5}. Moreover, some health care facilities are not adequately equipped to handle the needs of the population (especially in rural areas), and some of these facilities do not provide the health services that are required of them^{6, 7}. Most of the challenges are attributed to the apartheid regime, which is argued to have laid the backdrop for the "failing" and "inadequate" primary health care system⁸. The primary health care sector is often surrounded by much controversy when it comes to the availability of resources, especially in rural areas. Health resources are unequally "distributed among rural and urban areas," including the unequal distribution of health facilities⁹. This uneven distribution presents problems when it comes to health care use in both rural and urban areas. The availability and accessibility of health care facilities are some of the determinants of health care use in South Africa, but are even more so for most of the rural areas therein^{10, 11, 12}. Maternal mortality is defined as the death of women during pregnancy, childbirth, or within 42 days after delivery^{13, 14}. Throughout the world, women's health is a priority for many countries, especially those with high mortality rates¹⁵. Another author argues that maternal mortality is higher for women living in rural areas and those living in poorer communities¹⁶. The current maternal mortality rate for South Africa is estimated at 300 maternal deaths per 100 000 live births; this is a considerable decrease (at 73%) when compared to the 2008 estimates of 410 maternal deaths per 100 000 live births^{14, 17}. But still MMR is quite high in South Africa. Nevertheless, South Africa has high maternal mortality rates, which are rising and this presents major challenges in terms of the country's prospects of achieving the Millennium development Goals¹⁵. There are less than two years left to achieve the targets set under Millennium

Development Goal 5, and it seems that South Africa is not anywhere near to reaching these Millennium Development targets set by the United Nations. Reducing the current maternal mortality rate by 75% would mean that South Africa's target for 2015 would be 225 maternal deaths per 100 000 live births, which is still quite high. According to the 2010 estimates of the World Health Organization, the maternal mortality rate in developing countries (of 240 maternal deaths per 100 000 live births) is 15 times higher than that of developed countries 14. It seems that many developing countries will not be able to reach, by 2015, maternal mortality statistics similar to those of developed countries. Therefore, given the current maternal mortality statistics, the prospects of South Africa achieving the Millennium Development Goals relating to maternal mortality seem very unlikely 4,18. Moreover, South Africa's estimated average annual percentage change in the maternal mortality rate between 1990 and 2010 is 0.9%, which shows that the country is not making progress with regards to improving maternal health by 2015 14. Therefore, improving the country's current approaches to health care and specifically, implementing new directions in maternal health care could improve the country's health and thus ensure that, over time, the Millennium Development Goals are met 19. In planning for the reduction of maternal mortality, attention should be given to the socioeconomic-demographic determinants of health, as this will improve maternal health 20. A number of strategies could be implemented in this area, in order to try and reduce the high maternal mortality rates within developing countries. These strategies would play a big role in campaigning for better implementation and use of maternal health services. These could include (a) the inclusion of the media, (b) educational programmes aimed at enhancing the literacy skills of women (especially in rural areas), (c) implementing better policies that are aimed at shaping or focusing on the livelihoods of women, and (d) implementing better maternal health care delivery 11, 21. It is important that developing countries implement strategies that will improve the socio-economic status of women, in order to accelerate health care use and thus enhance the utilization of maternal health care services 22.

a) Aim of study

The aim of this study is to examine the determinants of the use of maternal health care services in the Eastern Cape of South Africa. Moreover, the study aims to examine the kind of barriers that contribute to the lack of (or minimal) use of maternal health care services.

b) Research questions

The research questions for the qualitative study are as follows

1. What determines maternal health care use in the Eastern Cape?

2. What can be done to improve maternal health use in the province?
3. What challenges or problems do mothers experience while attending maternal health services?

II. METHODS

The qualitative investigation used, which were obtained from a study that made use of both quantitative as well as qualitative methods to collect data. Researcher very much interested to emphasis on qualitative issue here. This study makes use of mixed methods to study the determinants of health care utilization among women from the Eastern Cape province of South Africa. A qualitative investigation was obtained from health care professionals, using one-on-one interviews as the data collection method. A range of questions were asked to gain the health professionals' views on certain maternal health care problems or challenges, as well as some of the barriers that hinder women from utilizing such services.

a) Qualitative Investigation

In this section makes use of transcriptions from health care professionals in the Eastern Cape. The instrument used was an individual interview, wherein a range of questions were asked from the respondents (health care professionals) with regards to their views on certain health matters involving women and their reproductive health. For the analysis and interpretation, responses from two health professionals from each of the different professions, i.e., two doctors, two nurses, and two maternal health care specialists) were selected. This made it possible for rich qualitative data to be extracted from these health professionals, thus ensuring that there was variety in terms of views expressed.

III. RESULTS

a) Qualitative Results: Determinants of maternal health care use

Staff shortages became a major theme in the investigation when it came to trying to understand the challenges faced by health professionals while working within the antenatal care unit. Shortages of health care staff can have serious implications for the utilization of maternal health services, and it serves as a barrier that hinders the use of maternal health services.

b) One nurse stated that

"The waiting time is long"

A study finding argues that shortages in health professionals and longer waiting times reduce patients' satisfaction with the health-related services they receive 22. Moreover, this then leads to a decline in the utilization of health services. For instance, this can influence patients not seeing the need to seek maternal

health care services, whereby they will argue, “Why go to the hospital if I will not receive the adequate care that I need?”

Additionally, one health professional argued that mothers who attend antenatal care services face a number of challenges, including having to travel long distances, financial problems, and the attitudes of health professionals. Having no money and having to travel (often by walking) long distances to get to a hospital plays a role in the delayed use of antenatal service.

c) *One nurse stated that*

“[Most mothers and those who are expecting face] financial problems, [whereby] they won’t come on time here when referred and they will tell us that they didn’t have money to come here.”

Digressing from time-sensitive appointments and not seeking the available help can also be viewed in relation to the attitudes of health professionals.

d) *One doctor observed that:*

“Sometimes the attitude from the health care workers [hinders the satisfaction and use of maternal health services].”

One could argue that, due to the shortage of health professionals, there is a burden placed on the available staff to attend to a large number of women seeking maternal health services. This then has a repercussion on the type and quality of service offered, whereby some staff members will often take their frustrations out on the patients. Moreover, the attitudes of health professionals towards their patients are often driven by stigmas surrounding some health problems and certain social circumstances. For instance, with regards to teenage pregnancy and HIV, there seems to be a general stigma among health professionals. The doctors interviewed noted that teenage pregnancy and HIV were among the challenges they faced while working within the antenatal care unit. With regards to the treatment received by pregnant and HIV-positive teenagers,

e) *One doctor observed the following*

“The clinic [staff], they just chase them away and send them without helping them, which is really a problem. And the other clinics the still chas[ing] them for being HIV positive to the hospital which ... is not a good thing because we’ve to treat people the same.”

The way in which patients are treated in most public hospitals—or rather, the way in which health care professionals behave in such settings—can be seen as a barrier that hinders the use of maternal health services. For instance, teenage girls are often treated differently for seeking health care when pregnant, as compared to adults. This leads to many cases of young girls not seeking medical care in time, due to being

ashamed of what the nurses will think or say about them. In most instances, the hospital that one uses is the same hospital in which friends, family, or acquaintances work. This creates a barrier to use, in the sense that a pregnant teenager might decide not to go to a hospital where there is someone who knows her, and would rather stay at home until there are too many complications.

The health professionals noted a number of other reasons why most women are not utilizing maternal health services available at the hospital. They cited a lack of knowledge as one of the factors that contribute to women not using maternal health services. Most women are not even aware that they should be going for antenatal care when they are pregnant.

f) *One maternal health care specialist argued that*

“Lack of knowledge [is the leading contributing factor to the less-frequent use of maternal health services]. They are not aware that they should be coming to maternal health care [sessions] when they are pregnant; they don’t even know what maternal health care is.”

Besides not being aware of maternal health services offered at the hospital, most women “are not being referred” to antenatal care by their nurses and doctors.

g) *One maternal health care specialist stated that pregnant women are:*

“Only referred for maternal health care services when there is a problem” or complication, and these problems could be prevented if referrals could be done at initial contact.

Common risk-factors during pregnancy and the role that maternal health care specialists can play.

Apart from HIV, women who seek medical attention at the hospital present with a number of other pregnancy-related complications. The health professionals noted the following list of the most common pregnancy-related complications that women who seek medical attention present with: hypertension (high blood pressure); TB due to HIV; intra-uterine growth restriction (properly diagnosed and sometimes not diagnosed); and gestational proteinuria hypertension. All of these complications are serious risk factors that can alter a woman’s pregnancy experience. Doctors and nurses in the study argued that maternal health care specialists can assist them in ensuring that the above-mentioned pregnancy-related complications are prevented. Both the nurses and doctors also said that exercises are a good starting point in ensuring that pregnant women become active, in terms of strengthening their bodies during pregnancy, thus preparing their bodies for delivery.

IV. DISCUSSIONS

The qualitative results suggest that there are several determinants of maternal health care utilization in the province. These determinants have an influence on the frequency with which women use the available maternal health services. Women who have had bad experiences tend to not utilize maternal health services as frequently as those who have had good experiences. For instance, a woman who must wait for a long time before being attended (perhaps even going home unattended) might see no need to return to the clinic, unlike the one that received attention within a reasonable amount of time. Moreover, if a patient receives poor treatment, then that patient will view the health system in a different light. In some cases, she might even choose to seek medical attention from privately paid doctors, as opposed to waiting for hours on end at public hospitals or clinics.

Financial problems were also noted as one of the most significant factors that influence the use of maternal health services. One health professional noted: "Most of the women who go to seek maternal health services have to travel long distances, since most of them stay in rural areas, and sometimes they do not have enough money to get to the clinics or hospitals."

The quantitative analysis shows that more than 40% of the women reported that they do not have a source of employment. In such situations, financial problems become a barrier to the frequency with which these women seek medical attention.

Besides financial problems, lack of information also poses a serious hindrance to the utilization of maternal health services. The study has indicated that most health professionals and patients are not aware of available maternal health services, and this lack of awareness leads to a minimal utilization of such services. Health professionals do not refer pregnant women for antenatal services, and when those women have delivered, they are not referred for postnatal care, which then contributes to less frequent use of maternal health services. Sometimes, women do not go back to clinics for check-ups after they have delivered; they see no need to do so, unless certain complications arise. This then contributes to the low numbers of those seeking postnatal care.

Cultural factors are an important variable when it comes to understanding the factors that influence the use of maternal health care services. In South Africa, culture is an important concept that influences the way people live, as well as their belief systems. For instance, there are women that believe in utilizing traditional birth attendants, rather than seeking professional health care, due to their cultural beliefs; these women tend to opt for home-based deliveries, assisted by traditional birth attendants, rather than going to a hospital or clinic. The independent variable Do you have access to medical

aid? has an influence on whether or not people have access to maternal health services. One could deduce a number of reasons for this relationship. Generally, people who have access to medical aid tend to use private doctors rather than public hospitals (or clinics). This is due to a number of reasons, some of which will be further explained in the qualitative phase of the discussion. One such reason is the kind of treatment people generally receive in public hospitals; for instance, public hospitals are short-staffed and as such, adequate care of patients is often overlooked.

V. CONCLUSION

A lot can be done to improve the delivery and utilization of maternal health services. One such improvement would be to ensure that information relating to the importance (and utilization) of maternal health services is properly disseminated, so that more people can have knowledge of such services. There also needs to be something done with regards to how information is distributed to the relevant stakeholders. Policy makers should extend their efforts and work within the objectives of campaigns such as CARMMA (adopted by the department of health) and other organisations, in trying to reduce the state of maternal mortality in the country. The health sector also needs to inspect the way in which public health care professionals conduct themselves. If one could change the attitudes of nurses, then utilization of maternal health services (especially among pregnant youths, and those who get stigmatised for being HIV positive) might increase and thus lead to a decrease in maternal mortality. Therefore, from an observational and evaluative point of view, South Africa needs more initiatives that will strengthen its objectives of reducing the country's maternal mortality before the target year of 2015. Such initiatives must speak and add on to what programmes such as CARMMA has already achieved. Currently, the maternal mortality rate is at a level where it seems unlikely that the country will achieve its Millennium development Goal of reducing maternal mortality rate by 75% by 2015. This means that there is a lot that needs to be done in terms of coming up with initiatives that will reduce this rate, and thus save many lives. Moreover, a number of strategies could play a significant role in campaigning for better use of maternal health services, especially in rural areas. These campaigns could include (a) the inclusion of the media, in terms of broadcasting information relating to maternal health services and the importance of such services, (b) educational programmes aimed at enhancing the literacy skills of women (especially in rural areas), (c) implementing better policies that shape or focus on the livelihoods of women, and (d) implementing better maternal health care delivery. The qualitative results showcase the status of the South African public health

system. The responses from the health professionals point to the fact that the public health system is not properly functioning. Many hospitals, especially those in townships and rural areas, tend to be neglected and usually do not have the adequate staff to render health services to scores of people who seek them. It should also be noted that health care use is determined by the way which health professionals treat patients. There are still stigmas associated with certain health conditions and social situations. For example, there are still health professionals who refuse to assist, or may even maltreat, young girls who fall pregnant in their teenage years; these professionals even go so far as to mistreat pregnant women who are HIV positive. These girls and women do not adequately receive the care they would otherwise receive if their status were different.

Authors' contribution

Mr. M Tsawe is a post graduate student. Professor A Sathiya Susuman is a thesis supervisor. MT and AS worked together, contributed equally.

a) Ethical considerations

Ethical issues such as plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc., have been completely observed by the authors.

VI. ACKNOWLEDGEMENT

The authors would like to thank Statistics South Africa for providing data sets and offer special thanks to Prof. Richard Madsen, University of Missouri, United States, for his valuable suggestions and critical comments.

VII. Funding source

This work was supported by the National Research Foundation, South Africa.

VIII. Competing interest

None declared.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Cadegan, M., English, R, Pillay, Y., & Barron, P. A Brief Summary of the Strategic Plan for Maternal, Newborn, Child and Women's Health (MNCWH) and Nutrition in South Africa 2012 – 2016. Series name: Kwik Skwiz, issue two. Kwazulu-Natal: Health Systems Trust. 2012.
2. Peltzer, K., Skinner, D., Mfecane, S., Shisana, O., Nqeketo, A., Mosala, T. 'Factors Influencing the Utilisation of Prevention of Mother-to-Child Transmission (PMTCT) Services by Pregnant Women in the Eastern Cape, South Africa'. *Health SA Gesondheid*, 2005, 10(1):26-40.
3. Nteta, T. P., Mokgatle-Nthabu, M., Oguntibeju, O. 'Utilization of the Primary Health Care Services in the Tshwane Region of Gauteng Province, South Africa'. *PLoS ONE*, 5(11): 1-8. e13909. doi:10.1371/journal.pone.0013909.2010.
4. Naledi, T., Barroni, P., & Schneider, H. 'Primary Health Care in SA since 1994 and implications of the new vision for PHC re-engineering'. In: Padarath A, English R, editors. *South African Health Review* 2011. Durban: Health Systems Trust. URL: <http://www.hst.org.za/publications/south-african-health-review-2011.2011>.
5. Tanser, F., Gijsbertsen, B., Herbst, K. 'Modelling and understanding primary healthcare accessibility and utilization in rural South Africa: An exploration using ageographical information system'. *Social Science & Medicine*, 2006, 63: 691–705.
6. Beksinska, M., Kunene, B., & Mullick, S. 'Maternal Care: Antenatal, peri and postnatal'. In: Ijumba, P., Padarath, A., editors. *South African Health Review* 2006. Durban: Health Systems Trust. URL: <http://www.hst.org.za/generic/29.2006>.
7. Myer, L., & Harrison, A. 'Why Do Women Seek Antenatal Care Late? Perspectives from Rural South Africa'. *Journal of Midwifery & Women's Health*, 2003, 48 (4): 268–272.
8. Kautzky, K., & Tollman, S. M. 'A Perspective on Primary Health Care in South Africa'. In: Barron, P., & Roma-Reardon, J. editors. *South African Health Review* 2008. Durban: Health Systems Trust. URL: <http://www.hst.org.za/publications/841.2008>.
9. Ngwena, C. 'The Recognition of Access to Health Care as a Human Right in South Africa: Is It Enough?' *Health and Human Rights*, 2000, 5(1): 26-44.
10. Munsur, A. M., Atia, A., & Kawahara, K. 'Relationship Between Educational Attainment and Maternal Health Care Utilization in Bangladesh: Evidence from the 2005 Bangladesh Household Income and Expenditure Survey'. *Research Journal of Medical Sciences*, 2010, 4(1): 33-37.
11. Mattson, J. 'Transportation, Distance, and Health Care Utilization for Older Adults in Rural And Small Urban Areas'. *Small Urban & Rural Transit Center: North Dakota State University*. 2010.
12. Hogan, M. C., Foreman, K. J., Naghavi, M., Ahn, S. Y., Wang, M., Makela, S. M., Lopez, A. D., Lozano, R., & Murray, C. J. L. 'Maternal mortality for 181 countries, 1980–2008: a systematic analysis of progress towards Millennium Development Goal 5'. *Lancet*, 2010, 375: 1609-1623.
13. World Health Organization. Trends in maternal mortality: 1990 to 2010. Estimates developed by: WHO, UNICEF, UNFPA and The World Bank. Geneva: World Health Organization Press. 2012.
14. Blaauw, D., & Penn-Kekana, M. 'Maternal Health'. In: Fonn, S., Padarath A, editors. *South African*

Health Review 2010. Durban: Health Systems Trust.
URL:<http://www.hst.org.za/publications/876.2010>.

15. Say, L., & Raine, R. 'A systematic review of inequalities in the use of maternal healthcare in developing countries: examining the scale of the problem and the importance of context'. Bulletin of the World Health Organization, 2007, 85(10): 812–819.
16. World Health Organization. Trends in Maternal Mortality: 1990 to 2008. Estimates developed by: WHO, UNICEF, UNFPA and The World Bank. Geneva: World Health Organization Press. 2010.
17. King, M. S., Mhlanga, R. E., & de Pinho, H. 'The context of Maternal and Child Health'. In: Ijumba, P., Padarath, A., editors. South African Health Review 2006. Durban: Health Systems Trust. URL: <http://www.hst.org.za/generic/29.2006>.
18. Chopra, M., Lawn, J. E., Sanders, D., Barron, P., Karim, S. S. A., Bradshaw, D., Jewkes, R., Karim, Q. A., Flisher, A. J., Mayosi, B. M., Tollman, S. M., Churchyard, G. J., & Coovadia, H. 'Achieving the health Millennium Development Goals for South Africa: challenges and priorities'. Lancet, 374: 1023–1031. 2009.
19. Karlsen, S., Say, L., Souza, J-P., Hogue, C. J., Calles, D. L., Gülmezoglu, A. M., & Raine, R. 'The relationship between maternal education and mortality among women giving birth in health care institutions: Analysis of the cross sectional WHO Global Survey on Maternal and Perinatal Health'. BMC Public Health, 11(606): 1–10. Available online: <http://www.biomedcentral.com/1471-2458/11/606> Accessed: (09 May 2013). 2011.
20. Dagne, E. 'Role of socio-demographic factors on utilization of maternal health care services in Ethiopia'. Dissertation. Umeå University: Sweden. 2010.
21. Celik, Y., & Hotchkiss, D. R. 'The socio-economic determinants of maternal health care utilization in Turkey'. Social Science & Medicine, 2000, 50: 1797–1806.
22. Gerein, N., Green, A., and Pearson, S. The Implications of Shortages of Health Professionals for Maternal Health in Sub-Saharan Africa. Reproductive Health Matters, 14(27): 40–50. PII: S 0968-8080(06)27225-2. 2006.



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY

Volume 15 Issue 3 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

A Case of a Non-Verbal Child with Autism Gaining Faculty of Speech with Applied Energy Medicine

By Rajalakshmi Kandaswamy

Abstract- A 9 year old child diagnosed as a case of Nonverbal Autism and Attention Deficit Hyperactivity Disorder (ADHD) at 2 years of age started speaking normally and coherently for the first time in his life with 20 Remote Healing sessions using the method of Applied Energy Medicine with Intent Healing(TM).

Keywords: autism, non-verbal autism, energy medicine, intent healing (tm), remote healing.

GJMR-K Classification: NLMC Code: WS 350.8.A8



Strictly as per the compliance and regulations of:



A Case of a Non-Verbal Child with Autism Gaining Faculty of Speech with Applied Energy Medicine

Rajalakshmi Kandaswamy

Abstract- A 9 year old child diagnosed as a case of Nonverbal Autism and Attention Deficit Hyperactivity Disorder (ADHD) at 2 years of age started speaking normally and coherently for the first time in his life with 20 Remote Healing sessions using the method of Applied Energy Medicine with Intent Healing(TM).

Keywords: autism, non-verbal autism, energy medicine, intent healing (tm), remote healing.

I. INTRODUCTION

The incidence of Non-Verbal Autism is on the increase with apparently no effective known scientific cure for the same. I present a case here in which a child with a diagnosis of Non-verbal Autism and Attention Deficit Hyperactivity Disorder ,who had never spoken in his life was healed by me operating in the newly emerging scientific discipline of Applied Energy Medicine, using the Intent Healing (TM) Remote Healing method, following which he started speaking for the first time in his life. The child lived in California, U.S and I did the Remote Healing sessions from Chennai, INDIA.

II. CASE

A 9 year old boy who was diagnosed as a case of Non-verbal Autism and Attention Deficit Hyperactivity Disorder at the age of two years presented with the complaints of never having spoken a single word in his life in spite of his hearing and all other faculties being examined by doctors and reported as normal. Other complaints were hyperactivity, not making eye-contact, not studying on his own or going to school (every school the parents took him to refused admission as he was too hyperactive, disruptive in class and would wander away), not communicating or socializing, clenching his teeth repeatedly, making repetitive noises and pinching things with his fingers.

III. MATERIALS

Energy Medicine is the art and science of restoring a being/system to its natural state of well-being

and wholeness by augmenting the innate ability of the being/system to heal itself on all levels by bringing about shifts in the energy fields in the being/system to resonate with its natural frequency of alignment, balance and harmony. Intent Healing(TM) is healing using the power of Intention, accessing energies prior to consciousness that is free from all limiting conditioning and which brings about the realignment in the energy fields of beings/systems by rewiring the neural network in the brain and gut, reprogramming the DNA and erasing faulty cellular memories. To understand the newly emerging scientific field of Applied Energy Medicine and the Intent Healing(TM) method further, it is recommended to peruse the editorials given under the references in this report. They may be downloaded from www.intenthealing.com/blog

IV. METHODS

Energy based assessments and reports were given to each parent individually after the observation of their energies over two days and nights. Similar energy assessment was done for the child as well and the report of the child's energy pattern was shared with both parents. The specific and individual energy disturbances seen in each of them and the reasons for the same was explained in detail in the report. Remote Healing sessions with Intent Healing(TM) was done. The healing sessions were of two types - Timed remote healing sessions and healing sessions over Skype . The sessions over Skype was done with the parents and the sessions for the child was done as Timed remote healing sessions when the child was lying in bed at night. The total number of healing sessions was 20 of which 5 sessions were done with each parent (biological mother and father) and 10 sessions for the child. The sessions were spread over a period of one month. The energy disturbances in each parent and the child was corrected and their energies re-aligned in the healing sessions over Skype and in the Timed remote healing sessions. Regular feedback about the shifts seen in the energy fields of each parent and in the child was given to the parents after each healing session done with them and the child.

Author: Sowajya Centre For Holistic Healing, Chennai 600119, INDIA. e-mail : snowy123@gmail.com website : www.sowjanya.in

V. RESULTS

The child started singing first and then speaking by the end of 15 healing sessions (5 with each parent and 5 for the child). All the other symptoms improved right from the first healing session done with each parent and teeth clenching, making repetitive noises and pinching things with his fingers disappeared completely. There was dramatic improvement in the remaining symptoms ; he started making good eye-contact, started studying on his own using the laptop and started initiating friendship with other children his age in his apartment block at home. The hyperactivity reduced by 80 percent. The energy disturbances in each parent and the child was corrected and their energies re-aligned in the healing sessions over Skype and in the Timed remote healing sessions. All the symptoms of illnesses that were present in each parent, relating to the digestive system in the mother and the low back pain and fatigue that was present in the father also completely disappeared in their individual healing sessions. All of them reported a definite and tangible improvements in energy levels and mood right from the first session.

VI. DISCUSSION

The basis of using the approach of Applied Energy Medicine in autism using the method of Intent Healing(TM) lies in the scientific fact that everything in the Universe is ultimately made of Energy and everything is connected to everything else at the quantum level with this common Energy Field. Latest discoveries in science has confirmed the existence of Dark Energy and Dark Matter that makes up 96 percent of the Universe. It is possible to access and connect with individual energy fields in order to assess the energy patterns and correct the disturbances and restore the energy fields into Alignment using the power of Intention and the tool of Remote Intent Healing. The symptoms that are seen in autism are due to disturbances in the energy fields of the child. This disturbance is caused mainly by the disturbances in the energy field of the parents and care givers of the child and also the surrounding environment. Once the energy disturbances are corrected in the parents first and then in the child, and each of them experience the aligned energy state individually, the limiting symptoms disappear in the child with Autism and ADHD. Not only that, with the realignment of energies in each parent to their natural, harmonious and balanced energy states, the symptoms of disease in the parents too disappears. This points to the fact that the root cause of other medical symptoms too has an underlying energy basis that can be corrected with Applied Energy Medicine with Remote Intent Healing.

VII. CONCLUSIONS

Applied Energy Medicine using Intent Healing method can heal the limiting symptoms in autism including making children with Non-Verbal Autism gain their faculty of speech and start talking normally and coherently. It would be worthwhile to bring this quote by Tesla to our consciousness " The day science begins to study non-physical phenomena, it will make more progress in one decade than in all the previous centuries of its existence." – Nikola Tesla

REFERENCES RÉFÉRENCES REFERENCIAS

1. Rupert Sheldrake. Science Set Free. 1st edn. New York: Deepak Chopra Books. 2012.
2. Lynne McTaggart. The Intention Experiment: Using Your Thoughts to Change Your Life and the World. 1st edn. New York: Free Press. 2008.
3. Rajalakshmi K. Editorial. The Way Forward in Autism: The Paradigm Shift from the Problem to the Solution in Autism. Autism-Open Access Journal. 2014, 4:3.
4. Rajalakshmi K. Editorial. Epigenetics as a Solution in Autism: Control above Autism Genes. Autism-Open Access Journal. 2015, 5:1.
5. Rajalakshmi K , Premanand P. Paper presentation. Remote Healing As Evidence Based Medicine In Early Recovery Of Difficult Cardiothoracic Cases. Proceedings of International Cardiothoracic Conference; Feb 2008. AFMC Pune, INDIA.
6. Rajalakshmi.K. Editorial. . The Third Brain In Autism : Opening The Doors To The Solution In Autism. Neurology : Open Access Journal, Vol 1.1
7. W.A. Tiller, Science and Human Transformation: Subtle Energies,Intentionality and Consciousness, Walnut Creek, CA, USA: Pavior Publishing, 1997
8. What Are Subtle Energies? William A.Tiller,Journal of Scientific Exploration, Vol. 7, No. 3, pp. 293-304, 1993
9. White Paper on A Brief Introduction to Intention-Host Device Research by William A. Tiller, Ph.D. and Walter E. Dibble, Jr., Ph.D.
10. Jahn, R.G., and Dunne, B.J. (1987). Margins of reality: The role of consciousness in the physical world. Harcourt Brace Jovanovich, Publishers, New York.
11. Hunt, V. (1989). in Energy Fields in Medicine. Eds. M.A. Morton and C. Dlouhy, Fetzer Foundation,Kalamazoo, MI.
12. Lyn Buchanan, The Seventh Sense: The Secrets of Remote Viewing as Told by a "Psychic Spy" for the U.S. Military.Pocket Books, New York, February 2003.
13. Rajalakshmi.K. Case Report. A Case Of A 17 Year Old Boy With Autism Becoming Completely Independent And Thriving On His Own With Applied

Intentional Epigenetics. International Journal Of
Advances In Case Report, 2015; 2(9).

GLOBAL JOURNALS INC. (US) GUIDELINES HANDBOOK 2015

WWW.GLOBALJOURNALS.ORG

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards “FARSM” title to individuals. The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The “FARSM” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:



FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.



You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.


We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.





The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.

As FARSM, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.



The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.

The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website <https://associationofresearch.org> which will be helpful to upgrade the dignity.



The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.





The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.

The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.

The “MARSM” is a dignified ornament which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.



MARSM accrediting is an honor. It authenticates your research activities. After becoming MARSM, you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefits can be availed by you only for next three years from the date of certification.



MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.





We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.



Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.



AUXILIARY MEMBERSHIPS

Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as “Institutional Fellow of Open Association of Research Society” (IFOARS).

The “FARSC” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.



The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as “Institutional Board of Open Association of Research Society”-(IBOARS).

The Institute will be entitled to following benefits:



The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA). The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.



The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of “Open Association of Research Society, U.S.A (OARS)” so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



Journals Research
inducing researches

The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.



We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as “Institutional Fellow” and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.

The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:

Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.



Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.

Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.



We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth \$ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.



- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10% discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in-depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

//

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of “Difference of Opinion [if any]” among the Board members, our decision will be final and binding to everyone.

//



PROCESS OF SUBMISSION OF RESEARCH PAPER

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not convenient, and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



PREFERRED AUTHOR GUIDELINES

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.

Author Guidelines:

1. General,
2. Ethical Guidelines,
3. Submission of Manuscripts,
4. Manuscript's Category,
5. Structure and Format of Manuscript,
6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

- 1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.
- 2) Drafting the paper and revising it critically regarding important academic content.
- 3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5. STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also. Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

- (a) Title should be relevant and commensurate with the theme of the paper.
- (b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.
- (c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.
- (d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.
- (e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.
- (f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;
- (g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.
- (h) Brief Acknowledgements.
- (i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.



The Editorial Board reserves the right to make literary corrections and to make suggestions to improve brevity.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than $1.4 \times 10^{-3} \text{ m}^3$, or 4 mm somewhat than $4 \times 10^{-3} \text{ m}$. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art. A few tips for deciding as strategically as possible about keyword search:



- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.



Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org.



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.



12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.



Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

- Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

- Use standard writing style including articles ("a", "the," etc.)
- Keep on paying attention on the research topic of the paper
- Use paragraphs to split each significant point (excluding for the abstract)
- Align the primary line of each section
- Present your points in sound order
- Use present tense to report well accepted
- Use past tense to describe specific results
- Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives
- Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript-- must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.



- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently. You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.



THE ADMINISTRATION RULES

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.
- Do not give permission to anyone else to "PROOFREAD" your manuscript.
- **Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)**
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILATION)
BY GLOBAL JOURNALS INC. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	C-D	E-F
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring



INDEX

A

Aerogenes · 59, 62, 63, 64, 65, 67, 68, 69, 72

C

Chloramphenicol · 61, 65, 71

E

Encephalin · 53
Enhancement · 2, 36
Enterobacteriaceae · 68, 69
Erythrocytosis · 40, 48, 49
Estrogen · 48, 50

H

Hemangioblastoma · 1, 40, 45, 46, 48, 49
Hemangiomas · 40
Hydrocodone · 52

K

Klebsiella · 59, 62, 63, 64, 65, 67, 68, 69

N

Nitrofurantoin · 61, 65, 71

O

Occasional · 44

P

Polycythemia · 1, 40, 46, 47, 48, 49
Pseudomonas · 60, 69, 73
Psychiatric · 54, 56, 58

S

Staphylococcus · 59, 62, 64, 65, 68, 69

U

Uropathogen · 59, 64, 65, 70



save our planet



Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.MedicalResearchJournal.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals