

# How Much of a Role Birth Asphyxia and Chronic Antenatal Hypoxia Disorders have in the Genesis of Cerebral Palsy?

Dr. Kulkarni R.S.<sup>1</sup>, Dr. Aditya. P. Kulkarni<sup>2</sup> and Dr. Rachana. R. Kulkarni<sup>3</sup>

<sup>1</sup> Civil Hospital Oras, Sindhudurg

*Received: 15 December 2013 Accepted: 1 January 2014 Published: 15 January 2014*

---

## Abstract

Objective: Analysis were under taken to determine the role of birth asphyxia and chronic antenatal hypoxia disorders in the genesis of Cerebral palsy, in a prospective study of 31,804 antenatal mothers and 30,080 live births. Material Methods: For this large-scale prospective study, proper documentation of all events in the antenatal, natal and postnatal period, a detail, stringent protocol was prepared and distributed to 49 Govt. Z.P. health institutes. The protocol was filled in for each antenatal mother by the doctor of antenatal clinic and who is attending the delivery. The same was collected back to us by above-mentioned institutes on a fixed date of every month, at the time of monthly review meeting.

---

*Index terms*— cerebral palsy, birth asphyxia, chronic antenatal hypoxic disorders.

## 1 Introduction

William John Little in 1862, an Orthopaedic Surgeon presented a group of Children with tonal and developmental abnormalities, which he described as spastic rigidity. (1) Many of these children had a history of prolonged labour, preterm delivery. Because of frequency of these perinatal problems, Little postulated that the motor defects resulted directly from difficulties in the birth process. This opinion was widely held for over a century. Yet there were early critics, chief among them Sigmund Freud, who speculated that, perinatal difficulties were the result of pre existing abnormalities in the foetus rather than the cause of cerebral palsy. (2) This study was undertaken to identify and quantitate the major causes of cerebral palsy. The analysis were based on specific disorders that might damage a child's brain. (3) The most widely discussed of these disorders is birth asphyxia, with some people claiming that it is a frequent and others could be misleading because it is possible that such disorders are being missed or that insufficient cases have been analysed to find a correlation between them and cerebral palsy. (4) The first goal of the present study was to determine how much of a role birth asphyxia has in the genesis of cerebral palsy. A second goal was to quantitate the roles of chronic antenatal hypoxia disorders, congenital disorders, hypoglycemia, oxytocin, toxemia of pregnancy, mal presentations and other prenatal factors as causes of cerebral palsy. (5) II. natal clinic and who is attending the delivery. The same was collected back to us by above mentioned institutes on a fixed date of every month, at the time of monthly review meeting at District head quarter with district Civil Surgeon. Thus from 1 st Feb 1998 to 31 st Jan 2000, a prospective study of 31,804 antenatal mothers were followed up till delivery and 30,080 live births were observed in Sindhudurg district. The total number of live births for the above mentioned period was 32366 as per the vital statistics department of District Health Officer, Sindhudurg, thus un accounting the total live births of 2286, which include deliveries in small dispensaries, other nursing homes of outside the districts and home deliveries. All the children born were seen and examined at every six months intervals to identify cerebral palsy by a systematic and uniform record keeping system. The last neurological examination in the study was conducted in February 2002. Data received from above mentioned institute enlisted for investigation, which became available for analysis in March 2002. Data for antenatal mothers and intranatal mothers compiled by the respective doctors in the stringent protocols updated, we are fairly confident that the protocol data accurately reflect the Cerebral palsy pattern

44 in Sindhudurg district. 1065 children could not be analysed because the mothers delivered at different hospitals  
45 other than above mentioned institutes.

## 46 **2 Material & Methods**

## 47 **3 III.**

## 48 **4 Results**

49 Analysis were undertaken in a prospective study of 31,804 ante natal mothers who delivered from 1 st Feb 1998  
50 to 31 st Jan 2000.. 246 children were identified as cerebral palsy in 30,080 live births at the end of 3 rd serial  
51 examination. Thus 33% (82/246) victims of Cerebral palsy had birth asphyxia the presumed cause of their  
52 cerebral palsy. Of this 26% (64/246) were cases of quadriplegic cerebral palsy and 7% (18/246) non quadriplegic,  
53 which was attributable to the birth asphyxia. There was quiet a significant association of cerebral palsy with  
54 chronic antenatal hypoxic disorders.. Congenital disorders explained about one third of quadriplegic cerebral  
55 palsy. Birth asphyxia was not a significant antecedent of non quadriplegic cerebral palsy.

56 Finally the findings of the present study under score the importance of making accurate measurements and  
57 observations on neonates to avoid mistakes attributing non asphyxial cerebral palsy to birth asphyxia. The  
58 overall incidence of cerebral palsy for Sindhudurg Dist. amount to 8.1 per thousand live births over a period of  
59 1998 to 2000.

## 60 **5 IV.**

## 61 **6 Discussion**

62 Most studies that have attempted to determine if birth asphyxia is a cause of cerebral palsy, have used low  
63 Apgar scores and foetal distress to identify asphyxia. Low Apgar scores and foetal distress are often non hypoxic  
64 in origin, so their use as indicators of birth asphyxia could misattribute some non asphyxial cerebral palsy to  
65 asphyxia. (6) We explored this possibility by seeing how many victims of cerebral palsy who had low Apgar  
66 scores had a non asphyxial disorder as the basis for their cerebral palsy.

67 During the past two decades, dramatic changes in obstetrical and perinatal care have included the increasing  
68 availability of foetal heart monitoring and foetal ultrasonography, the establishment of neonatal intensive care  
69 units, and the implementation of policies to encourage the regionalization of care and the transport of mothers  
70 carrying high-risk fetuses before delivery. If the occurrence of cerebral palsy reflected sub optimal obstetrical  
71 care, (6) then its prevalence would be expected to decline in response to these remarkable improvements in care,  
72 but it has not done so. (8) In an attempt to evaluate the relative contribution of all pregnancy-related factors,  
73 some epidemiologists have created analytic models that evaluate later events (for example, those occurring during  
74 the delivery) (9) in the light of earlier events (characteristics of the mother before pregnancy, firsttrimester events,  
75 and so on). (10) , in the victims of cerebral palsy, characteristic consequences of birth asphyxia were more often  
76 the result of non-asphyxial disorders. (11) These included muconium in the amniotic fluid, low 10 minute Apgar  
77 scores.

78 Another perspective is gained by looking at the relative risks of various risk factors for cerebral palsy. Birth  
79 asphyxia had the highest relative risk for quadriplegic cerebral palsy. However, the low frequency of birth asphyxia  
80 in the population as a whole (82 of 30804) gave birth asphyxia a much smaller role as a cause of quadriplegic  
81 cerebral palsy.

82 Difference in distribution of factors related to cerebral palsy is highly significant. Since these factors are not  
83 mutually exclusive i.e. same case of cerebral palsy can have more than one factor hence chi square test won't  
84 make any sense really. A child whose mother has long intervals between menses appears to be at increased risk  
85 for cerebral palsy. (12) The risk is increased if there has been an unusually short interval (less than three months)  
86 or an unusually long interval (more than three years) since the previous pregnancy. (13) In addition, mothers of  
87 children with cerebral palsy are more likely than other mothers to have a history of spontaneous abortion and  
88 stillbirth. These findings indicate that maternal menstrual and obstetrical factors convey information about the  
89 risk of cerebral palsy.

90 Twins are more likely than singletons to have antenatal peri ventricular leukomalacia (14) and cerebral palsy.  
91 (15) Some of the increased risk of cerebral palsy among twins probably results from their gestational age and  
92 intrauterine growth retardation. In one study, an increase in the cesarean-section rate in the delivery of twins  
93 was not associated with a reduction in the prevalence of cerebral palsy. (16) The greater concordance for cerebral  
94 palsy among monozygotic than dizygotic twins also suggests a genetic basis, but it is compatible with placental  
95 problems that are unique to monozygotic twins as well.

96 Mothers known to have been hyperthyroid or who were prescribed thyroid hormones or estrogen in pregnancy  
97 have been found to be at increased risk of giving birth to a child in whom cerebral palsy later develops.

98 Non-vertex and face presentations of the foetus are associated with an increased risk of cerebral palsy. (17)  
99 One interpretation of this fact is that an abnormal presentation does not cause cerebral palsy, but rather may be  
100 a marker of preexisting difficulties. According to this hypothesis, fetuses with hypotonia and other abnormalities  
101 that will later be manifested as cerebral palsy are less able than others to move into a vertex position.

102 The rate of cerebral palsy is 25 to 31 times higher among infants who weigh less than 1500 g at birth than  
103 among full-sized newborns. (18) Babies whose birth weight is less than 2500 g account for about one third of  
104 all babies who later have signs of cerebral palsy. (19) As a generalization, the lower the birth weight and the  
105 gestational age, the higher the risk of cerebral palsy (20) and peri ventricular leukomalacia. Thus it should  
106 not be surprising that a number of low birth weight and early gestational age children are associated with peri  
107 ventricular leukomalacia, even among babies born prematurely. ??21) Nelson and Ellen berg wrote in 1986 "Of  
108 the . . . mother-infant pairs in the 5 percent with the highest risk (for cerebral palsy) only 208 percent produced  
109 a child with cerebral palsy, the false positive rate was thus 97 percent." Epidemiological studies published since  
110 then have not provided any reasons to change the impression that our ability to identify modifiable presumed  
111 causes of cerebral palsy is limited.

112 The burden imposed by cerebral palsy on society has not abated despite recent advances in medical care.  
113 Indeed, the increased survival of preterm newborns at risk for the disease has resulted in an increased number of  
children with cerebral palsy, mainly of the spastic diplegic variety. (22)



Figure 1: W

1

Sr. No.	Factor	No	Incidence	Relative Risk
1	Prematurity	148/246	60.2%	54.4
2	Low birth weight	136/246	55.3%	52.2
3	Low Apgar score & abnormal foetal heart rate	86/246	34.9%	10.7
4	IUGR on USG	82/246	33%	15.1
5	History of spontaneous abortion & stillbirth	68/246	27.06%	5.7
6	Toxaemia of pregnancy	44/246	17.9%	8.1
7	Forceps application	42/246	17%	42.2
8	Muconium stained liquor	36/246	14.7%	1.2
9	Malpresentation	34/246	13.9%	8.6
10	Oxytocin drip during labour	26/246	10.7%	2.5
11	Unusually long or short interval between pregnancy	26/246	10.6%	1.9
12	Caesarian section	16/246	6.5%	0.78
13	History of taking thyroid / oestrogen hormones	16/246	6.5%	43.5
14	Vaccum application	12/246	4.9%	29.6
15	Post maturity	12/246	4.9%	1.6
16	Bleeding during 1st , 2nd, 3rd, trimester of pregnancy	8/246	3.3%	1.3

The following positive antenatal, intranatal findings noticed are suggestive of quite a significant association of cerebral palsy with chronic antenatal hypoxia disorders. 60.2% (148/246) were born prematurely before 32 weeks of pregnancy. 55.3% (136/246) were low birth weight babies (below 2500 grams). Low Apgar scores & abnormal foetal heart rate during labour were present in 34.9% (86/246). Evidence of IUGR on USG was diagnosed in 33% (82/246). History of spontaneous

abortion and still births were detected in 27.6% (68/246). Toxaemia of pregnancy was noted in 17.9% (44/246) ante natal mothers. Forceps were applied during deliveries in 17% (42/246). Muconium stained liquor during labour was seen in 14.7% (36/246). Mal presentations were seen in 13.9% (34/246). Oxytocin drip was started during labour in 10.6% (26/246). Unusually long or short interval between the pregnancies was seen in ante natal mothers cerebral palsy children in 10.6% (26/246). Caesarean section was performed

Figure 2: Table 1 :

---

**2**

Sr. No.	O	E	(O-F) <sup>2</sup> /E
1	136	49.5	151.1
2	148	49.5	196.0
3	26	49.5	11.1
4	68	49.5	6.9
5	34	49.5	4.8
6	12	49.5	28.4
7	86	49.5	26.9
8	36	49.5	3.6
9	16	49.5	22.6
10	8	49.5	34.7
11	44	49.5	0.6
12	82	49.5	21.3
13	42	49.5	1.1
14	12	49.5	28.4
15	16	49.5	22.6
16	26	49.5	11.1
			571.2

$EX = 792 \times =$   $= 49.5 \times 2 = 571.2, df = 15 p < 0.001$

b) Difference is highly significant statistically

Figure 3: Table 2



115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162

Thus, efforts to prevent cerebral palsy will require a focus on factors and events during pregnancy including those that predispose the mother and foetus to preterm delivery and low birth weight.

[Freeman et al. ()] , J M Freeman , K B Nelson , Intrapartum . *Pediatrics* 1988. 82 p. .

[Dale and Stanley ()] ‘An epidemiologic study of cerebral palsy in Western Australia, 1956-1975 II: spastic cerebral palsy and perinatal factors’. A Dale , F J Stanley . *Dev Med Child Neurol* 1980. 22 p. .

[Nelson Kb Ellenberg ()] ‘Antecedents of cerebral palsy: multivariate analysis of risk’. J H Nelson Kb Ellenberg . *N Engl J Med* 1986. 315 p. .

[Bejar R Wozniak P Allard ()] ‘Antenatal origin of neurologic damage in newborn infants. I Preterm infants’. M Bejar R Wozniak P Allard . *Am J Obstet Gynecol* 1988. p. .

[Cooke ()] ‘Cerebral palsy in very low birthweight infants’. R W Cooke . *Arch Dis Child* 1990. 65 p. .

[Riikonen R Raumavirta S Sinivuori Seppala ()] ‘Changing pattern of cerebral palsy in the southwest region of Finland’. T Riikonen R Raumavirta S Sinivuori Seppala . *Acta Paediatr Scand* 1989. 78 p. 5817.

[Blair et al. ()] ‘Cooke RW Effects of birth weight gestational age, and maternal obstetric history on birth prevalence of cerebral palsy’. E Blair , P O Stanley Fj ; Pharoah , T Cooke , L Rosenbloom . *Arch Dis Child* 1987. 62 p. . (Intrauterine growth and spastic cerebral palsy I. Association with birth weight for 21)

[Paneth ()] ‘Etiologic factors in cerebral palsy’. N Paneth . *Pediatr Ann* 1986. 194-5, 197-201. 15 p. 191.

[Freud ()] S Freud . *Infantile cerebral paralysis*, (Russin, Trans). Coral Gables) 1897. 1968. FL University of Miami Press

[Kubli F (ed.) ()] *Grant A The relationship between obstetrically preventable intrapartum asphyxia, abnormal neonatal neurologic signs and subsequent motor impairment in babies born at or after term*, Kubli F (ed.) 1988. Berlin, West Germany: Springer Verlag. p. . (Perinatal Events and Brain Damage in Surviving Children)

[Kulkarni (2005)] ‘How much of a role birth asphyxia and chronic antenatal hypoxia disorders have in the genesis of Cerebral Palsy’. R S Kulkarni . *At Asia Pacific Childhood Disability Update* 2005 at Mumbai, on 4th December 2005. (Presented a paper on)

[Kulkarni (2004)] ‘How much of a role birth asphyxia and chronic antenatal hypoxia disorders have in the genesis of Cerebral Palsy? at 1st Annual conference of K’. R S Kulkarni .K.M.S. of I.A.P.S.M . *Govt. RCSM Medical College Kolhapur on 1st, August 2004*. (Presented a paper on)

[Little] ‘On the influence of abnormal parturition, difficult labors, premature birth and asphyxia neonatorum, on the mental and physical condition of the child especially in relation to deformities’. W J Little . *Transactions of the Obstetric Society of London* 3 p. .

[Torgs et al. ()] ‘Prenatal and perinatal factors in the etiology of cerebral palsy’. C P Torgs , B Van Den Berg , F W Oechsli , S Cummins . *J Pediatr* 1990. 116 p. .

[Kulkarni (2001)] *Presented a paper on Incidence and risk factors contributing to the development of cerebral palsy -a prospective study of 30804 antenatal mothers*, R S Kulkarni . 22 Dec. 2001. (At 46th Annual Conference of Indian Orthopaedic Association)

[Rydhstrom H] Prognosis for twins with birth weight less than 1500 gm the impact of cesarean scction in relation to fetal presentat  
‘Rydhstrom H Prognosis for twins with birth weight less than 1500 gm the impact of cesarean scction in relation to fetal presentation’. *Am J Obstet Gynecol* 1990. 163 p. .

[Bennett et al. ()] ‘Spastic diplegia in premature infants: etiologic and diagnostic considerations’. F C Bennett , S L Chandler , N M Robinson , C J Sells . *Am J Dis Child* 1981. 135 p. .

[Stanley ()] ‘Survival and cerebral palsy in low birthweight infants: implications for perinatal care’. F J Stanley . *Paediatr Perinat Epidemiol* 1992. 6 p. .

[Grether et al. ()] ‘The California Cerebral Palsy. Project’. J K Grether , S K Cummins , K B Nelson . *Paediatr Perinat Epidemiol* 1992. 6 p. .

[Stanley Fj Watson ()] ‘Trends in perinatal mortality and cerebral palsy in Western Australia’. L Stanley Fj Watson . *Br. Med J* 1967. 1985. 1992. 304 p. .

[Stanley ()] ‘Using cerebral palsy data in the evaluation of neonatal intensive care: a warning’. F J Stanley . *Dev Med Child Neurol* 1982. 24 p. .