

A Study to Find Anatomic Measurement Correlating to Occlusal Vertical Dimension: Comparison in two Ethnic Groups of Nepal

Dr. Bishal Babu Basnet¹, Dr. Raj Kumar Singh² and Dr. Prakash Kumar Parajuli³

¹ b. p. koirala institute of health sciences

Received: 14 December 2013 Accepted: 4 January 2014 Published: 15 January 2014

Abstract

Establishment of proper occlusal vertical dimension is the important yet daunting task for successful prosthodontic therapy for function, esthetics and comfort to the edentulous patients. In the current study the relationship of various distances between facial landmarks to the OVD was tested in two ethnic groups of Nepal, viz. Aryans and Mongoloids. The result of this study can be useful in determining proper OVD in the patients who have lost the OVD. The aim of the study was to find the correlation between facial landmarks and OVD. Materials and methods: The OVD was measured using a Boley gauge from the prominent part of chin and nose. Other facial landmarks were also measured using the calipers in 200 adult volunteers. Results: The Pearson's product moment correlation coefficient was determined.

Index terms— occlusal vertical dimension, rima oris to pupil distance, eye-ear distance, vertical height of ear.

1 Introduction

Glossary of Prosthodontic Terms 1 defines the occlusal vertical dimension as the distance between two selected points when the occluding members are in contact and rest vertical dimension as the distance between two selected points when mandible is in physiologic rest position. For a successful denture in terms of esthetics and function correct vertical dimension should be determined. 2 In clinical practice, there exists no universal rule for determining correct vertical dimension at occlusion because of wide range of individual variation. 3 Use of pre-extraction records, 4 maximum biting force, 5 electromyographic method, 6 and cephalometric radiograph 7 are some of the techniques utilized for determining occlusal vertical dimension. Millet et al. 8 recommended using multiple methods, like physiologic rest position, swallowing, phonetic, esthetics and facial measurements. Distance between soft tissue landmarks of face was significantly correlated to occlusal vertical dimension. 9 Author ? ? ? : Department of Prosthodontics, B. P. Koirala Institute of Health Sciences, Dharan, Nepal. e-mail: bishal_bpkihs@yahoo.com

The purpose of this study was to evaluate the correlation between facial measurements and occlusal vertical dimension in two ethnic groups of Nepal.

2 II.

3 Materials and Methods

Two hundred subjects between 19 and 46 years (mean age 25 years) were randomly selected from among the students, staffs, patients and relatives of patients visiting the College of Dental Surgery, B. P. Koirala Institute of Health Sciences. They were divided into two groups according to ethnicity, Aryans and Mongoloids. The inclusion criteria were full set of dentition, straight profile on visual examination, absence of malocclusion, no history of orthodontic and prosthodontic treatment. Each subject was instructed to relax the lips and close mouth in centric relation. The occlusal vertical dimension was measured between most prominent part of chin and nose. Other

facial measurements measured were the distance between pupil of eye to corner of mouth (rima oris), distance between outer canthus of one eye to inner canthus of other eye, distance between outer canthus to midpoint of external auditory meatus and vertical height of ear. Independent sample t test was performed to test the difference of means of studied parameters in males and females and in Aryans and Mongoloids. Correlation between these facial measurements and occlusal vertical dimension was studied with Pearson's correlation coefficient (r) at the significance level ≥ 0.05 and 95% confidence interval.

4 III.

5 Results

The mean occlusal vertical dimension of studied sample was 64.87 ± 5.21 mm. In Aryans, it was mm and in Mongoloids mm. The tables 1-4 show the results of this study. It was found that males have significantly longer dimension of OVD. The difference of means of rima oris to pupil distance, vertical height of ear, distance between outer canthus of one eye to inner canthus of other eye and eye-ear distance between males and females was not statistically significant. As we compared two ethnicities (Table 3), it was found that OVD and rima oris to pupil distance were significantly longer in Mongoloids whereas eye-ear distance was significantly longer in Aryans than Mongoloids.

The correlation between the facial measurements and occlusal vertical dimension was found statistically significant in both ethnicities (Table 4). However, the correlation between rima oris to pupil distance was stronger than the others. It was also noted that correlation between other facial measurements (eye-ear distance, distance between canthi and vertical height of ear) in Mongoloids was relatively weaker. Overall, rima oris to pupil distance was significantly correlated to occlusal vertical dimension (Table ??) with stronger correlation than other facial measurements. p-value significant at ≥ 0.05 level. Sig.-significant, NS-not significant IV.

6 Discussion

The loss of teeth results in poor appearance of lower third of face. The restoration of correct vertical dimension in those who have lost it should be in harmony with the upper part of face. Prosthodontists have long been seeking for universally accepted method of determining the OVD. There were methods like use of anterior teeth measurements 10 , closest speaking space 11 , swallowing method 12 , patient's neuromuscular perception 13 , cephalometric radiographs 14 , intra-oral and extra-oral anatomic landmarks 15 and measurement of fingers. 16 All of these methods possess some kinds of disadvantages. The present study gave some idea about facial measurement and their use in determining occlusal vertical dimension.

In present study the occlusal vertical dimension was found 65.90 ± 4.31 mm in male and 63.35 ± 6.03 mm in female. These were found to be longer than the findings of Ladda et al. 16 (61.4 ± 4.2 mm in male and 56.7 ± 3.0 mm in female) and Kulakarni and Kohli 17 (54 to 59 mm in female and 62 to 65 mm in male). The distance between ANS and Me from cephalometric radiograph in Serbian population, was 67.87 ± 5.67 mm in male and 61.51 ± 5.06 mm in female. 7 Similar measurements were observed in Iraqi adult population reported by Al-Hamadany 18 where occlusal vertical dimension in 75 students was 66.74 ± 6.468 mm. The OVD in males was 68.25 ± 6.134 mm and in females, it was 63.99 ± 6.254 mm. These values are consistent to that of current study.

The correlation was statistically significant and positive with rima oris to pupil of eye ($r=0.557$), followed by distance between outer canthus of one eye to inner canthus of other eye ($r=0.449$) and eye-ear distance ($r=0.395$) and vertical height of ear ($r=0.378$). This is in contrast with the study done by Delic et al. 19 The correlation of eye-ear distance ($r=0.8676$) was stronger than pupil to rima oris distance ($r=0.4357$) in their study. Rima-oris to pupil distance was longer than lower facial height by 3 cm in a study done in Phillipinos by Tina-Olaivar et al. 20 The limitations of the study were only two ethnic groups were taken into consideration and the findings cannot be extrapolated to other races. Also, the sample investigations to endorse or refute the findings of present study in this part of world as most of studies are found pertaining to developed countries.

V.

7 Summary

Anthropometric measurements of facial landmarks were recorded for 200 subjects (100 Aryans and 100 Mongoloids) and correlation of facial measurements to occlusal vertical dimension was tested. Within the limitations of this study, we can conclude that the facial measurements are significantly correlated to the occlusal vertical dimension. Findings of this study will be helpful guide in determination of occlusal vertical dimension. size was very small to conclude the findings. The occlusal vertical dimension is not constant and dynamic. Therefore the anthropometric measurement in dentulous subjects can be an adjunct but not accurate for utilizing in edentulous patients. There need to be further¹

¹© 2014 Global Journals Inc. (US)



Figure 1: Figure 1 :



23

Figure 2: Figure 2 :Figure 3 :



4

Figure 3: Figure 4 :



5

Figure 4: Figure 5 :

1

Character	Number	Minimum	Maximum	Mean	Std. deviation
Age	200	19	46	25	4.78
OVD	200	48	77	64.87	5.21
RO-Pu	200	52.5	72	64.77	4.65
OC-IC	200	55	77	66.99	3.74
E-E	200	53	82	69.32	4.17
EH	200	48.4	71	60.25	3.75

Figure 5: Table 1 :

2

Parameters	Gender	N	Mean±S.D.	p value	Remarks
OVD	Male Female	120 80	65.90±4.31 63.35±6.03	0.001	Sig.
RO-Pu	Male Female	120 80	64.78±4.43 64.77±4.98	0.437	NS
OC-IC	Male Female	120 80	67.27±3.40 66.57±4.19	0.053	NS
E-E	Male Female	120 80	70.33±3.91 67.83±4.11	0.967	NS
EH	Male Female	120 80	60.93±3.75 59.23±3.55	0.421	NS

[Note: p-value significant at ?0.05 level. Sig.-significant, NS-not significant]

Figure 6: Table 2 :

3

Parameters	Ethnicity	N	Mean±S.D.	p value	Remarks
OVD	Aryan Mongoloids	100 100	65.86±3.61 63.89±6.29	<0.001	Sig.
RO-Pu	Aryan Mongoloids	100 100	66.90±3.56 62.65±4.65	0.003	Sig.
OC-IC	Aryan Mongoloids	100 100	67.56±3.45 66.42±3.95	0.062	NS
E-E	Aryan Mongoloids	100 100	68.54±3.24 70.12±4.81	<0.001	Sig.
EH	Aryan Mongoloids	100 100	60.21±3.75 60.29±3.78	0.200	NS

p-value significant at ?0.05 level. Sig.-significant, NS-not significant

Figure 7: Table 3 :

4

Year						
Volume	XIV	Issue	VI			
Version	I					
() J						
Measurements	Pearson's r	Aryans p-value	Significance	Pearson's r	Mongoloids p-value	significance
OVD	1			1		
RO-Pu	0.577	<0.001	Sig.	0.466	<0.001	Sig.
OC-IC	0.508	<0.001	Sig.	0.307	0.002	Sig.
E-E	0.498	<0.001	Sig.	0.320	0.001	Sig.
EH	0.471	<0.001	Sig.	0.278	0.005	Sig.

[Note: p-value significant at ?0.05 level. Sig.-significant, NS-not significant]

Figure 8: Table 4 :

4

Measurements	Pearson's r	p-value	Significance
OVD	1		
RO-Pu	0.557	<0.001	Sig.
OC-IC	0.499	<0.001	Sig.
E-E	0.395	<0.001	Sig.
EH	0.378	<0.001	Sig.

Figure 9: Table 4 :

-
- 95 [Rev Stomatol Chir Maxillofac (2010)] , *Rev Stomatol Chir Maxillofac* 2010 Nov-Dec.. 111 (5-6) p. .
- 96 [Tina-Olaivar and Olaivar (1998)] ‘A comparative study of the upper and lower vertical facial measurements of
97 the Filipinos as it is used in the Willis method for determining the vertical dimension of occlusion’. E O
98 Tina-Olaivar , O K Olaivar . *J Philipp Dent Assoc* 1998 Jun-Aug. 50 (1) p. .
- 99 [Ladda et al. ()] ‘A new technique to determine vertical dimension of occlusion from anthropometric measure-
100 ments of finger’. R Ladda , A J Bhandari , V O Kasat , G S Angadi . *Indian J Dent Res* 2013. 24 p.
101 .
- 102 [Basler et al. ()] ‘Cephalometric analysis of the vertical dimension of occlusion’. F L Basler , J R Douglas , R S
103 Moulton . *J Prosthet Dent* 1961. 11 p. .
- 104 [Strajnic et al. ()] ‘Cephalometric indicators of the vertical dimension of occlusion’. L Strajnic , D Stanisic-
105 Sinobad , D Markovic , L Stojanovic . *Coll Antropol* 2008. 32 (2) p. .
- 106 [Miljcovic et al. (2001)] ‘Comparison of physiologic methods of determination of occlusal vertical dimension in
107 edentulous persons’. Z Miljcovic , M Zeljkovic , M Anojcic . *Vojnosanit Pregl* 2001 Jul-Aug. 58 (4) p. .
- 108 [Al-Hamdany and Kassab ()] ‘Correlation of vertical dimensions of soft tissue facial profiles’. A K Al-Hamdany
109 , N H Kassab . *Al-Rafidain Dent J* 2010. 10 (2) p. .
- 110 [Al-Hamdany and Kassab ()] ‘Correlation of vertical dimensions of soft tissue facial profiles’. A K Al-Hamdany
111 , N H Kassab . *Al-Rafidain Dent J* 2010. 10 (2) p. .
- 112 [Fayz and Eslami (1988)] ‘Determination of occlusal vertical dimension: a literature review’. F Fayz , A Eslami
113 . *J Prosthet Dent* 1988 Mar. 59 (3) p. .
- 114 [Kulkarni and Kohli (2011)] ‘Estimation and correlation of individual facial height and total body height’. N
115 Kulkarni , M Kohli . *J Int Oral Health* 2011 Apr. 3 (2) p. .
- 116 [Delic et al. ()] ‘Evaluation of craniometric methods for determination of vertical of occlusion’. Z Delic , M
117 Simunovic-Soskic , R Perinic-Grzic , S Vukovojac , Z Rajic , T Kuna . *Coll Antropol* 2000. 1 p. . (suppl. 24)
- 118 [Ekfeldt and Karlsson ()] ‘Influence of lower lip support on recording of vertical dimension in edentulous patients’.
119 A Ekfeldt , S Karlsson . *J Oral Rehabil* 1992. 19 p. .
- 120 [Boos ()] ‘Intermaxillary relation established by biting power’. R H Boos . *J Am Dent Assoc* 1940. 27 p. .
- 121 [Shanahan ()] ‘Physiologic vertical dimension and centric relation’. Tej Shanahan . *J Prosthet Dent* 1956. 6 p. .
- 122 [Bissau (2004)] ‘Pre-extraction records for complete denture fabrication: a literature review’. M Bissau . *J*
123 *Prosthet Dent* 2004 Jan. 91 (1) p. .
- 124 [Heartwell and Rahn ()] ‘Syllabus of complete dentures. 2 nd ed’. C M Heartwell , A O Rahn . *Philadelphia: Lea*
125 *& Febiger* 1974. p. .
- 126 [The glossary of prosthodontics terms J Prosthet Dent (2005)] ‘The glossary of prosthodontics terms’. *J Prosthet*
127 *Dent* 2005 Jul. 94 (1) .
- 128 [Silverman (1953)] ‘The speaking method in measuring vertical dimension’. M M Silverman . *J Prosthet Dent*
129 1953 Mar. 3 (2) p. .
- 130 [Hurst ()] ‘Vertical dimension and its correlation with lip length and interocclusal distance’. W W Hurst . *J Am*
131 *Dent Assoc* 1962. 64 p. .
- 132 [Millet et al.] *Vertical dimension in the treatment of edentulous patient*, C Millet , A Leterme , C Jeannin , P
133 Jaudoin .
- 134 [Lytle (1964)] ‘Vertical relation of occlusion by the patient’s neuromuscular perception’. R B Lytle . *J Prosthet*
135 *Dent* 1964 Jan-Feb. 14 (1) p. .