

Prevalence of Beta-Lactamase Producers in ICUs of a Tertiary Care Teaching Hospital of North India

Dr. Poonam Sharma¹, Dr. Sarbjeet Sharma² and Dr. Jasvir kaur³

¹ GuruRamdas Institute of Medical Sciences And Research, Vallah, Amritsar, Punjab

Received: 9 December 2014 Accepted: 2 January 2015 Published: 15 January 2015

Abstract

The aim of this study was to identify prevalence rates, sites and types of infection, the most common bacteria, and their antimicrobial resistance patterns, especially betalactamase producers present in Intensive care units (ICUs) of a tertiary care hospital. Methods: Clinical samples obtained over a period of 12 months from January 2014 to December 2014 from six ICUs of a tertiary care hospital were analyzed. Identification of micro organisms to species level and their antimicrobial susceptibility testing was performed as per CLSI guidelines. For control, 500 samples each from the environment staff of ICU were collected processed in the same way.

Index terms— intensive care units, prevalence, betalactamase producers, cross-infection.

1 I. Introduction

Intensive care units (ICUs) are of great value in the control and treatment of the severe illnesses that affect the human body. In spite of their invaluable and well-established role, ICUs lead to some degree of morbidity to patients, and nosocomial infection is clearly related to that.

Exposure to life-saving invasive procedures, serious underlying disease of patients and increased patient contact with healthcare personnel put patients admitted to the ICU at higher risk of acquiring nosocomial infections. ICU staff and the equipment used for patient care during the hospitalisation are the primary sources of cross-transmission of nosocomial pathogens.

Nosocomial infections increase patient morbidity, length of hospital stay, hospital costs, and may also increase mortality rates. When serious infections are suspected, treatment must be commenced immediately to increase the likelihood of a satisfactory outcome for the patient. Empirical knowledge, to select appropriate antibiotics, must be used so that the most likely infecting organisms are treated.

The prevalence of ICU-acquired infections varies between 4.4% and 88.9% being significantly higher in developing countries than in developed countries (1). Furthermore, device-associated infection rates in developing countries, especially ventilator-associated pneumonia (VAP) followed by central venous catheter-related bloodstream infections (CRBSIs), occur at a higher frequency than in European countries and USA (2,3).

The aim of this study was to identify the prevalence rates, sites and types of infection, the most prevalent microorganisms, and the antimicrobial resistance patterns especially the beta-lactamase producers present in ICUs of a tertiary care hospital.

2 II. Materials and Methods

The present study was conducted in 6 intensive care units of a tertiary care teaching hospital. We retrospectively analyzed consecutive culture-positive isolates and studied the antimicrobial susceptibility patterns of microorganisms during the period from January 2014 to December 2014. The BacT/Alert (bioMérieux) was used for blood culture. Identification of micro organisms to species level and antimicrobial susceptibility testing was performed by the Vitek method. The interpretation of antimicrobial susceptibility results was done as per Clinical

and Laboratory Standard Institute (CLSI) guidelines (4). For control, 500 samples each from the environment & staff of ICU were collected & processed in the same way. Informed consent was taken from staff before collecting samples. This study was conducted on hospitalised patients from whom samples were received in microbiology department so the article didn't need an approval from ethical committee.

3 a) Statistical Analysis

Statistical analysis was conducted. The chisquare test and odds ratio was applied for the comparison of categorical variables. P values less than 0.05 were considered as statistically significant.

4 III. Results

Total number of clinical samples collected during one year period were 1545. Out of which, 522 were culture positive as shown in Table1. The prevalence of Gram positive and Gram negative isolates in environmental & staff samples is shown in Table 3. The prevalence of β -lactamase producing Gram negative and Gram positive isolates in clinical, environmental & staff samples is shown in Table 4 & 5 respectively. (D D D D) C 9.

Beta lactamase producers were maximum in clinical samples followed by environmental and staff samples as shown in Table 6. 10. By calculating the odds ratio, isolates from clinical samples were twice more likely to be beta lactamase producers than the ones from environmental samples (odds ratio=2.98) and isolates from clinical samples were four times (odds ratio=3.94) more likely to be betalactamase producers than from staff samples. 11. By using chi-square test no significant ($p=0.2$) association was found between isolates from environmental and staff samples.

5 V. Discussion

In this study, Gram-negative bacteria were more common than Gram-positive bacteria, which is in accordance with the other studies (5,6,7). Treatment becomes challenging in gram-negative organisms causing serious infections in ICUs including pneumonia, bloodstream infections, wound or surgical site infections and meningitis. These organisms also exhibit multidrug resistance, and therapeutic alternatives have declined due to stagnation in novel antimicrobial agents (7,8).

In the present study, blood stream infections were most common, followed by urinary tract, wound and respiratory infections. However in the study by Boyles S (9) respiratory infections were more common in ICU followed by wound, blood stream and urinary tract infections. The probable reason could be that the sample size in their study was much larger (14000 patients) than ours.

Acinetobacter species was the most common pathogen isolated which is in accordance with the study by Datta P et al (6,7). The prevalence of MRSA among clinical, environmental and staff samples was more than 50% which shows that more stress should be laid on hand hygiene in our ICUs as improved hand hygiene can reduce acquisition of antimicrobial resistant bacteria, particularly MRSA (10).

There has been increase in incidence of beta lactam resistant organisms in ICUs of our hospital which is in accordance with a study conducted by Shaikh S et al (11). Among the clinical isolates, 50.8% of Gram negative isolates and 56.4% of Gram positive isolates were beta lactamase producers. This data confirms that in the intensive care medical environment, the most virulent in general are the most resistant and are the most frequently found organisms (12,13).

Prevalence of beta-lactamase producers in environmental and staff samples apart from clinical samples indicates towards the possibility of cross infection by these organisms from the environment and staff to the patients. Infection control strategies such as hand hygiene, rational antibiotic use, training and performance feedback can lead to significant reduction in the cross infection rates. In addition patients colonized or infected with beta lactamase-producing organism should be placed under contact precautions to avoid hospital transmission (14).

Hand hygiene is the most important way of decreasing the spread of infections in ICUs especially in developing countries like ours. Initial empirical therapy with broad-spectrum antibiotics is a life-saving strategy, which improves clinical outcome and reduces selection of resistant organisms. However, it is imperative to deescalate these antibiotics according to culture and antibiotic susceptibility results. Antibiotic cycling can be used as an effective approach to control antibiotic resistance. Strict antibiotic policies in ICUs can prevent the use of long term antibiotics and shorten the duration of the antimicrobial therapy (15).

Conducting infection surveillance and control activities in ICUs and rational antibiotic utilisation policies are valuable measures for infection control. These measures provide current knowledge about antibiotic resistance patterns, early recognition and management of outbreaks, which is essential for infection control (16).

Increasing drug resistance and spreading of multidrug-resistant (MDR) pathogens in the ICU environment results in limited therapeutic options and prolonged hospitalisations.

Consequently, the prevalence of ICU-acquired infection, healthcare costs and mortality rates are higher in developing countries due to limited resources associated with the quality of care (17).

In conclusion, our study shows that multidrug resistant bacteria are on rise in our hospital. The rise in beta-lactamase producers emphasizes the importance of stringent infection control practices, rational prescribing policies and need for development of new drugs and vaccines. Also, there is a need for the continuous evaluation of the local antibiotic resistance patterns for the formulation of a rational antibiotic policy. Limiting use of

antibiotics to patients with clear evidence of infection rather than colonization is essential & discontinuation of antibiotics when their possible benefits have been obtained is also critical. Further, new drugs are required to replace the increasingly obsolete classes of antibiotics that currently exist and there is need for strict compliance with infection control practices.

6 Volume XV Issue III Version I

1

S.no.	Samples	Total no.	Positive
1	Blood	704	158 (22.4%)
2	Urine	498	118 (23.7%)
3	Pus	160	124 (77.5%)
4	Suction tip	100	83 (83%)
5	Respiratory	47	33 (70.2%)
6	Body fluids	27	02 (7.4%)
7	Central lines	09	04 (44.4%)
		1545	522 (33.8%)

The percentage of Gram negative organisms isolated from clinical samples was79.88% (417/522), the percentage of gram positive organisms was 19.37% (101/522) & that of Candida was 0.76% (4/522) a shown in

Figure 1: Table 1 :

22

S.No.	ORGANISM	TOTAL No.
1	ACINETOBACTER	151
2	E.COLI	105
3	S. AUREUS	93
4	ENTEROBACTER	89
5	PSEUDOMONAS	29
6	KLEBSIELLA	21
7	CITROBACTER	16
8	ENTEROCOCCUS	06
9	PROTEUS	06
10	CANDIDA	04
11	CONS???	02
	Total	522
???Coagulase Negative Staphylococcus		

Figure 2: Table 2 .Table 2 :

1

3

Year 2 015

Volume XV Issue

III Version I

D D D D) C

(

S.No.	ISOLATES	Environment	Staff
1	Gram positive	50(10%)	34 (6.8%)
2	Gram negative	78(15.6%)	208 (41.6%)
3	Positive samples	128 (25.6%)	242 (48.4%)
4	Sterile samples	372 (74.4%)	258 (51.6%)
	Total samples	500	500

Figure 3: Table 3 :

4

SN	ORGANISMS	CLINICAL ISOLATES TOTAL	BETA LACTAMASES PRODUCING	ENVIRONMENTAL ISOLATES TOTAL	BETA LACTAMASES PRODUCING	ISOLATES FROM STAFF TOTAL	BETA LACTAMASES PRODUCING
1	ACINETOBACTER	88	(58.3%)	12	0	20	2 (10%)
2	E.COLI	105	56 (53.3%)	22	08 (36.3%)	52	10 (19.2%)
3	ENTEROBACTER	31	(34.8%)	14	02 (14.3%)	10	0
4	PSEUDOMONA	29	15 (51.7%)	10	02 (20%)	46	10(21.7%)
5	KLEBSIELLA	21	13 (61.9%)	20	10 (50%)	70	22 (31.4%)
6	CITROBACTER	16	09 (56.3%)	0	0	06	0
7	PROTEUS	06	0 (0%)	0	0	04	0
	Total	417	212 (50.8%)	78	22 (28.2%)	208	44 (21.2%)

Figure 4: Table 4 :

5

S.No.	ORGANISMS	CLINICAL ISOLATES		ENVIRONMENTAL ISOLATES	
		TOTAL	BETA	TOTAL	BETA
		LACTAMASES PRODUCING			
1	MSSA	36	17(47%)		
2	MRSA	57	40 (70%)		
3	CONS	2	0 (0%)		
4	ENTEROCOCCI	10	1 (10%)		
	Total	101	57 (56.4%)		

IV. Observations

- Maximum number of infections in ICU were Blood stream infections followed by Urinary tract, wound and

Figure 5: Table 5 :

6

ISOLATES	TOTAL PER- CENTAGE	GRAM POSITIVE%	GRAM NEG- ATIVE %	BETA LACTAMASES PRODUCERS
CLINICAL	33.8	19.4	79.8	51.9
ENVIRONMENTAL	25.6	10	15.6	26.56
STAFF	48.4	6.8	41.6	21.48

Figure 6: Table 6 :

.1 Conflict of Interest Statement

We declare that we have no conflict of interest.

[Shaikh et al. (2015)] ‘Antibiotic resistance and extended spectrum beta-lactamases: Types, epidemiology and treatment’. S Shaikh , J Fatima , S Shakil , Smd Rizvi , M A Kamal . *Saudi Journal of Biological Sciences* January 2015. 22 (1) p. .

[Rahal (2009)] ‘Antimicrobial resistance among and therapeutic options against gram-negative pathogens’. J J Rahal . *Clin Infect Dis* 2009 Aug. 49 (1) p. . (Suppl)

[Clinical and Laboratory Standards Institute: Performance standard for antimicrobial susceptibility testing CLSI document M100 ‘Clinical and Laboratory Standards Institute: Performance standard for antimicrobial susceptibility testing’. *CLSI document M100-S18. 9 th ed*, (Wayne, Pa) 2008. Clinical and Laboratory Standards Institute (Eighteenth Informational Supplement)

[Tutuncu et al. (2011)] ‘Device-associated infection rates and bacterial resistance in the intensive care units of a Turkish referral hospital’. E E Tutuncu , I Sencan , B Ozturk , G C Senturk , A U Kilic . *Saudi Med J* 2011 May. 32 (5) p. .

[Leblebicioglu et al. (2007)] ‘Device-associated hospital-acquired infection rates in Turkish intensive care units. Findings of the International Nosocomial Infection Control Consortium (INICC)’. H Leblebicioglu , V D Rosenthal , O A Arikan , A Ozgultekin , A N Yalcin , I Koksall . *J Hosp Infect* 2007 Mar. 65 (3) p. .

[Alp et al. (2012)] ‘Economic burden of ventilator-associated pneumonia in a developing country’. E Alp , G Kalin , R Coskun , M Sungur , M Guven , M Doganay . *J Hosp Infect* 2012 Jun. 81 (2) p. .

[Luna et al. ()] ‘Gram-Negative Infections in Adult Intensive Care Units of Latin America and the Caribbean’. Carlos M Luna , Eduardo Rodriguez-Noriega , Luis Bavestrello , Manuel Guzmán-Blanco . 10.1155/2014/480463. *Critical Care Research and Practice* 2014. 2014 p. 12.

[Health care associated infections fact sheet ()] *Health care associated infections fact sheet*, [pg1.www.who.int/gpsc/country_work/gpsc_ccisc_fact_sheet_en.pdf](http://www.who.int/gpsc/country_work/gpsc_ccisc_fact_sheet_en.pdf) 2011. WHO

[Datta et al. (2014)] ‘Health care associated infections: Risk factors and epidemiology form an ICU in northern India’. P Datta , H Rani , R Chauhan , S Gombhar , J Chander . *Indian J Anaesth* 2014 Jan-Feb. 58 (1) p. .

[Siegel et al. ()] *Health Infection Control Practices Advisory Committee. Management of multidrug-resistant organisms in healthcare settings*, J D Siegel , E Rhinehart , M Jackson , L Chiarello . <http://www.cdc.gov/ncidod/dhqp/pdf/ar/MDROGuideline2006.pdf> 2006.

[Alp et al. (2011)] ‘Importance of structured training programs and good role models in hand hygiene in developing countries’. E Alp , A Ozturk , M Guven , I Celik , M Doganay , A Voss . *J Infect Public Health* 2011 Jun. 4 (2) p. .

[Alp et al. ()] ‘Infection control practice in countries with limited resources’. E Alp , H Leblebicioglu , M Doganay , A Voss . *Ann Clin Microbiol Antimicrob* 2011. 10 p. 36.

[Boyles ()] *Infections are common in ICUs-Web MD*, S Boyles . www.webmd.com/healthy-aging/news.Dec32009.

[Derde et al. ()] ‘Interventions to reduce colonisation & transmission of antimicrobial-resistant bacteria in ICUs: an interrupted time series study & cluster randomised trial’. L P Derde , B S Cooper , H Goossens . *Lancet Infect Dis* 2014. 14 p. 31.

[León-Rosales et al. ()] ‘Prevalence of infections in intensive care units in Mexico: a multicenter study’. S P León-Rosales , F Molinar-Ramos , F Domínguez-Cherit . *Crit Care Med* 2000. 28 (5) p. .

[Hassanzadeh et al. (2009)] ‘Prevalent bacterial infections in ICUs Of Shiraz university of medical sciences teaching hospitals’. P Hassanzadeh , M Motamedifar , N Hadi . *Jpn J Infect Dis* 2009 Jul. 62 (4) p. .

[Vincent et al. ()] ‘The prevalence of nosocomial infection in intensive care units in Europe -results of the European Prevalence of Infection in Intensive Care (EPIC) Study’. J-L Vincent , D J Bihari , P M Suter . *JAMA* 1995. 274 (8) p. .