

Dr. Gadangi Indira¹

¹ Pingle Govt College for Women Warangal. A.P

Received: 6 December 2014 Accepted: 4 January 2015 Published: 15 January 2015

Abstract

The frequency of superficial fungal infections is more in hot and humid climate. Seasonal variation of skin diseases, a subject of much epidemiological interest, has been studied for centuries. The present study report is on dermatophytoses in different clinical conditions such as age, sex and seasonal impact in Warangal District, A.P. A retrospective study was done on all new outpatients visiting a single dermatologist Dr. Ramesh, at the Ramesh skin hospital, Warangal, Andhra Pradesh. The study population comprised of all new subjects visiting the outpatient department, from January to December, from 2008- 2010. The total number of cases was recorded according to the month wise prevalence.

Index terms— dermatophytosis, tinea or ringworm infections, dermatophytes, woods lamp and skin scrapings.

1 Introduction

dermatophytoses are a superficial cutaneous mycoses confined to the outer layers of skin, hair, and nails, and do not invade living tissues. The fungi are called dermatophytes. Dermatophytes, or more properly, keratinophilic fungi, produce extracellular enzymes (keratinases) that are capable of hydrolyzing keratin. These infections are commonly called as tinea infections or ringworm infections. Basing on the area or site of infections these are categorized as tinea capitis (plate-1), tinea corporis (plate-2), tinea manuum (plate-3), tinea cruris (plate-4), tinea facie (plate-5), tinea pedis (plate-6). Dermatophytes are by far the most significant fungi because of their widespread involvement of population at large and their prevalence all over the world. They are assuming greater significance both in developed and developing countries particularly due to the advent of immunosuppressive drugs and disease. Hot and humid climate in the tropical and subtropical countries like India makes dermatophytosis a very common superficial fungal skin infection. The prevalence of dermatophytic infections are governed by environmental conditions [18], personal hygiene [17], and individual susceptibility from place to place. The isolation of different dermatophytes also varies markedly from one ecological niche to another niche depending on their primary habitat [3]. The isolation of different dermatophytes also varies markedly from one ecological niche to another niche depending on their primary habitat [3]. The Warangal district in Andhra Pradesh (India) is predominantly a rural area with tropical climate. Though the ringworm infections are more prevalent, no systemic study and analysis has been made so far. The present investigations were undertaken to identify the size and magnitude of the dermatophytoses problem in this region. Further objectives were to:

i. study the incidence of dermatophytosis according to age and sex factors. ii. study the effect of seasonal variations in clinical types of dermatophytosis. iii. study the incidence of dermatophytoses in rural and urban areas.

2 II. Materials and methods

3 a) Study group

The present study was conducted in Warangal, which has favorable environment for development of superficial mycoses. The climate in the district is hot and humid for most of the year. It receives moderate to heavy rainfall in monsoon season. The present study was conducted on 400 clinically diagnosed patients with dermatophytoses who visited as out patients at Ramesh Skin Hospital (Dr.Ramesh, Dermatologist.) during the two-year period: January 2008 to December 2010. Most of the patients belong to low and middle socioeconomic groups coming

from Warangal town and surrounding villages of Warangal district. The data from the patients was collected by supplying a data sheet regarding name, age, sex, address, occupation, family history, socioeconomic background, duration of illness, personal contact at home, work place /school, previous medication like antifungal therapy, history of using immunosuppressive drugs and involvement of more than one site. After collecting the information the patients were examined regarding the lesions, types, and presence of inflammatory margins for apparent diagnosis of infection. If the outbreak is large a random sample was examined and data was collected.

4 b) Sample collection

The samples from patients were collected in aseptic conditions from infected areas such as skin, nail and hair (Robinson1988, Murray 1999)

5 Results and Discussions a) Age and sex wise prevalence of dermatophytoses

The results of analysis made through sex and age wise prevalence of dermatophytosis are presented in Table 1 and Table 2. Out of the 400 patients included in this study, 234 (67%) of the patients were male and 166 (33%) were female. Male to female ratio was 2.03:1. This statistical analysis was shown in Table ??1. This difference may be due to increased physical activity and more exposure to the out door activities leading to exposure to infectious etiological agents or fungi. Similar to the present observations, the study on dermatophytosis at Tirupati reported male preponderance with female ratio was 2:1. Dermatophytosis was more prevalent in men (63.4%) than in women (27). Similarly Gupta et al., (2001) have reported male preponderance with female ratio of 2.3:1. Males were affected at least 3 times more frequently than the females, according to the reports of Sentamilselviet al ??1997). In most of the other studies there was a predominance of male population than females. In the present study also there was male predominance over female population. This was due to more vulnerable infections such as higher exposure to army, school activities, increased sweating and types of shoes and socks they use [20,21]. ??19.5). Most of patients were in the age range of 20-29 years [27] which confirms to the results of some researches in Iran [15,1] But in the early years 0-10 years the cases reported were less 36 (9%) than the adolescent age 11-20 with 78 cases (19.5%) as the environmental exposure and physical activities were more in this age group. In the fifth and sixth decades of life there was a gradual decrease in the incidence of disease, so 28 cases (7%) were seen in 51 -60 and 22 cases (5.5%) in 61-70 years. Least incidence reported with only 2 cases (0.5%) in 71-80 age groups due to the less survival rate at this age group. Ranganathan et al ??2001). ??harma and Gupta (1983) in their investigations revealed that the maximum number of patients was in the age group of 20 -30 (30.7%). Table 1

This increased incidence of fungal infections may be due to the hormonal change and or increase in sebum secretion [7]. The age group most commonly affected was between 20 and 40 years of age. Females were affected more between the ages of 30 to 40 years based on an Indian study report by Sentamilselviet al ??1997). Similarly Verenkaret al ??1991), in their study on dermatophytosis reported that ringworm infection were common in third decade of life. Gupta et al (1993) [19, ??2]. Senet al, (2005) also reported the male predominance over females, although some authors found higher incidences in the second decade of age [6].

6 b) Seasonal incidence of dermatophytosis

Seasonal variation of skin diseases, a subject of much epidemiological interest, has been studied for centuries. The data was collected for two years ??2008) ??2009) ??2010) and the total number of the cases were recorded according to the month wise prevalence and were analyzed according to seasons. The seasonal variations and their impact on dermatophytosis were presented in the Table -4 and the critical analysis showed the high incidence of disease in month of April (49 cases, 12.25%) followed by May (43 cases, 10.75%) and June (40 cases, 10%). Correlated with seasonal variations majority of our patients with fungal infections were reported to be more in summer. It is known that warm, humid climates create the environment for the development of fungal infections as the temperature, humidity, ultraviolet radiation (UVR), flora and fauna all change with season [11]. The high prevalence of dermatophytosis in June month could be attributed to the extended summer season in Warangal district of Andhra Pradesh.

7 (C)

The frequency of fungal infections varies with seasons. The critical analysis of Table-5 reveals that, the highest number of cases of tinea pedis and tinea cruris occurred in the summer months, while tinea capitis, tinea corporis and tinea unguium occurred in the spring and winter months [28]. Our findings, Table-1 and fig- ??, even corroborate with the study report on dermatophytoses in Khammam district, Andhra Pradesh, India by Sumana and Singaracharya (2004). The incidence was more during the months of March to July in patients who were agricultural labourers living in rural areas. The incidence of dermatophytosis in the months of November with 34 cases and December with 36 cases can be explained, as the infections with anthrophilic species being commoner during summers while infections with zoophilic species peaking during the autumn and winter months (Jang, 2000). The higher prevalence of dermatophytoses of pets like cats and dogs, coupled with greater contact of

humans with their pets during the winter months have been presumed to result in common tinea infection by zoophilic species. The higher incidence of dermatophytic infections in winter than rainy season is explained that low temperature and lower humidity results in the extensibility, resistance to fissuring and hydration of the stratum corneum, thus contributing to damage of the epidermal barrier thus causing more lesions in winter..

8 Studies

of dermatophyteinfecti.

9 Distribution of dermatophytosis among rural and urban population .

The prevalence of tinea infections was more in urban area with 278 cases (69.5%) than in rural with 122 cases (30.5%) and the data was presented in Table-6. The incidence of dermatophytosis is very high among the patients of low socioeconomic status in rural population with 91.8% than in urban with 75.89%. This is due to the poor hygienic conditions and overcrowding. Even the people were infected with ringworm infections by soil (geophilic), animals (zoophilic) and contact by man (anthrophilic). In the middle income group the incidence of dermatophytosis was more in urbanites (19.42%) than the rural people ??8.19%). The reasons could be attributed to the high humid conditions due to concrete buildings and overcrowding of the population in urban areas. The prevalence of dermatophytoses in urban and rural schools was reported to be 14.3% and 10% respectively [29]. Our study report is similar with the report on extent and pattern of pediatric dermatoses in rural areas of central India by Vikas Bhatia (1989). The nutrition also has a significant role in the incidence of ringworm infections. The rural children who were lacking the sufficient diet were more prone to the tinea infections in Udaipur district [5].

Children of primary school age are usually between the ages of 3 to 14 years or above in rural areas, and thus, are more susceptible to various infections due to their close contact with each other and low immune status when compared with adults. Some other factors such as enlightenment, customs, and tradition of people, hygiene level and environmental sanitary conditions may influence the prevalence of dermatophytosis [2]. Also, their inability to keep themselves clean always, and their frequent contact with soil and infected pets like dogs and cats at home. The study was carried out to determine the prevalence of dermatophytosis among primary school children [8] at Nigeria, revealed that close contact with the soil and pets increased the incidence of dermatophytoses among the school children.

10 D D D D

IV.

11 Conclusion

The study revealed that the ringworm or tinea infections are very common in the age group of 21-30 years and less common in two extremities of the age group i.e., children and old age people. The reason for this incidence of dermatophytosis can be attributed to the increased physical activities and increased opportunity for exposure and due to the hormonal change. The male preponderance was observed in ringworm infections with male to female ratio as 2.03:1. The increased incidence in the males might be explained as increased outdoor physical activities and exposure to infectious etiological agents, as mostly they are the breadwinners of the family. Prevalence tinea infections were observed to be more in summer season than in winter. It is known that warm, humid climates create the ideal environment for the development of fungal infections. The high prevalence of dermatophytosis in June and July months could be attributed to the extended summer season in Warangal district of Andhra Pradesh. The prevalence of tinea infections were found to be more in urban area with 278 cases (69.5%) than in rural with 122 cases (30.5%). But the incidence of dermatophytosis is very high among the patients of low socioeconomic status in rural population with 91.8% than in urban with 75.89%. This is due to the poor hygienic conditions and overcrowding. In the middle income group the incidence of dermatophytosis was more in urbanites (19.42%) than the rural people ??8.19%). The reasons could be attributed to the high humid conditions due to concrete buildings and overcrowding of the population in urban areas.

V. ^{1 2}

¹© 2015 Global Journals Inc. (US)

²Volume XV Issue 2 Version I © 2015 Global Journals Inc. (US) Year 2015



Figure 1: Figure :

Plate-1: Tineacapitis



Plate-3 TineaMannum



Plate-5 Tinea facie



Plate-2: Tineacorporis



Plate-4: Tineacruris



Plate-6 Tineapedis



Figure 2:

Year 2015
 Volume XV Issue 2 Version I
 Total no. of cases = 400; Total no. of males = 234
 (67%);
 Total no.of females = 166 (33%)

Figure 3: :

(C)			
234	166	400	2.03:1
Percentage 67	33	100	

Figure 4: Males Females Total Ratio of male/female No. of cases

2

Figure 5: Table 2 :

Age(years) No.	of cases	Percentage
0-10 yrs	36	9.0
11-20 yrs	78	19.5
21-30 yrs	102	25.5
31-40 yrs	78	19.5
41-50 yrs	54	13.5
51-60 yrs	28	7.0
61-70 yrs	22	5.5
71-80 yrs	02	0.5
Total	400	100
		(C)

Figure 6:

3

Figure 7: Table 3 :

61-70	Male	13	59.09
yrs	Female	09	40.91
71-80	Male	02	100
yrs	Female	nil	0
The incidence of dermatophytosis gradually declined in fifth and sixth decades of life with males 18 (64.28%), females 10 (35.71%) in 51-60 yrs age group and were 13 in 70 the age group of 0-10 yrs male			
Year were 22 (61.1%) and females 14 (38.98%). The least incidence of infection was recorded in the age group 2015			
males. In a study report by Kamothi (2010) on prevalence of dermatophyte infection in Rajkote District, Volume 10, young adult in age group of 21-30 years were mainly affected. Male to female ratio was 2.03:1. In this study			
XV			
Is-			
sue			
2			
Ver-			
sion			
I			
	Male	22	61.1
	Female	14	38.98
11-20	Male	46	58.97
yrs	Female	32	41.02
21-30	Male	57	55.88
yrs	Female	45	44.11
31-40	Male	45	57.69
yrs	Female	33	42.30
41-50	Male	31	57.40
yrs	Female	23	42.59
51-60	Male	18	64.28
yrs	Female	10	35.71

Figure 8: Age group Sex No.of cases Percentag e Total no. of cases Total percentage 0-10 yrs

4

Month	No. of cases *	Percentage
January	33	8.25
February	27	6.75
March	38	9.5
April	49	12.25
May	43	10.75
June	40	10
July	32	4
August	25	6.25
September	21	5.23
October	22	5.5
November	34	8.5
December	36	9

Figure 9: Table 4 :

5

Clinical types	Tinea corporis	Tinea cruris	Tinea inguinum	Tinea pedis	Tinea capitis	Tinea manuum	Tinea faciei	Total cases
Summer (March-July)	32	43	28	48	32	16	03	202
Winter (Nov-Feb)	19	21	23	20	34	12	01	130
Rainy season (Aug-Oct)	12	14	11	10	14	07	-	68

Figure 10: Table 5 :

6

Socio economic Status	No. of cases in rural	Percentage	No. of cases in urban	Percentage
Low income	112	91.80	211	75.89
Middle income	10	8.19	54	19.42
High income	-	-	13	4.67
Total no. of cases	122	30.5	278	69.5

Figure 11: Table 6 :

.1 Acknowledgment

This research work was carried under the supervision of my guides Prof. S. Ram Reddy, Department of Microbiology and Prof. G. Raghu ramulu, Department of Zoology, Kakatiya University, Warangal (India) and I extend my heartfelt thanks to them. Thanks are also due to Dr.Ramesh, dermatologist Warangal (India) for facilities and encouragement.

- [Sentamilselvi et al. ()] , G Sentamilselvi , A Kamalam , K Ajithadas , C Janaki , A S Thambiah . 1998. 1997-1998.
- [Gupta et al. ()] , A K Gupta , Y Kohli , J Faergemmann , R C Sumerbell . *Epidemiology of Malassezia yeast associated with Pityriasis Versicolor on Ontario. Can.Med. Mycol* 2001. p. .
- [Sen and Rasul ()] , S S Sen , E S Rasul . *Dermatophytosis in Assam. Ind. J. Med. Microbiol* 2006. 24 p. .
- [Aghamirian and Hashemi ()] ‘A survey of dermatophytosis in patients referring to Dermatology Clinic in Ghazvin Avicenna Hospital in 2001-2002’. D Aghamirian , H J Hashemi . *Proceedings of the 5th National Iranian Congress on Parasitic Diseases*, (the 5th National Iranian Congress on Parasitic Diseases) 2002. p. . ShahidBeheshti University of Medical Sciences
- [Bindu and Pavithran ()] ‘Clinico -mycological study of dermatophytosis in Calicut’. V Bindu , K Pavithran . *Indian J DermatolVenereolLeprol* 2002. 68 p. .
- [Verenkar et al. ()] ‘Clinico-microbiological study of dermatophytoses. Ind’. M P Verenkar , M J Pinto , S Rodrigues , W P Roque , I Singh . *J. Pathol.Microbiol* 1991. 34 p. .
- [Chander ()] ‘Clinicomycological profile of dermatophytosis in Jaipur’. J Chander . *Mehta publishers, New Delhi. Epidemiologicalstudies on Dermatophytoses in Warangal District, Telangana State*, India 12 Jain, N, M Sharma, V N Saxena (ed.) 2002. 2008. 74 p. . (Text Book of Medical Mycology)
- [Peerapur et al. ()] ‘Clinicomycological study of dermatophytosis in Bijapur. Ind’. B V Peerapur , Ac , P V Inamadar , B Pushpa , Srikant . *J. Med. Microbiol* 2004. 22 p. .
- [Robinson and Pandhye (ed.) ()] *Collection, transport and processing of clinical specimens*, B E Robinson , A A Pandhye . B.B. Wentworth (ed.) 1998. (Diagnostic procedures for mycotic)
- [Sentamilselvi et al. ()] ‘Deep dermatophytosis caused by Trichophytonrubrum-a case report’. G Sentamilselvi , C Janaki , A Kamalam , A S Thambiah . *Mycopathologia* 1998. 142 p. .
- [Patwardhan and Dave (1999)] ‘Dermatomycosis in and around Aurangabad’. N Patwardhan , R Dave . *Ind. J. Pathol. Microbiol* 1999. Oct. 42 (4) p. .
- [Sumana and Ma Singaracharya ()] ‘Dermatophytes in Khammam (Khammam district’. V Sumana , Ma Singaracharya . *Andra Pradesh. India). Ind. J. Pathol. Microbiol* 2004. 47 p. .
- [Chukwu et al. (2011)] ‘Dermatophytoses in rural school children associated with livestock keeping in Plateau State’. D Chukwu , O O C Chukwu , A Chuku , B Israel , B I Enweani . *Nigeria Journal of Yeast and Fungal Research* 2011. January 2011. 2 (1) p. .
- [Sepahvand et al. ()] ‘Dermatophytosis in Western Part of Iran’. A Sepahvand , J Abdi , Y Shirkhani , Sh , M Fallahi , S Tarrahi , Soleimannejad . *Khorramabad.Asi. J. Biological Sciences* 2009. 2 p. .
- [dermatophytosis: an Indian study] *dermatophytosis: an Indian study*, 140 p. .
- [Shahindokhtbassiri-Jahromi et al. ()] ‘Epidemiological survey of dermatophytosis in Tehran’. Ali Shahindokhtbassiri-Jahromi , ; | Asgharkhaksari , Page . *Ind. J. Dermatol. Venereol. Leprol* 2009. 2000 to 2005. 75 p. .
- [Aly et al. ()] ‘Epidemiology of tineacapitis’. R Aly , Rj , A Hay , R Palacio , Galimberti . *Med Mycol* 2000. 38 p. .
- [Vikas ()] ‘Extent and pattern of pediatric dermatoses in rural areas of Central India’. B Vikas . *Ind. J. Dermatol. Venereol. Leprol* 1997. 63 p. .
- [Ranganathan and Gokulsanker ()] *Fungal diseases of the skin*, S Ranganathan , M S Gokulsanker , Ranjith . 2001. (The Hindu on line edition of Indian national news paper)
- [Padhye et al. ()] ‘Hair perforation as a diagnostic criterion in the identification ofEpidermophyton,Microsporum, andTrichophyton species’. A A Padhye , L Young , Ajello . *Pan. Am Health Org Sci Publ* 1980. 396 p. .
- [Gupta et al. ()] ‘Mycological aspects of dermatomycoses in Ludhiana’. B K Gupta , S Kumar , R Kumar , S Khurana . *Ind. J. Pathol. Microbiol* 1993. 36 p. .
- [Bhandari and Jain ()] ‘Nutrition and health status of rural school boys of Udaipur district’. B Bhandari , A M Jain . *Ind. J. Paed* 1975. 42 p. 186.
- [Sharma and Garg ()] ‘Pattern of skin diseases in urban school children’. N K Sharma , N Garg . *Ind. J Dermatol . Venereol .Leprol* 1986. 6 p. .

11 CONCLUSION

- [Kamothi ()] 'Prevalence of dermatophytic infection in District Rajkote'. M Kamothi . *Electronic Journal of Pharmacology and Therapy* 15. Mohmoudabadi A.Z. (ed.) 2010. January 1. 2010. 2005. 160 p. . (Mycopathologia.)
- [Hay et al. ()] *Rook's Textbook of Dermatology*, R J Hay , Mk , Moore , T Burns , S Breathnach , N Cox , C Griffiths . Science: pp-31.1-31.101. infections. 2004. Washington DC: American public health association. 2 p. . (Mycology. 7th edn)
- [Phipot ()] 'Some aspects of the epidemiology of tinea'. C M Phipot . *Mycopathologica* 1977. 62 p. .
- [Sharma and Gupta ()] 'Superficial mycoses in shimla'. L N Sharma , . L Gupta . *Indian J DermatolVenereolLeprol* 1983. 49 (6) p. .
- [Shokohi et al. ()] 'The study of 101 cases of onychomycosis and associate factors in patient referred to BoaliSina Hospital and Toba dermatology outpatient clinics'. T Shokohi , Z Hajheidari , I Haghani , A R Khalilian , S R Agilli , S Mayahi . *Sari . J. Mazandaran University of Med. Scie* 2009. 19 (71) p. .
- [Oyeka ()] 'Tineacapitis in Awka local Government Area of Anambra state'. C A Oyeka . *West Afr . J. Med* 1990. 9 p. .
- [Jang et al. ()] 'Tineapedis in Korean children Int'. K A Jang , D H Chi , J H Choi , K J Sung , K C Moon , J K Koh . *J. Dermatol* 2000. 30 p. .
- [Alabi ()] 'Trends in the Pattern of Skin Diseases in Nigeria'. G O Alabi . *Nig. Med. J* 1980. 10 (5) p. .
- [Murray et al. ()] 'Trichophyton, Microsporum, Epidermophyton and Agents of Superficial Mycoses'. P R Murray , E J Baron , M A Pfaller , F C Tenouer , R H Youlken . *Manual of Clinical Microbiology* 1999.
- [Asawananda et al. ()] 'Wood's light in dermatology'. P Asawananda , Cr , Taylor . *Int. J. Dermatol* 1999. 38 p.