



GLOBAL JOURNAL OF MEDICAL RESEARCH: G
VETERINARY SCIENCE AND VETERINARY MEDICINE
Volume 15 Issue 2 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

A Cross-Sectional Survey of Bovine Fasciolosis at Elkadaro Abattoir, Khartoum State, Sudan

By Mohammed B. Badreldeen & Abdelhamid A. Elfadil
Sudan University of Science and Technology, Sudan

Abstract- A cross-sectional study was conducted from May to July 2011. The multivariate analysis exposed significant associations by postmortem examination: age >4 years ($P=0.000$, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed ($P=0.000$, OR= 77.6, 95% CI= 9.3, 81.6), light weight ($P=0.000$, OR= 3.0, 95% CI= 1.9, 5.3), Ethiopian cattle ($P=0.000$, OR= 76.1, 95% CI= 8.4, 83.7) and small size ($P=0.000$, OR= 3.1, 95% CI= 1.9, 5.3). while, the coprology demonstrated significant associations among: age >4 years ($P=0.000$, OR= 28.8, 95% CI= 3.9, 34.2), foreign breed ($P=0.000$, OR= 94.4, 95% CI= 13.2, 102.6), light weight ($P=0.000$, OR= 53.4, 95% CI= 7.1, 61.3), Ethiopian cattle ($P=0.000$, OR= 60.2, 95% CI= 8.3, 70.1) and small size ($P=0.000$, OR= 54.9, 95% CI= 6.3, 63.8). A higher prevalence was recorded by postmortem examination ($X^2=1.669$, $P=0.000$). This study determined the prevalence of bovine fasciolosis.

Keywords: fasciolosis, cattle, prevalence, risk factors, abattoir, khartoum, sudan.

GJMR-G Classification : NLMC Code: QW 70



Strictly as per the compliance and regulations of:



A Cross-Sectional Survey of Bovine Fasciolosis at Elkadaro Abattoir, Khartoum State, Sudan

Mohammed B. Badreldeen ^α & Abdelhamid A. Elfadil ^ο

Abstract- A cross-sectional study was conducted from May to July 2011. The multivariate analysis exposed significant associations by postmortem examination: age >4 years (P= 0.000, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed (P= 0.000, OR= 77.6, 95% CI= 9.3, 81.6), light weight (P= 0.000, OR= 3.0, 95% CI= 1.9, 5.3), Ethiopian cattle (P= 0.000, OR= 76.1, 95% CI= 8.4, 83.7) and small size (P= 0.000, OR= 3.1, 95% CI= 1.9, 5.3). while, the coprology demonstrated significant associations among: age >4 years (P= 0.000, OR= 28.8, 95% CI= 3.9, 34.2), foreign breed (P= 0.000, OR= 94.4, 95% CI= 13.2, 102.6), light weight (P= 0.000, OR= 53.4, 95% CI= 7.1, 61.3), Ethiopian cattle (P= 0.000, OR= 60.2, 95% CI= 8.3, 70.1) and small size (P= 0.000, OR= 54.9, 95% CI= 6.3, 63.8). A higher prevalence was recorded by postmortem examination ($\chi^2= 1.669$, P= 0.000). This study determined the prevalence of bovine fasciolosis.

Keywords: fasciolosis, cattle, prevalence, risk factors, abattoir, khartoum, sudan.

I. INTRODUCTION

Fasciolosis is among the most neglected important tropical diseases. Although it has significant economic impact on livestock industry, particularly cattle and sheep and occasionally can infects human beings (CDC, 2013). Among the estimated 91.1 million humans at risk for infection worldwide, as many as 17 million may be infected (Tolan, 2011). However, the disease is also consider as one of the major parasitic diseases contributing to loss in productivity estimated at over 200 US\$ million per annum worldwide (Sturat, 1998). The infection is due to the food- and water-borne route. The two species most commonly implicated, as the etiological agents of fasciolosis are *F. hepatica* and *F. gigantica* (CDC, 2013).

Sudan possesses one of the highest livestock populations in Africa but productivity is low as a result of diseases, malnutrition and other management problems. Fasciolosis is one of these diseases which is responsible for considerable economic losses in livestock production (Boray, 1985). The disease is proved to be endemic in certain districts that characterized by intensive sheep or cattle production, in addition to the existence of favorable habitats for the snails host, like: White Nile, Eljazeera and Sennar

regions (Ali, 1983). Another negative economic impacts on indigenous livestock result from inefficient conversion of feed, retarded growth, death, condemnation of infected livers, cost of preventive and treatment programs, reduced production, predisposition to other diseases, restricted use of infested lands and protein deficiencies among livestock dependant people (Saad, 2004).

Diagnosis is made serologically most often, although fecal examination for the eggs is fruitful if obtained when the adult worm is laying eggs (Tolan et al., 2011).

Therefore, the present study was designed to estimate the prevalence of bovine fasciolosis among cattle slaughtered at Elkadaro Abattoir, to identify potential risk factors associated with the occurrence of fasciolosis and to evaluate the accuracy of fecal examination.

II. MATERIALS AND METHODS

a) Study Abattoir

Elkadaro Abattoir was designed for a processing capacity of 30 cattle per hour, with shifting work system every 10 hours (two shifts per day), in five days weekly. The abattoir was classified as code no.1, certified via O.I.E categories. The plant is situated within an open free disease area (recognized as free zone referring to O.I.E scientific terms). The main task of the abattoir is to process fresh and frozen meat, mainly for export orders. The abattoir is managed directly by the Federal Ministry of Animal Resources and Fisheries.

b) Duration of Study

A field investigation was launched in Sunday 21/ May and completed in Saturday 23/ July 2011.

c) Study Animals

According to the latest estimation of the livestock population in Khartoum State 2010, about 33800 of cattle are raised. The target animals were provided from different sites in and out of Sudan. The local cattle were fetched from: Nyala, Kordofan, Kosti, Kassala, Eljazeera, Khartoum and Upper Nile (South Sudan). While, the foreign ones were from Ethiopia (eastern neighboring country).

d) Sample Size Determination and Sampling Methods

The sample size was calculated using the formula: $4PQ/L^2$, given by (Martin et al., 1988) where:

Author α : College of Veterinary Medicine, Sudan University of Science and Technology, P.O. Box: 204, Khartoum north 13314, Sudan.
e-mails: mohammed_bushra2000@yahoo.com,
aamelfadil@yahoo.com

P= prevalence

Q= 1-P

L²= allowable error, for systematic random sampling with 7.4% reported prevalence (Eldoush, 1995) and 3% allowable error. Accordingly, the sample size was determined to be 307.

The sampling procedure was carried out in such a way that from daily 120 slaughtered cattle, 10 were randomly selected. There are five slaughter days a week and accordingly, 50 cattle were examined weekly. Hence, 307 cattle were examined within two months of the study period. A fresh feces was collected instantly after slaughtering of the selected animals. Then livers were subjected to detailed postmortem examination.

e) Study Methodology

Coprology: Fecal samples for parasitological examination were collected directly from the rectum of each animal immediately after slaughter using disposable plastic gloves and placed in new plastic bags. Prior to slaughter, each selected animal was given an identification number. Then each fecal sample was clearly labeled with the cattle identification number. Samples were kept at the room temperature and examined fresh. In laboratory, coproscopic examinations were performed to detect *Fasciola* eggs using standard sedimentation technique as previously described (Coles, 1986).

Liver Inspection: Liver of each cattle was strictly examined for the presence of liver flukes separately to correlate the coprology and postmortem examination of each animal. Examination of livers for *Fasciola* was carried out immediately after removal of liver from abdominal cavity. The inspection was made according to the procedures certified by FAO (2003).

f) Sensitivity and Specificity of the Fecal Examination Method

One of the objectives of this study was to evaluate the accuracy of the direct coprological examination method, which is routinely employed at field to examine the presence of *Fasciola* species eggs in feces. The sensitivity and specificity of the method was computed by taking liver inspection at postmortem as gold standard for the diagnosis of fasciolosis. Kappa statistic was used to determine the degree of agreement between the two methods of liver fluke diagnosis. The kappa value was interpreted as: slight agreement (K <0.2); fair agreement (K= 0.2→0.4); moderate agreement (K= 0.4→0.6); substantial agreement (K= 0.6→0.8); and almost perfect agreement (K >0.8) (Thrusfield, 2005).

g) Data Management and Analysis

Both fecal examination and liver inspection results were recorded on specially designed forms and preliminary analysis was done in Microsoft Excel. The

outcome variable was the cases of fasciolosis detected during routine postmortem inspection (positive or negative) and fecal examinations for *Fasciola spp* eggs (positive or negative). Descriptive statistics were carried out to summarize the prevalence and proportion of infection in each category of investigated potential risk factors. Univariate and multivariate logistic regression analysis were conducted to see the significance and strength of association between potential risk factors and the occurrence of the infection. 95% confidence interval and p-value (P=<0.05) were used to notice the significance of association. Also, Odds Ratios (Exp B) was employed to assess the strength and direction of this association using SPSS statistical software (SPSS 16.0).

III. RESULTS

a) Abattoir (Postmortem) Prevalence

Of the total 307 slaughtered cattle that subjected to detailed postmortem examination at Elkadaro Abattoir, 31.6% (97/307) were found positive for fasciolosis (Table I). The highest prevalence was recorded in age greater than 4 years (44.5%), foreign breed (64.1%), light weight (47.4%), Ethiopian source (65.5%) and small size (47.8%) (Table III).

b) Risk Factor Analysis for the postmortem results

The occurrence of fasciolosis significantly varied with age, breed, weight, source and animal size (P=<0.05). The likelihood of fasciolosis occurrence was significantly higher in age >4 years (P=0.000, OR= 18.5, 95% CI= 3.1, 21.7), foreign breed (P=0.000, OR= 77.6, 95% CI= 9.3, 81.6), light weight (P=0.000, OR= 3.0, 95% CI= 1.9, 5.3) Ethiopian source (P=0.000, OR=76.1, 95% CI=8.4, 83.7), Kassala source (P=0.027, OR= 1.6, 95% CI= 1.1, 3.1), small size (P=0.000, OR=3.1, 95% CI= 1.9, 5.3) (Table III).

c) Prevalence by Coprology

Of the total 307 collected fecal samples 20.2% (62/307) were positive for coprological examination by sedimentation technique (Table II). The highest prevalence was recorded in age >4 years (28.9%), foreign breed (42.1%), light weight (43.8%), Ethiopian source (43.0%) and small size (44.1%) (Table IV).

d) Risk Factors Analysis for the Coprological Results

The results of coprological examination revealed significant association (P=<0.05) between the occurrence of fasciolosis and the risk factors: age, breed, source, weight and animal size. The likelihood of fasciolosis occurrence was significantly higher in age >4 years (P= 0.000, OR = 28.8, 95% CI = 3.9, 34.2), foreign breed (P= 0.000, OR = 94.4, 95% CI= 13.2, 102.6), Ethiopian source (P= 0.000, OR = 60.2, 95% CI = 8.3, 70.1) light weight (P= 0.000, OR = 53.4, 95% CI= 7.1, 61.3) and small size (P= 0.000, OR = 54.9, 95% CI= 6.3, 63.8) (Table IV).

e) *Difference in Prevalence between the Two Diagnostic Methods*

Based on the proportions comparison test there was significant difference ($X^2=1.669$, $P=0.000$) between fasciolosis prevalence estimated by coprology and postmortem examinations. Hence, in this study, higher prevalence of infection was observed by postmortem examination (31.6%) than by coprology (20.2%) (Table V).

f) *The Sensitivity and Specificity of the Fecal Examination Technique Considering the Presence of Fasciola spp in the liver as a Gold Standard Test*

As indicated in (Table VI), no animal that was positive with fecal examination and negative during postmortem examination. This revealed that postmortem examination was the golden test for diagnosis of fasciolosis when compared with coprology. The table set out the number of positive and negative tests in animals with and without flukes in their livers (Smith, 1995). The sensitivity and the specificity of fecal examination were found to be 63.9% and 100%, respectively. The calculated Kappa value ($Kappa=0.69$) indicated substantial agreement between the two techniques.

IV. DISCUSSION

Fasciolosis is a wide spread ruminant health problem and causes significant economic losses to the livestock industry in some areas in Sudan. The abattoir prevalence of fasciolosis obtained from the present study (31.6%) is very high compared to 7.4% (Eldoush, 1995) and nearly similar with 30% (Elmannan, 2001) and slightly lower than 34.4% (Abu-rigaila, 1983). These differences within the country could be attributed mainly to variations such as altitude, rainfall and temperature, although differences in livestock management system and the ability of the meat inspectors to detect the infection may play a part (Abu-rigaila, 1983). From African countries, a higher prevalence of 63.8% from Tanzania (Keyyu et al., 2006) and 53.9% from Zambia (Phiri et al., 2006) were reported. The observed prevalence may reflect suitable ecological and climatic conditions for the snail intermediate host in the areas from which the study animals came from.

Regarding the risk factors analysis, the results of this study indicate that the occurrence of fasciolosis in cattle varied with sex, age groups, breeds, weights, sources, size of animals and other diseases concurrent with fasciolosis.

The association between the occurrence of bovine fasciolosis and sex of the animals by both coprological and postmortem examinations revealed that the prevalence of *Fasciola* infection was found to be higher in males in agreement with a previous report (Shiferaw et al., 2009). The significant effects of sex on the prevalence of bovine fasciolosis might be attributed

to the management system in which males are kept outdoor while females are kept indoor at the beginning of lactation (Balock et al., 1985).

Higher *Fasciola* infection rate was recorded among older cattle (>4 years) with both postmortem and coprological examinations in agreement with a previous report (Andrade et al., 2002). The higher prevalence in older animals by both examinations could be associated with the degree of exposure to the parasite which is normally greater in old animals than young animals. In contrast to our finding, higher prevalence in young cattle than older ones has been reported (Mulugeta et al., 2011).

The current study determined a higher prevalence of *Fasciola* infection among Ethiopian zebu cattle by both postmortem and coprological examinations in agreement with previous reports (Kassaye et al., 2012) and (Chakiso et al., 2014). This might be attributed to the difference in resistance to parasitic infection between different breeds (Tasawar et al., 2007).

In this study a higher prevalence of fasciolosis was observed among Ethiopian source animals by postmortem and coprological examinations in consistency with another finding (Yilma et al., 2000). The higher prevalence of the disease among Ethiopian source animals could be associated with the existence of suitable ecological conditions for the intermediate snail host in the areas where animals graze (Abebe et al., 2010).

We determined a higher prevalence of *Fasciola* infection rate among light weight animals with both postmortem and coprological examinations in agreement with previous reports (Kassaye et al., 2012) and (Nega et al., 2012). This signifies that the light weight animals are more susceptible to the infection.

The results of our study also showed a higher infection rate among small size animals by postmortem and coprological examinations in agreement with a previous report (Bekele et al., 2012). This attributed to low resistance against the disease.

In this study the association between the occurrence of fasciolosis with concurrent infections with both coprological and postmortem examinations revealed that there is no significant difference between the concurrent infections with fasciolosis by both examinations.

The prevalence of fasciolosis reported by using coproscopy was lower than that obtained by the abattoir results indicating that the latter is more sensitive in detecting the disease. The detection of *Fasciola* eggs can be unreliable as the eggs are expelled intermittently, depending on the evacuation of the gall bladder (Briskey et al., 1998). Similar study suggested that about 36% infected animals may pass undetected with single fecal examination technique. This might be attributed partly to the fact that *Fasciola* eggs only appear in feces 8-15

weeks post infection, so most of pathological lesions had already occurred (Mulugeta et al., 2011).

The present sensitivity value (63.9%) is comparable to other reports: 65.9% from Ethiopia (Regassa et al., 2012) and 69% from Switzerland (Rapsch et al., 2006). The latter stated that traditional coproscopy can be very efficient if there is repeated sampling, resulting in sensitivity of approximately 92%. Therefore, worm counts at liver necropsy can only be considered as a gold standard if the livers are sliced and soaked. Even then very light or prepatent infections could still be missed, affecting the calculated sensitivity and specificity of the evaluated tests.

The current study revealed that the infection with fasciolosis varies according to different regions in Sudan and demonstrated that the disease had a high prevalence among Ethiopian cattle. It also elucidated that coprological examination for the parasite eggs has significant limitations in detecting the presence or absence of fasciolosis in animals. On the other hand, it has helped to illustrate the usefulness of meat inspection in monitoring disease situation and demonstrating possible long term trends. This study also showed that bovine fasciolosis is significantly associated with age, breed, weight, source and size of animal.

V. ACKNOWLEDGEMENTS

Special appreciation to Elkadaro Abattoir officials and all the workers for their willingness and collaboration which was indispensable inputs for the accomplishment of this study. All contributions and supports are gratefully acknowledged.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Abebe, R., F. Abunna., M. Berhane ., S. Mekuria., B. Megersa., and A. Regassa. 2010. Fasciolosis: Prevalence, financial losses due to liver condemnation and evaluation of a simple sedimentation diagnostic technique in cattle slaughtered at Hawassa Municipal abattoir, Southern Ethiopia. *Ethiopian Veterinary Journal* **14**:39-5.
2. Abu-rigaila, H. A. 1983. Studies on naturally occurring and experimental ovine Fasciolosis in Sudan. M.V.Sc Thesis. University of Khartoum, 7 p.
3. Ali, M. A. 1983. Epidemiological studies on bovine fasciolosis in the White Nile Province. M.V.Sc Thesis. University of Khartoum, 5 p.
4. Andrade, S. R., A. Paz-Silva., L. J. Suarez., R. Panadero., J. Pedreira., C. Lopez., P. Diez-Banos., and P. Morondo. 2002. Influence of age and breed on natural bovine fasciolosis in an endemic area (Galicia, NW Spain). *Veterinary Research Communications* **26**:361-370.
5. Balock, F. C., and R. J. Arthur. 1985. A survey of fasciolosis beef cattle killed at abattoir in southern Queensland. *Australian Veterinary Journal* **62**:324-326.
6. Bekele, M., H. Tesfay., and Y. Getachew. 2012. Bovine Fasciolosis: Prevalence and its economic loss due to liver condemnation at Adwa Municipal Abattoir, North Ethiopia. *Ethiopian Journal of Applied Science and Technology* **3**:319-324.
7. Boray, J. C. 1985. Flukes of Domestic Animals. In: Gaafar, S.M., Howard, W.S. and R.E. Marsh, (Eds.) *Parasites, Pests and Predators*. Elsevier 178-218.
8. Briskey, D. W. 1998. Diagnosis of liver fluke infections in cattle, *Veterinary Bulletin* 68-72.
9. Centre of Disease Control and Prevention. Va. USA. 2013. Available @www.dpd.cdc.gov/dpdx/html/fasciolosis.htm.
10. Chakiso, B., S. Menkir., and M. Desta. 2014. On farm study of bovine fasciolosis in Ilemo districts and its economic loss due to liver condemnation at Hossana Municipal Abattoir, Southern Ethiopia. *International Journal of Current Microbiology and Applied Sciences* **3**:1122-1132.
11. Coles, E. 1986. *Veterinary Clinical Pathology*. Professor of Clinical pathology and Immunology. 4th ed. Kansas State University, 374-376 p.
12. Eldoush, M. E. 1995. Some studies on liver and gastro-intestinal parasites of cattle at Atbara town, Northern State, Sudan. M.V. Sc Thesis. University of Khartoum, 6 p.
13. Elmannan, A. M. 2001. Studies on biological control of lymnea natalensis snails using a molluscicidal indigenous plant pulicaria crispa (forsk) oliv compositae. Ph. D Thesis. University of Khartoum, 9 p.
14. FAO. 2003. *Diagnostic Manual on Meat Inspection for developing countries*.
15. Kassaye, A., N. Yehualashet., D. Yifat., and S. Desie. 2012. Fasciolosis in slaughtered cattle in Addis Ababa abattoir, Ethiopia. *Global Veterinaria* **8**:115-118.
16. Keyyu, J. D., A. A. Kassuku., L. P. Msalilwa., J. Monrad., and N. C. Kyvsgaard. 2006. Cross-sectional prevalence of helminth infections in cattle on traditional, small-scale and large-scale dairy farms in Iringa District, Tanzania. *Veterinary Research Communications* **30**:45-55.
17. Martin, W. S., L. A. Meek., and P. Willeberg. 1988. *Veterinary Epidemiology*. 1st ed. Ontario: Ontario Veterinary College, 33 p.
18. Mulugeta, S., F. Begna., and E. Tesgaye. 2011. Prevalence of Bovine Fasciolosis and its Economic Significance in and Around Assela, Ethiopia. *Global Journal of Medical Research* **11** (3):1-7.
19. Nega, M., B. Bogale., M. Chanie., A. Melaku., and T. Fentahun. 2012. Comparison of Coprological and Postmortem Examinations Techniques for the

- Determination of Prevalence and Economic Significance of Bovine Fasciolosis. *Journal Advanced Veterinary Research* **2**:18-23.
20. Phiri, A. M., I. K. Phiri., and J. Monrad. 2006. Prevalence of amphistomiasis and its association with *Fasciola gigantica* infections in Zambian cattle from communal grazing areas. *Journal of Helminthology* **80**:65-73.
 21. Rapsch, C., G. Schweizer., F. Grimm., L. Kohler., C. Bauer., P. Deplazes., U. Braun., and P. R. Torgerson. 2006. Estimating the true prevalence of *Fasciola hepatica* in Cattle slaughtered in Switzerland in the absence of an absolute diagnostic test. *International Journal of Parasitology* **36**:1153-1161.
 22. Regassa, A., T. Woldemariam., S. Demisie., N. Moje., D. Ayana., and F. Abunna. 2012. Bovine Fasciolosis: Coprological, Abattoir Survey and Financial loss Due to Liver Condemnation in Bishooftu Municipal Abattoir, Central Ethiopia. *European Journal of Biological Sciences* **4**:83-90.
 23. Saad, A. A. 2004. A survey of diseases prevalent in dairy cows in the White Nile and Gezira State, Sudan. M.V.Sc Thesis. University of Khartoum, 4 p.
 24. Shiferaw, S., A. Kumar., and K. Amssalu. 2009. Organs condemnation and economic loss at Mekelle Municipal Abattoir, Ethiopia. *Haryana Veterinarian* **48**:17-22.
 25. Smith, R. 1995. *Veterinary clinical pathology. A problem-oriented Approach*, 2nd ed. Florida: CRC Press-Boca Raton, 52 p.
 26. Sturat, J. A. 1998. The life cycle of *F. hepatica*. In Dalton J.P. (Ed), *Fasciolosis*. CBA International Publication Cambridge, p. 1.
 27. Tasawar, Z., U. Minir., C. Hayat., and M. H. Lashari. 2007. The prevalence of *Fasciola hepatica* in Goats around Multan. *Pakistan Veterinary Journal* **27**:5-7.
 28. Thrusfield, M. 2005. *Veterinary Epidemiology*. 3rd ed. UK: Black Well Science, 183p.
 29. Tolan, R.W. Jr. 2011. Fasciolosis Due to *Fasciola hepatica* and *Fasciola gigantica* Infection: An Update on this 'Neglected' Neglected Tropical Disease. *Laboratory Medicine* **42** (2):107-116.
 30. Yilma, J., A. Mesfin. 2000. Dry season bovine fasciolosis in North Western part of Ethiopia. *Revue de Médecine Vétérinaire* **6**:493-500.

Table 1 : Prevalence of fasciolosis in 307 cattle examined by postmortem examination at Elkadaro abattoir, Khartoum, Sudan

Disease	Frequency	Relative frequency (%)	Cumulative frequency (%)
Negative	210	68.4% (210/307)	68.4%
Positive	97	31.6% (97/307)	100%
Total	307		

Table 2 : Prevalence of fasciolosis in 307 cattle examined by coproscopic examination at Elkadaro abattoir, Khartoum, Sudan

Disease	Frequency	Relative frequency (%)	Cumulative frequency (%)
Negative	245	79.8% (245/307)	79.8%
Positive	62	20.2% (62/307)	100%
Total	307		

Table 3 : Multivariate analysis for the association between fasciolosis (examined by postmortem) and potential risk factors using logistic regression

Risk Factors	No. tested	No. positive (%)	Exp (B)	95% C.I. for Exp (B)	p-value
Age (years)					
<2	25	0	---	---	1.000
2-4	71	3 (4.2%)	1	---	.999
>4	211	94 (44.5%)	18.5	3.1, 21.7	0.000
Breed					
Local	128	3 (2.3%)	1	---	1.000
Cross	34	1 (2.9%)	1.3	0.4, 3.2	0.641
Foreign	145	93 (64.1%)	77.6	9.3, 81.6	0.000
Weight					
Light	137	65 (47.4%)	3.0	1.9, 5.3	0.000
Medium	139	32 (23.0%)	1	---	1.000
Heavy	31	0	---	---	1.000
Source					
Nyala	79	2 (2.5%)	1	---	1.000
Ethiopia	142	93 (65.5%)	76.1	8.4, 83.7	0.000
Khartoum	35	0	---	---	1.000
Kordofan	13	0	---	---	0.999
Eljazeera	1	1 (100%)	---	---	1.000
White Nile	9	0	---	---	0.999
Kassala	25	1 (4 %)	1.6	1.1, 3.1	0.027
South Sudan	3	0	---	---	1.000
Animal Size					
Small	136	65 (47.8%)	3.1	1.9, 5.3	0.000
Medium	141	32 (22.7%)	1	---	1.000
Large	30	0	---	---	1.000

Table 4 : Multivariate analysis for the association between fasciolosis (examined by coprology) and potential risk factors using the logistic regression

Risk Factors	No. tested	No. positive (%)	Exp (B)	95% C.I. for Exp (B)	p-value
Age (years)					
<2	25	0	---	---	1.000
2-4	71	1 (1.4 %)	1	---	1.000
>4	209	61 (28.9%)	28.8	3.9,34.2	0.000
Breed					
Local	128	1 (0.7%)	1	---	1.000
Cross	34	0	---	---	1.000
Foreign	145	61 (42.1%)	94.4	13.2,102.6	0.000
Weight					
Light	137	60 (43.8%)	53.4	7.1,61.3	0.000
Medium	139	2 (1.4%)	1	---	1.000
Heavy	31	0	---	---	1.000
Source					
Nyala	79	1 (1.2%)	1	---	1.000
Ethiopia	142	61 (43.0%)	60.2	8.3,70.1	0.000
Khartoum	35	0	---	---	1.000
Kordofan	13	0	---	---	1.000
Eljazeera	1	0	---	---	1.000
White Nile	9	0	---	---	1.000
Kassala	25	0	---	---	1.000
South Sudan	3	0	---	---	1.000
Animal Size					
Small	136	60 (44.1%)	54.9	6.3,63.8	0.000
Medium	141	2 (1.4%)	1	---	1.000
Large	30	0	---	---	1.000

Table 5 : The difference in fasciolosis prevalence estimated based on coprology and postmortem examination at Elkadaro abattoir during May -July 2011

Test Type	No. Positive	Prevalence	Diff. (95% CI)	X ²	p-value
Coprology	62	20.2%	7.3 (3.1, 9.4)	1.669	0.000
Postmortem	97	31.6%			

Diff = difference

Table 6 : The presence or absence of *Fasciola spp* eggs in the feces of cattle with and without *Fasciola* in the liver for measuring the sensitivity, specificity and agreement level of the two tests

Faecal examination	Presence of <i>Fasciola spp</i> in liver		Total
	Fluke (+ve)	Fluke (-ve)	
Egg present (+ve)	62	0	62
Egg absent (-ve)	35	210	245
Total	97	210	307

This page is intentionally left blank