

GLOBAL JOURNAL

OF MEDICAL RESEARCH: A

Neurology & Nervous System

Neural Networks With Rule

Treatment for Neuropsychiatric

Highlights

Severe Traumatic Brain Injury

Health Status of Multiple Sclerosis

Discovering Thoughts, Inventing Future

VOLUME 25 ISSUE 1 VERSION 1.0



GLOBAL JOURNAL OF MEDICAL RESEARCH: A
NEUROLOGY AND NERVOUS SYSTEM

GLOBAL JOURNAL OF MEDICAL RESEARCH: A
NEUROLOGY AND NERVOUS SYSTEM

VOLUME 25 ISSUE 1 (VER. 1.0)

OPEN ASSOCIATION OF RESEARCH SOCIETY

© Global Journal of Medical Research. 2025.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website <http://globaljournals.us/terms-and-condition/menu-id-1463/>

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)
Sponsors: [Open Association of Research Society](#)
[Open Scientific Standards](#)

Publisher's Headquarters office

Global Journals® Headquarters
945th Concord Streets,
Framingham Massachusetts Pin: 01701,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals Pvt Ltd
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at local@globaljournals.org

eContacts

Press Inquiries: press@globaljournals.org
Investor Inquiries: investors@globaljournals.org
Technical Support: technology@globaljournals.org
Media & Releases: media@globaljournals.org

Pricing (Excluding Air Parcel Charges):

Yearly Subscription (Personal & Institutional)
250 USD (B/W) & 350 USD (Color)

EDITORIAL BOARD

GLOBAL JOURNAL OF MEDICAL RESEARCH

Dr. Apostolos Ch. Zarros

DM, Degree (Ptychio) holder in Medicine,
National and Kapodistrian University of Athens
MRes, Master of Research in Molecular Functions in
Disease, University of Glasgow FRNS, Fellow, Royal
Numismatic Society Member, European Society for
Neurochemistry Member, Royal Institute of Philosophy
Scotland, United Kingdom

Dr. Alfio Ferlito

Professor Department of Surgical Sciences
University of Udine School of Medicine, Italy

Dr. Jixin Zhong

Department of Medicine, Affiliated Hospital of
Guangdong Medical College, Zhanjiang, China, Davis
Heart and Lung Research Institute, The Ohio State
University, Columbus, OH 43210, US

Rama Rao Ganga

MBBS
MS (University of Health Sciences, Vijayawada, India)
MRCS (Royal College of Surgeons of Edinburgh, UK)
United States

Dr. Izzet Yavuz

MSc, Ph.D., D Ped Dent.
Associate Professor, Pediatric Dentistry Faculty of
Dentistry, University of Dicle Diyarbakir, Turkey

Sanguansak Rerksuppaphol

Department of Pediatrics Faculty of Medicine
Srinakharinwirot University
NakornNayok, Thailand

Dr. William Chi-shing Cho

Ph.D.,
Department of Clinical Oncology
Queen Elizabeth Hospital
Hong Kong

Dr. Michael Wink

Ph.D., Technical University Braunschweig, Germany
Head of Department Institute of Pharmacy and Molecular
Biotechnology, Heidelberg University, Germany

Dr. Pejcic Ana

Assistant Medical Faculty Department of Periodontology
and Oral Medicine University of Nis, Serbia

Dr. Ivandro Soares Monteiro

M.Sc., Ph.D. in Psychology Clinic, Professor University of
Minho, Portugal

Dr. Sanjay Dixit, M.D.

Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine
Web: pennmedicine.org/wagform/MainPage.aspx?

Antonio Simone Laganà

M.D. Unit of Gynecology and Obstetrics
Department of Human Pathology in Adulthood and
Childhood “G. Barresi” University of Messina, Italy

Dr. Han-Xiang Deng

MD., Ph.D
Associate Professor and Research Department
Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical
Neurosciences
Northwestern University Feinberg School of Medicine
Web: neurology.northwestern.edu/faculty/deng.html

Dr. Roberto Sanchez

Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University
Web: mountsinai.org/

Dr. Feng Feng

Boston University
Microbiology
72 East Concord Street R702
Duke University
United States of America

Dr. Hrushikesh Aphale

MDS- Orthodontics and Dentofacial Orthopedics.
Fellow- World Federation of Orthodontist, USA.

Gaurav Singhal

Master of Tropical Veterinary Sciences, currently
pursuing Ph.D in Medicine

Dr. Pina C. Sanelli

Associate Professor of Radiology
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,
School of Medicine and Biomedical Sciences
Web: weillcornell.org/pinasanelli/

Dr. Michael R. Rudnick

M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine
Web: uphs.upenn.edu/

Dr. Seung-Yup Ku

M.D., Ph.D., Seoul National University Medical College,
Seoul, Korea Department of Obstetrics and Gynecology
Seoul National University Hospital, Seoul, Korea

Santhosh Kumar

Reader, Department of Periodontology,
Manipal University, Manipal

Dr. Aarti Garg

Bachelor of Dental Surgery (B.D.S.) M.D.S. in Pedodontics
and Preventive Dentistry Pursuing Phd in Dentistry

Sabreena Safuan

Ph.D (Pathology) MSc (Molecular Pathology and Toxicology) BSc (Biomedicine)

Getahun Asebe

Veterinary medicine, Infectious diseases, Veterinary Public health, Animal Science

Dr. Suraj Agarwal

Bachelor of dental Surgery Master of dental Surgery in Oromaxillofacial Radiology.
Diploma in Forensic Science & Oodntology

Osama Alali

PhD in Orthodontics, Department of Orthodontics, School of Dentistry, University of Damascus. Damascus, Syria. 2013 Masters Degree in Orthodontics.

Prabudh Goel

MCh (Pediatric Surgery, Gold Medalist), FISPU, FICS-IS

Raouf Hajji

MD, Specialty Assistant Professor in Internal Medicine

Surekha Damineni

Ph.D with Post Doctoral in Cancer Genetics

Arundhati Biswas

MBBS, MS (General Surgery), FCPS, MCh, DNB (Neurosurgery)

Rui Pedro Pereira de Almeida

Ph.D Student in Health Sciences program, MSc in Quality Management in Healthcare Facilities

Dr. Sunanda Sharma

B.V.Sc.& AH, M.V.Sc (Animal Reproduction, Obstetrics & gynaecology),
Ph.D.(Animal Reproduction, Obstetrics & gynaecology)

Shahanawaz SD

Master of Physiotherapy in Neurology PhD- Pursuing in Neuro Physiotherapy Master of Physiotherapy in Hospital Management

Dr. Shabana Naz Shah

PhD. in Pharmaceutical Chemistry

Vaishnavi V.K Vedam

Master of dental surgery oral pathology

Tariq Aziz

PhD Biotechnology in Progress

CONTENTS OF THE ISSUE

- i. Copyright Notice
- ii. Editorial Board Members
- iii. Chief Author and Dean
- iv. Contents of the Issue
- 1. Evaluating Psilocybin as a Treatment for Neuropsychiatric Symptoms in Parkinson's Disease. **1-7**
- 2. Artificial Intelligence and Neural Networks with Rules based Filter are useful to Find a Correlations, between Diet and the Health Status of Multiple Sclerosis Patients, by Treating the Problem as a Surface Detector, One Can also Draw Indications on the Search for Oil and Recognize Radioactive Atoms on Neutral Surfaces. **9-16**
- 3. Toward a Neurocounseling Paradigm: The Science Behind Step One in Addiction Recovery. **17-34**

- v. Fellows
- vi. Auxiliary Memberships
- vii. Preferred Author Guidelines
- viii. Index



GLOBAL JOURNAL OF MEDICAL RESEARCH: A NEUROLOGY & NERVOUS SYSTEM

Volume 25 Issue 1 Version 1.0 Year 2025

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Evaluating Psilocybin as a Treatment for Neuropsychiatric Symptoms in Parkinson's Disease

By Nayiri Barton

Abstract- Parkinson's Disease (PD) is a progressive neurodegenerative disorder marked by motor symptoms due to dopaminergic degeneration and non-motor symptoms such as depression, anxiety, and cognitive impairment, which significantly affect patients' quality of life. Traditional dopaminergic therapies address motor symptoms but offer limited efficacy for neuropsychiatric manifestations. Psilocybin, a serotonergic compound with strong affinity for the 5-HT2A receptor, has emerged as a promising candidate for addressing the complex symptomatology of PD, including its neuropsychiatric components. This review examines the pharmacological effects of psilocybin, particularly its ability to modulate serotonergic and dopaminergic systems, enhance neuroplasticity, and reduce neuroinflammation, offering a potential therapeutic approach for PD. While clinical research in PD remains limited, evidence from related conditions such as Major Depressive Disorder (MDD) and Substance Use Disorder (SUD) supports the notion that psilocybin could modulate both motor and non-motor symptoms in PD.

GJMR-A Classification: NLMC: WL 359



Strictly as per the compliance and regulations of:



Evaluating Psilocybin as a Treatment for Neuropsychiatric Symptoms in Parkinson's Disease

Nayiri Barton

Abstract- Parkinson's Disease (PD) is a progressive neurodegenerative disorder marked by motor symptoms due to dopaminergic degeneration and non-motor symptoms such as depression, anxiety, and cognitive impairment, which significantly affect patients' quality of life. Traditional dopaminergic therapies address motor symptoms but offer limited efficacy for neuropsychiatric manifestations. Psilocybin, a serotonergic compound with strong affinity for the 5-HT2A receptor, has emerged as a promising candidate for addressing the complex symptomatology of PD, including its neuropsychiatric components. This review examines the pharmacological effects of psilocybin, particularly its ability to modulate serotonergic and dopaminergic systems, enhance neuroplasticity, and reduce neuroinflammation, offering a potential therapeutic approach for PD. While clinical research in PD remains limited, evidence from related conditions such as Major Depressive Disorder (MDD) and Substance Use Disorder (SUD) supports the notion that psilocybin could modulate both motor and non-motor symptoms in PD. Furthermore, psilocybin's ability to induce brain network hyperconnectivity and regulate dopamine release offers mechanistic insight into its potential benefits. Despite the promising neurobiological underpinnings, ethical concerns and regulatory constraints remain barriers to widespread clinical use. Future research should prioritize disease-specific trials to explore psilocybin's therapeutic efficacy, optimal dosing, and safety profile in PD, potentially redefining the treatment landscape for this underserved population.

I. INTRODUCTION

Parkinson's disease (PD) is a progressive neurodegenerative disorder primarily characterized by both motor and debilitating non-motor symptoms, including depression, anxiety, and cognitive decline. These symptoms often appear before the onset of motor deficits and contribute significantly to reduced quality of life. While dopaminergic therapies such as levodopa remain the cornerstone of motor symptom management, they offer limited benefit for neuropsychiatric symptoms and may lead to complications with prolonged use. As the limitations of conventional therapies become more evident, interest has grown in exploring alternative approaches such as psychedelic-assisted therapy. This narrative review examines the current evidence surrounding psilocybin's neurobiological mechanisms and therapeutic potential

in the context of Parkinson's disease. Psilocybin, a serotonergic compound derived from certain species of mushrooms, has shown promising effects in treating mood disorders and substance use disorders by acting on the 5-HT2A receptor. PD, MDD, and SUD all involve underlying dopaminergic dysfunction, suggesting a potential mechanistic overlap. Psilocybin's ability to modulate serotonergic activity and indirectly influence dopaminergic pathways may offer a novel strategy for addressing the complex symptom profile of PD.

II. PSILOCYBIN: PHARMACOLOGICAL OVERVIEW

Psilocybin is a naturally occurring psychedelic compound found in several species of mushrooms, colloquially referred to as "magic mushrooms." After ingestion, psilocybin is rapidly converted in the body to its active form, psilocin, which acts primarily as a serotonin receptor agonist, with high affinity for the 5-HT2A receptor*. Activation of this receptor is thought to underlie the drug's characteristic effects on perception, mood, and cognition. Psilocybin's influence on neural plasticity, including increased dendritic growth and synaptogenesis, has drawn significant attention for its potential in treating psychiatric and neurodegenerative disorders. These structural changes are believed to be mediated in part through downstream signaling cascades involving brain-derived neurotrophic factor (BDNF) and the mTOR pathway, both of which play crucial roles in synaptic remodeling and neuronal resilience [2].

Pharmacokinetically, psilocin typically reaches peak plasma concentration within 1 to 2 hours after oral administration, with a terminal half-life of approximately 2 to 3 hours. Despite its relatively short half-life, the subjective effects can last 4 to 6 hours, aligning well with the temporal needs of a therapeutic session. This brief but intense window of altered consciousness allows clinicians to structure therapy sessions with predictable onset and offset, reducing risks associated with prolonged intoxication. The short duration of action, combined with the low potential for physiological

Author: e-mail: nayiribarton@gmail.com
ORCID: 0009-0004-2110-3495

* (although mechanisms of the 5-HT1A and 5-HT2C receptors play roles in altering perception, 5-HT2A's potential will be primarily highlighted within this review)*



dependence and a favorable safety profile in controlled settings, supports the use of psilocybin in supervised clinical environments [3].

In a double-blind study on the effects of psilocybin, participants found their experience “[had] substantial personal meaning and spiritual significance and attributed to the experience sustained positive changes in attitudes and behavior consistent with changes rated by community observers” [4]. Its ability to induce meaningful psychological experiences, when administered with appropriate therapeutic support, has made it a leading candidate in the emerging field of psychedelic-assisted therapy.

III. NEUROBIOLOGICAL BASIS

a) Pathophysiology of Parkinson's Disease

Parkinson's Disease is characterized by the progressive degeneration of dopaminergic neurons in the substantia nigra, leading to a significant reduction in dopamine levels in the striatum. This dopaminergic cell loss is the hallmark of PD, which disrupts the basal ganglia's motor control pathways, resulting in motor symptoms such as bradykinesia, tremors, and rigidity. PD, however, “not only affects dopamine systems but also includes alterations in serotonergic, cholinergic, and other neurotransmitter systems, exacerbating symptoms such as depression and cognitive dysfunction,” [5]. This broader involvement of neurotransmitter systems contributes to a diverse range of both motor and non-motor symptoms, including mood disorders, sleep disturbances, and cognitive decline, all of which complicate the management of PD. Moreover, the progressive nature of the disease is compounded by mechanisms like neuroinflammation, mitochondrial dysfunction, and oxidative stress, [6]. Additionally, the aggregation of, “misfolded alpha-synuclein proteins plays a central role in the progression of Parkinson's disease, leading to neuronal death and dysfunction,” [7]. These multifaceted molecular processes: dopaminergic degeneration, altered neurotransmitter systems, neuroinflammation, mitochondrial dysfunction, and alpha-synuclein aggregation, collectively contribute to the complex pathophysiology of Parkinson's disease, making it a challenging disorder to treat and manage effectively.

b) 5-HT2A Receptor in Psilocybin: Mechanism of Action

Parkinson's disease involves not only dopaminergic neuron degeneration in the substantia nigra but also early and significant changes in the serotonergic system. Degeneration of serotonergic neurons in the *raphe nuclei*, a primary source of brain serotonin, occurs early in the disease and contributes to widespread serotonin deficits. As noted, “serotonergic dysfunction is a prominent feature of Parkinson's disease, contributing significantly to both motor and nonmotor symptoms” [6]. This dysfunction underlies

symptoms such as depression, anxiety, and cognitive decline. Altered serotonin receptor density and signaling, including changes in 5-HT2A receptor expression, have been observed in cortical and subcortical regions affected by PD, highlighting the critical role of serotonergic-dopaminergic interactions in the disease's pathophysiology and therapeutic approaches.


The 5-HT2A Receptor, commonly referred to as the serotonin-2A receptor, is a G-protein-coupled-receptor (GPCR) activated by the neurotransmitter serotonin (5-hydroxytryptamine). Psilocybin, a serotonergic psychedelic, exerts its primary pharmacological effects through partial agonism at the 5-HT2A receptor, densely expressed in cortical and subcortical brain regions. Activation of this receptor initiates a cascade of intracellular events, including phospholipase C stimulation and subsequent increases in intracellular calcium levels, which influence gene transcription and synaptic plasticity [7]. These signaling pathways are implicated in enhanced cortical connectivity and neuroplasticity, mechanisms increasingly viewed as therapeutically valuable in neurodegenerative diseases. Psilocybin-induced 5-HT2A receptor activation has also been shown to indirectly affect dopaminergic systems, particularly within the mesocorticolimbic and nigrostriatal pathways, suggesting a potential mechanism by which psilocybin could modulate motor and affective symptoms in Parkinson's disease, [7]. In this context, the serotonergic-dopaminergic interplay is especially relevant, as serotonin receptors may help normalize dysregulated dopamine transmission: a hallmark of PD pathophysiology. 5-HT2A activation can promote, “[mediate] Psychedelic-induced hyperconnectivity in the brain...[this] may facilitate neural rewiring, which could offer therapeutic benefits for Parkinson's disease,” [9]. Collectively, these findings underscore the 5-HT2A receptor as a key mediator of psilocybin's neurophysiological effects and a potential target in the development of novel interventions for Parkinson's disease.

IV. PSilocybin IN THE TREATMENT OF NEUROLOGICAL AND PSYCHIATRIC DISORDERS

Historically, Psilocybin has been actively used in treating various medical and mental health conditions. Specifically focused on Substance Abuse Disorder (SUD) or addiction issues, psilocybin (specifically the properties of 5-HT2A) has been successfully proven to help mitigate symptoms of withdrawal or craving, as well as improve neuroplasticity after the effects of substance abuse damages the brain. The “proposed working mechanisms of psychedelics, including psilocybin, for improving psychiatric symptoms such as depression

and SUD are both biological (e.g., by inducing brain neuroplasticity through elevating Brain-Derived Neurotrophic Factor (BDNF) levels, which are diminished in psychiatric conditions) and psychological...," [10]. Neuroplasticity is seen in patients of addiction and neurological damage significantly after the use of Psychedelic Administration Therapy (PAP) due to psychoactive properties. Psilocybin's psychoactive properties have demonstrated clinically meaningful antidepressant effects in controlled studies. Psilocybin's favorable safety profile, particularly in controlled therapeutic settings, supports its

investigational use for patients with MDD (Major Depressive Disorder). In a randomized psilocybin trial study conducted at Braxia Health in Ontario, Canada, patients (not participating in pharmacological treatment) were administered "one, two or three psilocybin sessions with a fixed dose of 25 mg synthetic psilocybin powder dissolved in water....accompanied by one preparatory therapy session (1–2 h), a supportive dosing session (6–8 h) and two integration therapy sessions (1–2 h each)...," [12]. After observing longer term effects of overall mental health, the following results can be viewed below:



Based on the mean standard errors for both depression severity and suicidality, the trial demonstrated significant reductions from baseline levels, highlighting the potential of Psilocybin therapy as a promising intervention for psychological well-being.

V. CLINICAL PARALLELS: WHY SUCCESS IN SUD AND MDD MATTERS FOR PD

Though there has not been extensive direct research on the specific effects of psilocybin in Parkinson's disease (PD), neurobiological, mechanistic, and clinical parallels provide a biologically plausible rationale for its therapeutic exploration. Psilocybin promotes neuroplasticity by increasing dendritic spine growth and enhancing synaptic connectivity, primarily through activation of serotonin receptors such as 5-

[11] Psilocybin with psychological support for treatment-resistant depression: An open-label feasibility study.


HT2A, which leads to elevated brain-derived neurotrophic factor (BDNF) levels. These neuroplastic changes have demonstrated benefits in conditions marked by impaired plasticity and neurodegeneration, including major depressive disorder (MDD) and PD [14].

Importantly, affective symptoms such as apathy and anhedonia are shared across MDD, substance use disorder (SUD), and PD, although their clinical presentations and underlying neurocircuitry may differ. In PD, apathy often correlates with dopaminergic degeneration in nigrostriatal pathways, while in MDD and SUD, disruptions in frontostriatal and limbic circuits contribute to these symptoms. Despite these differences, all three conditions involve dysregulated reward processing and motivation pathways modulated by serotonergic and dopaminergic signaling. As reviewed, "apathy and anhedonia manifest across these

disorders through overlapping but distinct neural mechanisms, highlighting transdiagnostic relevance," [13].

Regarding BDNF, its function diverges between neurodegenerative and mood disorders. In mood disorders like MDD, reduced BDNF expression is linked primarily to impaired synaptic plasticity and neuronal resilience, contributing to mood dysregulation. In contrast, in PD, BDNF deficiency additionally exacerbates dopaminergic neuron loss and neurodegeneration. Psilocybin's ability to elevate BDNF may thus serve a dual role: restoring synaptic plasticity in mood disorders while potentially promoting neuronal survival in neurodegenerative disease [13].

Specific mechanism depiction can be seen below:



[12] Augmentation in depression: Esketamine, a new standard?

On a psychological level, a multicenter trial led by Dr. James Rucker of King's College London, found that a 25 mg dose of psilocybin on human participants, alongside psychological support, significantly impacted depression symptoms in participants with treatment-resistant depression. The study observed greater reductions in depression scores three weeks after treatment in the high-dose group compared to the placebo group [16].

VI. CURRENT PAP AND PD CONDUCTED RESEARCH

While direct research on psilocybin's effects in Parkinson's Disease remains limited, the existing neurobiological, mechanistic, and clinical evidence suggests promising potential for its application in this population. The observed ability of psilocybin to enhance neuroplasticity, modulate inflammatory pathways, and alleviate depressive symptoms in related conditions like MDD and SUD highlights its potential as a multifaceted therapeutic approach for the non-motor symptoms of Parkinson's. A pilot study at UC San

Francisco, led by Ellen R. Bradley, investigated the effects of psilocybin on mood, cognition, and motor function in individuals with Parkinson's disease. Participants experienced clinically significant improvements in mood and motor symptoms that lasted for weeks after treatment. The study suggests that psilocybin may help repair brain function and alleviate non-motor symptoms associated with PD. After administering psilocybin to a moderately sized group of participants with PD and observing results, "non-motor (MDS-UPDRS Part I: -13.8 ± 1.3 , $p < 0.001$, Hedges' $g = 3.0$) and motor symptoms (Part II: -7.5 ± 0.9 , $p < 0.001$, $g = 1.2$; Part III: -4.6 ± 1.3 , $p = 0.001$; $g = 0.3$) as well as performance in select cognitive domains (Paired Associates Learning $[-0.44 \pm 0.14$, $p = .003$, $g = 0.4$], Spatial Working Memory $[-0.52 \pm 0.17$, $p = 0.003$, $g = 0.7$], and Probabilistic Reversal Learning $[2.9 \pm 0.9$, $p = 0.003$, $g = 1.3$]) improved post-treatment," supporting the plausibility of extending findings from related conditions to PD [8]. Additionally, on a preliminary case-report done on a middle-aged woman (PD patient) with a lack of depressive-symptoms, positive effects of PAP were seen. Post-treatment, the

patient reflected on a more positive outlook on the severity of her disorder, worrying less about mortality or her future, "promoting profound decentration from habitual thoughts and emotions, improving mood and PD acceptance," [8]. Though these observed effects could be attributed to potential placebo, the existence of general outlook improvement is not one to overlook, and is often a pivotal tool for the wellbeing of those with chronic neurodegenerative disorders like PD.

VII. NEUROIMAGING AND MECHANISTIC EVIDENCE

Psilocybin has been shown to enhance brain network connectivity, particularly within the default mode network (DMN), as demonstrated by fMRI studies. These studies reveal that psilocybin induces hyperconnectivity across multiple brain regions, including the prefrontal cortex, which is associated with subjective alterations in cognition and perception, [8]. This heightened connectivity may be particularly beneficial for PD, where reduced brain connectivity contributes to both motor and non-motor symptoms, [15]. These findings support the idea that psilocybin's neuroplastic effects may provide relief from symptoms often associated with neurodegenerative conditions.

VIII. DOPAMINE-SEROTONIN SYSTEM INTERACTIONS

Psilocybin's effects are mediated primarily through serotonin 5-HT2A receptor activation, which in turn influences dopamine release, particularly in regions involved in motor control. This interaction is crucial for Parkinson's Disease, where dopamine dysregulation plays a central role in both motor and non-motor symptoms. It's highlighted that psilocybin, "primarily activates serotonin 5-HT2A receptors, which modulate dopamine release, potentially alleviating symptoms related to PD" [12]. The regulation of both serotonin and dopamine systems could improve symptoms in PD patients, especially those who do not respond to conventional dopamine-based therapies, with "serotonin-dopamine interactions in Parkinson's Disease [being critical] for understanding treatment efficacy, with psilocybin potentially [modulates] both neurotransmitter systems to improve motor and non-motor symptoms" [14].

IX. SAFETY, ETHICS, AND THERAPEUTIC CONSIDERATIONS

As interest in psilocybin as a therapeutic tool for neurodegenerative disorders like PD grows, safety and ethical considerations must be carefully addressed. Psilocybin, like other psychedelics, carries potential risks, particularly when used outside of controlled environments or by individuals with certain health

conditions. Common side effects include transient anxiety, confusion, and disorientation, which may be particularly concerning for individuals with cognitive impairments, such as those with PD. It's not to be forgotten that psilocybin possesses psychoactive properties, with potential for hallucinogenic and psychedelic symptoms if taken at a high or non-controlled dose. However, studies have generally found that psilocybin is well-tolerated when administered in a supervised clinical setting, with careful screening to ensure patient suitability.

Additionally, the issue of cognitive decline, neural plasticity damage, and deficit as symptoms of PD become present in the ethical issue of consent by participants. Depending on the stage/onset of a participant's PD, it can potentially be a violation of conscious consent to administer a substance like psilocybin considering the patient may not be at the same level of consciousness as pre-diagnosis. In terms of decision-making, based on the pathophysiology of the disorder, "a progressive loss of cells in the midbrain producing the neurotransmitter dopamine...the controlled release of dopamine into synapses connecting the cerebral cortex and the basal ganglia plays a central role during the selection and reinforcement of actions," [10]. Due to potential decision-making impairment, the clinician in charge of administering a patient with said psychedelic must be perceptive on the patient's level of cognitive decline and ability to make a conscious decision in order to avoid ethical or HIPAA violations. Once these factors are taken into account, however, PAP appears to be an effective and low risk strategy towards cognitive and emotional recovery/stability.

X. LIMITATIONS IN CURRENT RESEARCH

Psilocybin has emerged as a leading candidate in contemporary psychedelic research. Considering this, there are still major limitations on clinical exploration/trials for neurodegenerative disorders like PD. As mentioned above, the cognitive decline and ambiguous or impaired decision-making capacity in cognitively affected individuals may raise ethical concerns regarding informed consent. Due to the severity of PD's neurodegenerative impacts, sample sizes and lengths of trials are significantly shorter than desired, therefore altering the professionalism and accuracy of results. Short trial durations and limited sample sizes restrict statistical power, increase variability, and limit the generalizability of findings. Additionally, the long-term safety profile of repeated or high-dose psilocybin administration remains insufficiently characterized, particularly for populations already vulnerable to cognitive decline or psychosis. Regulatory constraints and the classification of psilocybin as a Schedule I substance also pose



significant barriers to large-scale, multi-site studies. Until these limitations are addressed through more rigorous, targeted, and longitudinal trials (including those specific to PD), clinical recommendations for psilocybin use in neurodegenerative contexts must remain cautious and exploratory.

XI. FUTURE DIRECTIONS AND RESEARCH PRIORITIES

To fully assess psilocybin's therapeutic potential for Parkinson's Disease, future research must prioritize disease-specific clinical trials that evaluate both motor and non-motor outcomes. These studies should include larger, more diverse patient populations, with extended follow-up periods to better understand long term efficacy and safety. There is also a pressing need for trials that examine optimal dosing regimens, frequency of administration, and the role of psychological support during psychedelic-assisted therapy. Particularly, the effects of repeated dosing on serotonergic neurotoxicity, emotional lability, or precipitating latent psychosis in predisposed individuals remain poorly understood. Mechanistic studies using advanced neuroimaging and biomarker analysis, particularly those exploring serotonin-dopamine interactions and network level brain changes, will be critical in determining how psilocybin affects PD specific neurobiology. In addition, comparative trials that evaluate psilocybin against existing pharmacological and behavioral therapies could help clarify its position within the broader treatment landscape. As regulatory frameworks continue to evolve, ethical and equitable access to psychedelic therapy must remain a central concern, ensuring that any future therapeutic model is both scientifically grounded and socially responsible.

XII. CONCLUSION

Psilocybin emerges as a promising candidate with systematically supported potential for treating neuropsychiatric symptoms in PD. Its mechanism of action, centered on 5-HT2A receptor activation, initiates a cascade of neuroplastic and neuromodulatory processes that extend beyond mood regulation to potentially influence dopaminergic function, cortical network connectivity, and inflammatory pathways. While direct research on psilocybin in PD remains limited, strong clinical and mechanistic parallels from studies in depression and substance use disorders offer a valuable foundation for future inquiry. Although limited, early pilot data suggests that psilocybin may positively affect both motor and non motor symptoms, warranting further investigation. However, ethical, regulatory, and methodological challenges must be addressed through rigorous, disease-specific clinical trials. In particular, the use of psilocybin in vulnerable populations such as those with PD necessitates robust consent frameworks,

clearly defined inclusion criteria, and stringent monitoring protocols to ensure patient safety and uphold ethical standards. Further research should prioritize Phase I safety studies in early-stage PD, with a focus on affective symptoms, quality of life metrics, and neurobiological markers of change. With appropriate safeguards and continued interdisciplinary research, psilocybin could help redefine the therapeutic landscape for Parkinson's Disease, offering a novel avenue for enhancing quality of life in a population long underserved by conventional treatments.

REFERENCES RÉFÉRENCES REFERENCIAS

1. American Psychological Association. Press release: Parkinson disease. Published March 2010. Accessed May 17, 2025. <https://www.apa.org/news/press-releases/2010/03/parkinson-disease>
2. Ly, C., Greb, A. C., Cameron, L. P., Wong, J. M., Barragan, E. V., Wilson, P. C., ... & Olson, D. E. (2018). *Psychedelics promote structural and functional neural plasticity*. *Cell Reports*, 23(11), 3170–3182. DOI: 10.1016/j.celrep.2018.05.022
3. Holze, F., Becker, A. M., Kolaczynska, K. E., Duthaler, U., & Liechti, M. E. (2023). Pharmacokinetics and Pharmacodynamics of Oral Psilocybin Administration in Healthy Participants. *Clinical pharmacology and therapeutics*, 113(4), 822–831. <https://doi.org/10.1002/cpt.2821>
4. Griffiths, R. R., Richards, W. A., McCann, U., & Jesse, R. (2006). Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance. *Psychopharmacology*, 187(3), 268–292. <https://doi.org/10.1007/s00213-006-0457-5>
5. Mastinu A, Premoli M, Bonini SA, et al. Psilocybin for treating psychiatric disorders: Pharmacological profile and clinical perspectives. *Neurosci Biobehav Rev*. 2023; 149:105320. doi:10.1016/j.neubiorev.2023.105320
6. Politis, M., Wu, K., Loane, C., et al. (2010). Serotonergic dysfunction in Parkinson's disease and its relevance to nonmotor symptoms. *Proceedings of the National Academy of Sciences*, 107(37), 16087-16092. <https://doi.org/10.1073/pnas.1007378107>
7. Fleury P, Goff DC, Murray RM. Psychedelic compounds as treatment strategies for neuropsychiatric symptoms in Parkinson's disease: Rationale and emerging evidence. *Neuropharmacology*. 2025; 221:109444. doi:10.1016/j.neuropharm.2024.109444
8. Bradley PC, Holtman JR, Haroutounian S. Psychedelic-assisted therapy for neurodegenerative diseases: Challenges and promises. *J Neural Transm*. 2025; 132(1):87–95.
9. Elsevier. Psychedelic drug-induced hyperconnectivity in the brain helps clarify altered subjective experiences. Elsevier Press Release. Published April 2024. <https://www.elsevier.com/about/press-releases/psychedelic-drug-induced-hyperconnectivity-in-the-brain>
10. Brengel D, Keller L, Möller HJ. Alpha-synuclein pathology and implications for neurodegeneration. *Parkinsonism Relat Disord*. 2025;108:56-64.
11. Carhart-Harris RL, Bolstridge M, Rucker J, et al. Psilocybin with psychological support for treatment-resistant depression: An open-label feasibility study. *Lancet Psychiatry*. 2016;3(7):619-627. doi:10.1016/S2215-0366(16)30065-7
12. Rosenblat JD, McIntyre RS. Augmentation in depression: Esketamine, a new standard? *Med (N Y)*. 2023;4(12):852-854. doi:10.1016/j.medj.2023.11.002
13. Husain, M., & Roiser, J. P. (2018). *Neuroscience of apathy and anhedonia: A transdiagnostic approach*. *Nature Reviews Neuroscience*, 19(8), 470–484. <https://doi.org/10.1038/s41583-018-0029-9>
14. Fereydouni S, Andrade C, Osuch EA. Serotonergic psychedelics and dopamine: An interaction with therapeutic implications ons. *Biol Psychiatry*. 2025; 97(4):251-25
15. De Deus JM, Polari A, Maia FM. Neuroinflammatory and plasticity pathways affected by psychedelics: Potential relevance to Parkinson's disease. *Brain Behav Immun Health*. 2025; 32:100813.
16. Herz DM, Eickhoff SB, Løkkegaard A, Siebner HR. Functional neuroimaging of motor control in Parkinson's disease: A meta-analysis. *Hum Brain Mapp*. 2017;38(7):3227-3247. doi:10.1002/hbm.23588



This page is intentionally left blank





GLOBAL JOURNAL OF MEDICAL RESEARCH: A NEUROLOGY & NERVOUS SYSTEM

Volume 25 Issue 1 Version 1.0 Year 2025

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Artificial Intelligence and Neural Networks with Rules based Filter are useful to Find a Correlations, between Diet and the Health Status of Multiple Sclerosis Patients, by Treating the Problem as a Surface Detector, One Can also Draw Indications on the Search for Oil and Recognize Radioactive Atoms on Neutral Surfaces

By Francesco Pia

Abstract- In this article we address three different problems of "Surface Analysis" thanks to Artificial Intelligence with Neural Networks related to the diet of the person newly affected by multiple sclerosis, a diet that can be myelinating or demyelinating; the second has the objective of recognizing whether oil, or water, can be present in a site and the presumed depth; the third objective is to classify radioactive atoms, such as Plutonium, on a neutral surface, as a sort of "Identity Card". The data were collected, not on sites or authors that deal with the topic in question, but thanks to what emerges in summary form in the most common browsers, called in this work "data found on the volatile web".

Keywords: *food, myelinating, demyelinating, radioactivity.*

GJMR-A Classification: NLMC: WL 360



Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

Artificial Intelligence and Neural Networks with Rules based Filter are useful to Find a Correlations, between Diet and the Health Status of Multiple Sclerosis Patients, by Treating the Problem as a Surface Detector, One Can also Draw Indications on the Search for Oil and Recognize Radioactive Atoms on Neutral Surfaces

Francesco Pia

Abstract- In this article we address three different problems of "Surface Analysis" thanks to Artificial Intelligence with Neural Networks related to the diet of the person newly affected by multiple sclerosis, a diet that can be myelinating or demyelinating; the second has the objective of recognizing whether oil, or water, can be present in a site and the presumed depth; the third objective is to classify radioactive atoms, such as Plutonium, on a neutral surface, as a sort of "Identity Card". The data were collected, not on sites or authors that deal with the topic in question, but thanks to what emerges in summary form in the most common browsers, called in this work "data found on the volatile web". It should be noted that the following chapters of this article do not aim to be precise and complete, but only to provide some ideas presented in a preliminary form to be explored in further studies. Each reader can draw the various indications, interpreting, thanks to the preliminary results, to modify the various "Intelligent" systems and proposing modifications to the various proposed architectures; This aspect must be considered carefully and recursively in all three chapters that follow. In a second stage, the bibliographical sources will also have to be studied in depth, finding help also from the various authors to be taken into consideration. remembering that the various problems, present in the three chapters, have been addressed by transforming them into a "surface" analysis.

Keywords: food, myelinating, demyelinating, radioactivity.

I. INTRODUCTION

Our time is precious, and we thank you for the one you will spend reading this. Since I was 7 years old I have always liked to study, and now I am the only author of the works present in the bibliography [1]-[14]. I suffer from nrsSPMS that blocks me and many patients, families, all over the world. I have opened a fundraiser on "www.gofundme.com" with the keyword "Artificial intelligence against multiple sclerosis" and thanking you, I promise that I will make every effort to ensure that the funds raised will be spent on the next works that will surely be expensive.

Author: prof. PhD Eng. Gonnosfanadiga (SU), Italy.
www.gofundme.com keyword "Artificial intelligence vs Multiple Sclerosis". e-mail: piafranc@hotmail.com

In this work three applications of a versatile "intelligent" system as a "surface analyzer" will be presented, in different chapters: the first on Multiple Sclerosis, the second as a detector of oil, water or other, the third for the identification of heavy and radioactive atoms.

II. DIET AND MULTIPLE SCLEROSIS

In this chapter we will try to adapt the intelligent systems present in [12], [14] in order to improve the analysis capabilities and to further test the "seed of discernment" present in [12] and [14] in "surface analysis" problems. and in particular, in this chapter, in the link between the diet assumed by a newly diagnosed multiple sclerosis patient and his state of well-being by virtue of myelinizing or demyelinizing foods.

In the following lines we will try to give some indications for the newly diagnosed or not that emerged from the experience of the undersigned that each reader will evaluate independently or followed by his neurologist.

1. Doing sports, fitness and any activity with your body
2. Stem cell implant, after medical opinion from different bell towers
3. Assumer TOLEMABRUTINIB® by SANOFI®, PIPE 307® Contineum Therapeutics, Inc. of hoping my neurologist agrees, I'm waiting anxiously.
4. Resorting to the intermittent diet.
5. Use, with the help of a nutritionist, mainly myelinizing foods.

In this chapter we address the problem of diet in "newly" diagnosed patients with multiple sclerosis, which can be myelinating or demyelinating depending on dietary intake, addressing it with artificial intelligence with an architecture based mainly on Back-Propagation neural networks. The inputs used are to be considered "pseudo cybernetic" but of natural origin that can be for or against the increase of myelin in circulation. Through the use of NNs connected with appropriate architectures, a system could be built that highlights the

reaction, in the medium-long term, of the human body to natural inputs available such as food; without machines, weight indicators, strength, and only thanks to the musculoskeletal response felt and self-assessed by the subject who adheres to this diet here simply called "diet".

The research of information, and data, was conducted thanks to the "volatile" WEB, a type of information that is not necessarily collected in specific sites, so there will be no bibliographic references nor authors and their works, only thanks to the various navigators that bring the data to light. It is very important to take into consideration the fact that these pages do not want to give precise, exhaustive indications on the eating habits of newly diagnosed people. In fact, some people may like some foods that are indicated as demyelinating, for example by eliminating them, and in that case they would have others, difficult to correlate with positive and demyelinating effects. Each patient, and not only, can freely interpret under his experience the indications that the system will provide him; as he could, on the contrary, feed himself with myelinating elements that give him more strength than others, not indicated or unfortunately given as demyelinating by the system; therefore, if he does not find anything to replace them, or integrate them, he will be able to interact with the system by leaving a feedback. The data used in this work are those that emerge from browser searches and displayed in summary form: this type of data is called, in this work, "volatile"; that is, high-level information, if they were low-level (such as saying that one would have the duty to eat... as in the bibliography would find...), with these data it will be a duty for the user, after entering them into the system and following the nutritionist's instructions.

So, whoever goes on a diet is considered a "natural" cybernetic system with input and programmed because the diet is still a nutritional reprogramming of the body whose outputs are deduced by the user who perceives a possible benefit or not..

Foods are either remyelinating or demyelinating. In this work, few elements are evaluated; in fact, it is a system that can be modified or expanded.

The "seed of discernment [12]" has been used in all three chapters: the use of two NNs, appropriately interconnected, which give a nuanced response, where the response is the motor improvement or not that the user can report. This scheme is present in [12] where the same type of approach is used, but in this work no laboratory tests will be performed, but reference will be made to the state of well-being reported by the user. The Sulfurin is a rare, myelinating substance that can repair myelin; the naturally occurring molecule blocks the activity of an enzyme that is overactive in areas of myelin damage. This same enzyme also contributes to the growth and spread of cancer cells, meaning the discovery has implications beyond MS. Study co-author

Angela Hoffman, a professor at the University of Portland who had been examining nearly unobtainable plant-based hyaluronidase inhibitors, and Steve Bryson and colleagues at Oregon Health & Science University say it's exciting that the plant-based molecule can inhibit the growth of myelin-producing cells [14].

Our body contains about 21 mg of Potassium 40, and emits β rays and neutrinos (cosmic objects) about 400 million/day; it could also repair itself, and implement remedies with the right stimulations. In this work we use foods in two groups as stimuli: remyelinating and demyelinating, the first ones help myelination the second ones damage it. Artificial Intelligence can be useful to verify if this reasoning is coherent. We do not want to question foods that are eaten daily by those who are not diagnosed with MS; such as cereals for many people, however, certain foods that the network indicates as such are considered myelinating. The research, in this work, is also sometimes superfluous since for the simulation it would be enough to insert generic foods, for example from the Mediterranean diet; dividing them into categories: this is important because the present system is made up of two Neural Networks that use the so-called "Seed of Discernment" [12]: the first work where MS is studied with AI.

Given the paradigm: move and you will move, stay still and you will stay still: let's assume the metabolization of the food taken in the diet. This paradigm can be used as an input, with any type of diet, drug, etc. etc. without exercise it is almost all useless; that is, if you take a "new drug" and do not follow adequate physiotherapy it is almost useless. Instead, considering physical activity as a myelinating factor and staying still as a demyelinating one, it can be used as an input, in a specific APP, metabolized by movement. It would not be wrong to imagine that since a myelinating food introduced, not metabolized does not go into circulation, but is only digested; but if metabolized it is more likely to have a myelinating effect.

a) A Linear Algebra for Surface Analysis

From the following table tab.1 it is possible to extract a square matrix to obtain a function that highlights the trend of the new data, compared to the average of the previous data, from which we consider the average. The matrix can be constructed as a square by joining the rows with low quantity impact (for example dietary) and adding columns that take into account the response of the various outputs (for example parts of the patient's body subjected to the diet) and other non-linearly dependent on the previous ones, useful for describing the state of the system (for example the patient's health).

Let's proceed by making the data in array form. Given the original array $\begin{bmatrix} \varrho \end{bmatrix}$, a random reallocation of its various cells is carried out, obtaining $\begin{bmatrix} \overrightarrow{R} \end{bmatrix}$ of which we would calculate the average and normalization.

I then subtract that of the new day of the same patient $\begin{bmatrix} \overline{G} \end{bmatrix}$:

$$\begin{bmatrix} \overline{R} \end{bmatrix} - \begin{bmatrix} \overline{G} \end{bmatrix}$$

($j+1 \rightarrow i$) for any days more than one; from which we can calculate:

$$(a) \begin{bmatrix} P \end{bmatrix} = \nabla \left(f \left(\text{div} \begin{bmatrix} \overline{R} \end{bmatrix} - \begin{bmatrix} \overline{G} \end{bmatrix} \right) \right) \begin{cases} > 0 & *? \\ < 0 & \end{cases}$$

After proper reallocation to the original cells of patient:

Gistificare la riallocazione, così che non vengano considerate le posizioni sempre presenti specificare meglio l'analisi di superficie


$$(b) \begin{bmatrix} \overleftarrow{P} \end{bmatrix}$$

$$(c) \begin{cases} > 0 & \text{perhaps it has a negative impact} \\ = 0 & \text{it's almost indifferent} \\ < 0 & \text{perhaps it has a positive influence} \end{cases}$$

From the examination of (c) we can use NNs to get more information, we call Nnhope a neural network with multi-layer perceptron architecture with three layers and back-propagation learning algorithm with inputs and outputs equal to the number of cells of the matrices used so far and trained "artfully" to recognize what are the hypotheses, for better or for worse, examined so far on the myelinating diet or not with advantages, or less, on the various musculoskeletal districts examined for each patient subject to the diet.

Using the following scheme we will have to obtain more and more detailed information on how to adjust the diet to follow:

*? The use of certain mathematical operators, matrix ones, and not others, has as its objective the obtaining of a scalar; it would probably be sufficient to use only the calculation of the determinant of the matrix which is the argument of the operator



From the examination of the matrix [hope] the salient data can be reported to the nutritionist and neurologist for the corrections of the case and the various conclusions. The undersigned follows, not at 100%, the indications reported in table 1 and are not detecting worsening but very slight improvements that fall within the fluctuating clinical picture of multiple sclerosis, however the results are expected in the long term. It is believed that this approach will most likely give positive and more visible results in mild forms and in newly diagnosed than in those who have been affected for years by the nrSPMS form like the undersigned who after the end of this work in addition to continuing the diet will dedicate himself more to rehabilitative physiotherapy and physical exercise.

III. METHODS AND TOOLS

This paragraph will describe the main scheme of the setup that will be used, also in the next works and

the present one which mainly describes the idea, and the second one which involves the use of neural networks and the drafting of an algorithm especially by virtue of the fact that the patients will be virtually encapsulated while the third will be much more challenging because the use of "real" patients; at this point in the exposition it is not easy to use real patients.

In the following figures everything is particularly "simplified" because only the preliminary project that will be described is represented, represented in the figures fig. 1, 2. All this is obviously simple compared to the scale of the overall project. As said in the introduction, it is not very useful to describe in depth these blocks that are part of the drawings represented in fig. 1 and 2 because the difficulties that will be encountered will not be few and above all the methods used to describe and realize the various components will not be simple, and the type of representation and its representation is unpredictable.




Fig. 1: This Figure Represents the Original Idea to Train A Neural Network to Distinguish an MS Patient from a Healthy One, as Well as "Memorizing" the Cases Seen in Training

Tab. 1: This Table Represents the "i-th" Patient

Patient _i		Diet _i	Response	Example of a rule indicated by the nutritionist
Food _i	MY/DEMY	RULES/GRAMS	Diet _i	
Dried fruit	My	rule ₁		...
animal proteins	My	rule ₂		0 ≤ 20 [g]
Fruits and vegetables	My	rule ₃		...
Sulphuretin	My	rule ₄		...
Fish + Ω ₃	My	rule ₅		...
Ω ₃	My	rule ₆		...
Biotin	My	rule ₇		0 ≤ 400 [mg]
Vitamins B1, B2, B3, B5, B6, B12, C, Niacin	My	rule ₈		...
Legumes	My	rule ₉		...
Grape seeds	MY/DEMY	rule ₁₀		...
Salt	DEMY	rule ₁₁		0
Sugar	DEMY	rule ₁₂		0
Butter	DEMY	rule ₁₃		...
Oil	DEMY	rule ₁₄		...
Refined foods	DEMY	rule ₁₅		...
Beer	DEMY	rule ₁₆		0
Dairy products	DEMY	rule ₁₇		...
pork sausages	DEMY	rule ₁₈		...
Cereals	DEMY	rule ₁₉		...
Carbohydrates	DEMY	rule ₂₀		...
1 ≤ PATIENT _i ≤ 30	2100 KCal	0 ≤ KCal _i ≤ g _i	RESPONSE 0 ≤ R ≤ 1	$\sum_{i=1}^{20} KCal_i \leq 2100 [KCal]$

Many foods, such as legumes "for example", are very healthy if present in the diet but the effects on myelin are not known, this fact we here would call the "bean problem"

For The patients' response is certainly subjective, but the intelligent system will most likely be able to give useful indications for MD clinicians will be used, hoping to limit the number of inputs, and since they are numerous, it will be necessary to ensure that the NN [6], [12] has a variable and selectable range for the inputs.

At this point in the simulation, it is recommended to use 10 patients for the system training phase, 10 for validation and 10 for testing.

At this point the first neural network should be able to associate food with the response of the sick patient at the onset of multiple sclerosis, but this is not what we would like only. In fact, thanks to the second NN, the most myelinating foods and their quantity to be taken daily should be noted with the supervision of the nutritionist and the neurologist. The following figure further highlights the potential of diet-patient matching highlighted in Figure fig. 1. The male/female ratio is an

important factor for patient selection and the impact of MS, which must be represented in the patient population and therefore will be implicit in the selected sample.

As for training, validation should be done on MS patients, while for the testing phase it would be interesting to insert some patients with variable expressed parameters in the simulations: *new*, *old* and *healthy*, to try to represent the whole population. Overfitting should be avoided by shifting the just evoked parameters and their respective inputs and by a somewhat broader representation of the output and inputs.

The following figure fig. 2 shows the system just exposed; that is, a system able to indicate significant parameters to be provided to the clinician with the totality of patients and simulated data thanks to *Tab. 1*.





Fig. 2: This Figure Represents the Second Part of the System Which Could Give Important Indications to the DIET

To clarify, training can be done on "newly diagnosed" patients and validation on those "diagnosed not long ago" the Test is a somewhat nuanced middle ground, a bit of the first group and a bit of the second, assuming that the system in fig.1 is able to distinguish a healthy patient from a sick one by his diet, then we ask ourselves where the information resides?

The information and the result of the correct training of the NN n°1 of the successful learning of the diet-patient pairing; and up to this point after having carried out the training: then an average of the input vectors of the arrays of the patients and their diets is


made and the significant food could be hidden, in truth the input of interest appears: then a new case is presented and at this point the network will say the most myelinizing diet and the difference is made between the representative vector of the new case minus the average of the patients, then we will see what are the variables in play that determine this difference between the representative vectors. The schemes proposed in fig. 1 and 2 should be considered a common place that can also be used for other pathologies, this aspect is very important to underline.

IV. INDICATIONS ON THE SEARCH FOR OIL



Well	Coordinates		PROFONDITA	
P1	x	y	P	
P2				
P3				
P4				
P5				
...				

V. RECOGNITION OF RADIOACTIVE ATOMS ON NEUTRAL SURFACES



VI. CONCLUSION

Since we intend to proceed, at the end of this mainly descriptive work on the idea of using NNs, other steps are substantially planned that will concern the introduction of over-the-counter drugs into the entrances and then prescribed by the neurologist and with a lot of work to do. Once the correct functioning of the virtual encapsulator of the patient and the entire system has been verified, and the presence of sufficient funds has been verified, to then try to concretely implement the procedure that should answer, in part, the question of the title of this work, thus giving indications to clinical doctors who are experts in the sector covered in this article.

The undersigned, who has been following this diet for a month and more, has noticed great improvements, even if... due to autosuggestion. After all, due to waiting for drugs that never arrives: there's no harm in trying. Funds are needed for software simulations and the undersigned, in addition to making himself available, strongly recommends contacting competent doctors in the various fields of medicine. And like many things in the life of those affected by MS, even a diet must not deprive us of a few moments of happiness. For example, a pizza party with friends (pizza and beer): two non-myelinating foods and we do not participate, aware or not we are not happy but sad and depressed by the renunciation. Such suffering could make us lose the results obtained after months of myelinating diet: the diet should perhaps be faced with a serene, friendly spirit, like the life of a patient, and not only, of MS. We would say balance and serenity: therefore a yes with moderate behavior to pizza and beer with friends.

ACKNOWLEDGMENT

An extraordinary thank you to all those who take care of people with physical and mental disabilities. In particular at my lovely family.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Francesco Pia "I will be back", pp. 149, Ed. Amazon® Kdp, January 2019.
2. Francesco Pia "The liar knee", pp. 224, Ed. Amazon® Kdp, July 2019.
3. Francesco Pia "Genetic geometry", pp. 80, Ed. Amazon® Kdp, January 2019.
4. Francesco Pia "JEWEL: a cosmological model due to the geometrical displacement of galactic object like black, white and worm holes", ICCEA 2021: 15. International Conference on Cosmology and Extragalactic Astronomy; July 19-20, 2021 in Toronto, Canada.
5. Francesco Pia "How to Try to Get Antigravity From Electrical Circuits by an Almost Magnetic Monopole or from Anti Matter" in IRJAES, Volume 7, Issue 3, Date:25/06/2022
6. Francesco Pia "A Computer System Based On Neural Network And Rules System Based For Find An Optimum Mix of Vaccine And/Or Drugs For mRNA Like Covid-19 Virus And Other Pathologies " in IRJAES, Volume 7, Issue 2, Date, 09/06/2022
7. Francesco Pia "POLIFEMO: a Teleportation Model Based on Object Like Black, White and Worm Hole for Electric Signal and Object Also Thinking and With Conscience" ID: IRJAES-V7N3P66Y22 June-Aug 2022 Issue Vol. 7 Issue 3 <https://irjaes.com/volume-7-issue-3/>

8. Francesco Pia "A Symmetry of Cosmological Model Named Jewell That Allows Re Entry Into The Starting Parallel Universe With Adequate Teleportation Based on Creation of Useful Stellar Object Like Black, Withe Holes", ICES-23-P1159, ICES2023, 6-8 February 2023, Riyadh, Saudi Arabia.

9. Francesco Pia "Neutron Concentrator, a Hypothetical Small Neutron "Star" based on the Emission of Neutrons in Minerals such as Granite and Other Commonly Available Objects" Global Journal of Researches in Engineering: E Publisher: Global Journals Civil And Structural Engineering Volume 23 Issue 1 Version 1.0 Year 2023 Type: Double Blind Peer Reviewed International Research Journal Online ISSN: 2249-4596 & Print ISSN: 0975-5861

10. Francesco Pia "A Natural Model for Dark Matter And Its Interaction With Stellar Objects Such as The Galaxy Known As With Name "The Milky Way" American Journal of Biomedical Science & Research, 2023 19(5) JBSR.MS.ID.002630, DOI: 0.34297/AJBSR.2023.19.002630 RECEIVED: JULY 28, 2023; PUBLISHED: AUGUST 07, 2023 PAG. 616-621 FINAL PDF: PP-RW-23-0513 ISSN: 2642-1747

11. Francesco Pia "A Time-Machine, An Electro-Mechanical System That Have A Strange Behaviour Respect At Time" American Journal of Biomedical Science & Research, 2023 20(2) AJBSR.MS.ID. 002681, DOI: 10.34297/AJBSR.2023.20.002681 Received: September 16, 2023; Published: September 21, 2023, pag. 122-126 Final PDF: PP-RW-23-0513 ISSN: 2642-1747

12. Francesco Pia "A Neural Networks And Rules Based System Used To Find A Correlations, And Therefore Try To Maintain The State Of Health, In Patient Affect By Multiple Sclerosis At The Origins Of Well-Being At A Certain Time Daily Time With Clinical And The Musculoskeletal Exams"Global Journal of Medical Research: A Neurology & Nervous SystemVolume 24 Issue 1 Version 1.0 Year 2024 Type: Double Blind Peer Reviewed International Research Journal Publisher: Global Journals Online ISSN:2249-4618 & Print ISSN: 0975-5888

13. Francesco Pia "HOW THE UNIVERSE WORKS", Generis Publishing, 979-8-89248-502-9, 06 October 2024, pp. 66

14. Steve Bryson & colleague, Alec Petersa, Fatima Baninea· Kanon Yasuharaa, Angela Hoffmanc, Basappad Prashant, K. Metrid, Lily Gunningc, Ava Huffmanc, Jake Van Campenc, Clinton C, Shocke Stephen, A. Backf, Larry S. Sherman (Oregon National Primate Research Center from l'Oregon Health & Science University (OHSU)) "Distinct chemical structures inhibit CEMIP hyaluronidase and promote maturation of oligodendrocyte progenitor cells", Journal of Biological Chemistry 28 October 2024; Articles in Press 107916 October 2024, DOI: 10.1016/j.jbc.2024.107916 Also available on Science Direct Copyright: © 2024 THE AUTHORS. Published by Elsevier Inc on behalf of American Society for Biochemistry and Molecular Biology. User License: Creative Commons Attribution (CC BY 4.0) Elsevier's open access license policy.

15. Francesco Pia "How The Artificial Intelligence Whit a Neural Networks And Rules Based System Are Used To Find A Correlations, And Therefore Try To Maintain Ed Improve The State Of Health, In Patient Affect By Multiple Sclerosis At The Origin Of The Food Taken By The Patient", accepted for publication by Global Journal of Medical Research

16. <https://it.wikipedia.org/wiki/Plutonio>





GLOBAL JOURNAL OF MEDICAL RESEARCH: A NEUROLOGY & NERVOUS SYSTEM

Volume 25 Issue 1 Version 1.0 Year 2025

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Toward a Neurocounseling Paradigm: The Science Behind Step One in Addiction Recovery

By Ashton, C. & Duffie, D.

Abstract- Facilitating successful early and sustained remission from substance use disorder remains an extraordinarily difficult puzzle for both clients and supporting persons to navigate. Ninety years since Alcoholics Anonymous developed its twelve step, spiritual approach to recovery from the devastation of alcoholism, initiating and sustaining abstinence from addictive substances remains a tremendous challenge. While the 12-step community continues to support addicted persons at no cost through meetings and relationships, psychology, counseling and medicine are continuing to develop approaches for people to achieve recovery from addiction. To date, there is minimal convergence in these approaches leaving affected individuals with often confusing choices in seeking necessary recovery support. The unfortunate result is the predominant view of substance use disorder as a chronic relapsing disorder whereby relapse is embedded as fundamental to its nature. Greater societal stressors and increasingly dangerous substance access is resulting in death by overdose becoming a major public health issue.

GJMR-A Classification: NLM: WM 270



Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

Toward a Neurocounseling Paradigm: The Science Behind Step One in Addiction Recovery

Ashton, C.^a & Duffie, D.^a

Abstract- Facilitating successful early and sustained remission from substance use disorder remains an extraordinarily difficult puzzle for both clients and supporting persons to navigate. Ninety years since Alcoholics Anonymous developed its twelve step, spiritual approach to recovery from the devastation of alcoholism, initiating and sustaining abstinence from addictive substances remains a tremendous challenge. While the 12-step community continues to support addicted persons at no cost through meetings and relationships, psychology, counseling and medicine are continuing to develop approaches for people to achieve recovery from addiction. To date, there is minimal convergence in these approaches leaving affected individuals with often confusing choices in seeking necessary recovery support. The unfortunate result is the predominant view of substance use disorder as a chronic relapsing disorder whereby relapse is embedded as fundamental to its nature. Greater societal stressors and increasingly dangerous substance access is resulting in death by overdose becoming a major public health issue.

Rapid advances in diagnostic and imaging research techniques over the past two decades, have given tremendous insights into central nervous system functioning at cellular and systems levels. Building on pioneering work in the behavioural neuroscience of addiction, this paper enriches and unifies the often-divergent approaches taken by addicted persons and loved ones, supporting professionals, and peer support groups in addiction recovery. Through a systems neurobiological approach, we outline the progressive neuroadaptations in the limbic system which lead to the condition described by Alcoholics Anonymous as powerless. Understanding the veracity of this central issue in addiction provides an approach we name as Neurocounseling, providing the opportunity for agreement and synergy across approaches. With neuroplasticity as a guiding principle both in progression of the disorder, as well as in the attainment of recovery, addiction can be more appropriately viewed as a life experience rather than a life sentence.

I. INTRODUCTION

In 1939, Alcoholics Anonymous (AA) World Services Inc. published its text which, remarkably, is still in broad use today and shows no signs of losing popularity. Within the profession of caring for people suffering from addictions of all types in residential and ambulatory care sites, the Twelve Steps still remain a gold or, at minimum, a comparison standard. On the other hand, the field of addiction counselling is emerging in response to significantly growing population needs. Addiction Counsellors have developed professional certifications and licensing through

embracing well developed therapeutic modalities (CCPA, 2025) and the scientific method. For the most part, the two approaches are believed to complement each other; it is rare that any current addictions counsellor would dissuade a client from attending a free support group.

The two predominant approaches, counselling and twelve step programs differ significantly: AA clearly advocates its approach through the twelves steps as spiritual in nature (Big Book, Appendix II, 2001). Counselling and psychotherapy are the skilled and principled use of relationship to facilitate self-knowledge, emotional acceptance and growth and the optimal development of personal resources (CCPA, 2025). Counseling modalities have embraced the scientific method for both developing and assessing the effectiveness of its various modalities (Fordham, B. et al, 2021). However, robust long term outcomes studies have not yet taken place which clearly identify evidence for best approaches for addressing addiction since Project Match (*Project Match Research Group*, 1998).

Project Match was a multi-centre study conducted from 1989 to 1997 which has received considerable criticism of its methods in the scientific field (Walters, G., 2002). Regardless, it failed to show any significant difference among any counselling methods and 12 Step programs in outcomes: most enrolled alcoholics showed minimal change in their drinking patterns at the one-year point. Studies since in the areas of treatment outcomes for addictions have been significantly smaller in terms of participants and time length. Our own commissioned work of 2014 (Ashton, C. and Duffie, D., 2014) exploring potential best practices in residential treatment for addictions revealed very few articles, and those centres that did publish reliable outcomes showed a success rate around 20% overall at one year for clients.

In the absence of clear directions for counseling best practices and a multitude of potential options, the treatment industry has currently defaulted to a position which now assumes that each person finds their own path into recovery. Choice of types of therapy is mainly at the client's direction and consensus with the counsellor. There has, however, been a potentially paradigm-shifting body of knowledge which has emanated from advanced brain imaging techniques (PET and functional magnetic resonance imaging (fMRI)) and microbiology research technologies over the

Author a: e-mail: chrisashton@hfhg.ca



past 20 years. What can be collectively termed the neuroscience of addiction has provided incredible insights into relatively uniform and consistent pathways of addiction in the limbic system of the brain. That being said, we have yet seen little of the potential of neuroscience incorporated into present day programs (Verdejo-Garcia, A., 2019).

Our experience in treatment has strongly indicated that those persons who do achieve sustained recovery attribute their success to one predominant ideology that leaves the other far secondary in terms of perceived impact. This may be a suboptimal outcome in many cases. We have developed a novel and extraordinarily successful methodology to addiction counselling whereby the two philosophies merge through the neuroscience of addiction. Counseling and educating clients in the neuroscience behind Step One of AA is an approach that clients find fascinating and immensely gratifying through finally understanding the reasons for the complex array of compulsive and self-destructive thoughts, emotions and behaviours in addiction. Effective in facilitating sustained remission for 83% of clients who complete the 30-day program, understanding the science of recovery through progressive, healthy neuroadaptations allows anticipation of challenges and accelerates the process toward healthy, neurotypical function.

The fields of psychology, addiction medicine and addiction counselling continue to make strides through research and practice. Counselling modalities such as cognitive behavioural therapy (CBT), dialectical behavioural therapy (DBT), motivational interviewing (MI) and acceptance commitment therapy (ACT) have become more refined, formalized and have a growing literature base. Their specific application at the varying stages of change (Prochaska and DiClemente, 1983) has become better articulated.

Drawing on the extraordinary work published in the field, our Step One method uses a robust neuroscientific evidence base to unify current counseling modalities with AA's first step to develop a common, highly effective framework. Coming to terms with the detailed organicism of addiction allows a tangible comprehension of the seeming incorporeal influence of substances over affected persons. At the same time, our program stimulates broader existential questions in contemplating the complexity and terrible perfection of this disorder, merging with the spiritual approach of AA.

a) *Introduction to Applied Behavioural Neuroscience*

Our development of Step One Therapy began with a consideration of AA's Step 1:

"We admitted we were powerless over alcohol—that our lives had become unmanageable."

A difficult admission for people faces even more resistance when the basis for recovery according to the Big Book is dependence on higher power and a spiritual

solution. While it still gains favour among many, an abstract solution to excruciatingly painful, tangible human issues often gathers only minor commitment and unfortunately does not work for the vast majority of persons who try the program (Project Match Research Group, 1998). On the other hand, the counselling profession, as advanced as it is becoming, has yet to produce robust evidence of better efficacy with addiction than twelve step methods since Project Match. Current systematic reviews (Balandeh et al, 2021) of CBT applied to addictions treatment still refer to it as a 'promising practice.' Since Project Match, a thorough review of the literature still shows a paucity of studies whereby adding the two therapies may prove summative.

Years of attending AA and Narcotics Anonymous and listening to people's narratives convinced us that the first half of Step 1, 'we admitted we were powerless,' was true. Since the AA text represented the only means of communicating to a potentially mass audience about this new solution for alcoholism in 1939, it was very carefully written. The Oxford English Dictionary of that era defines powerless as: 'without power or ability; devoid of power, helpless.' AA explains this as losing 'the power to choose whether he will drink or not.' The AA text expands on this state throughout the text speaking of the 'peculiar mental twist already acquired' whereby alcohol always wins out over willpower. Importantly, writing two years after AA formed, the text speaks to its members as 'were powerless,' indicating they had recovered (Alcoholics Anonymous, 1939).

The second part of Step 1, 'that our lives had become unmanageable,' is consistently given witness by the narratives of the varying terrible life circumstances that one comes into recovery with. Our notion in this regard was that this was attributable to more complexity than simply the cost of alcohol or behaviour when highly intoxicated on drugs or alcohol. Rather, when combined with the first half of Step 1, we surmised that the reason behind powerlessness and unmanageable life circumstances was fundamentally a broader disorder of decision making, preceding and progressive with the development of addiction.

Our framework builds on established models, in particular the iRISA (impaired Response Inhibition and Salience Attribution) (Goldstein, R. and Volkow, N., 2001; Kwako, L. et al, 2016; Ceceli, A. et al, 2023) model where the neurobiology of salience of drug cues is postulated to overpower that of other reinforcers with a concomitant decrease in self-control. The iRISA model underpins the familiar cycle in addiction as shown in Figure 1 below:



(Goldstein, R. and Volkow, N., 2001)

Figure 1: Behavioral Manifestations of the I-RISA (Impaired Response Inhibition and Salience Attribution) Syndrome of Drug Addiction

In further exploring the neurobiology of Step One, we additionally postulated that the severe dysphoria, often encountered in withdrawal, had a neurobiological basis which contributed to the overpowering of inhibitory circuitry demonstrated in the prefrontal cortex (Ceceli, A. et al, 2023). Contemplating the seemingly inexplicable relapses in persons in early

recovery just as positive life changes seemed to be occurring, our exploration in this regard was guided by previous work on the anti-reward system in addiction (Koob, G. and Le Moal, M., 2008). The anti-reward system refers to behavioural processes, such as aversion and avoidance, that are part of the normal homeostasis in reward function.




Figure 2: Progressive development of the anti-reward system from initial substance experience (top) to increasing dysphoria and allostatic load (internal physiologic stress) over time with repeated substance use. (Koob, G., 2015)

Just as progressive neuroadaptations develop over time, return to neurotypical functioning has been shown to fail to return to normal until months of abstinence have passed (Koob, G. and Mason, M., 2016). Time to return to normal function is multifactorial and includes the extent of anti-reward development and environmental stresses in recovery (Wemm, S. and Sinha, R., 2019). Portending precarious underlying neurobiology predisposing persons to relapse in the first

months of abstinence, our framework expanded this through careful and often lengthy interviews with clients seeking greater insight surrounding prior relapse events.

Exploring the thought process prior to the decision over which people become 'powerless' in severe substance use disorder, that is taking the first drink or using a substance, revealed one commonality: while people could recall the circumstances surrounding the substance use, they most often had no recollection

of their thoughts immediately preceding consumption. It became quite apparent that there was a momentary conscious lapse, that the decision was made automatically. It consistently appeared as if some other part of the brain was making the decision and conscious thought was overridden. Returning to the iRISA model (Goldstein, R. and Volkow, N., 2001), neurobiological changes in the brain over the course of addiction had pushed inhibition beyond 'impaired' to 'complete,' meaning no conscious resistance whatsoever in many cases. This marked our first direct indication validating the veracity of AA's beliefs with neurobiology.

We then postulated that a discrete neural circuit to drive the fateful decision and actions, existed in the brain which had become empowered unto itself beyond conscious control. Furthermore, looking at the downward trajectory of people's lives in addiction, we postulated this was associated with fundamental decision-making processes that became increasingly self-destructive. Decisions made over the course of addiction became geared solely to creating the conditions for the circuit to activate. Thus, there was a broader impairment in decision making which made life truly 'unmanageable.' Addicted persons had maladaptive brain executive function to the point of being unable to have enough clarity of mind to stop substance use without external help, barring an act of God. Exploring the thought processes of addiction thus

gave a working hypothesis for the biology of Step One, powerlessness and unmanageability being the result of specific brain disorders.

Continuing to expand this hypothesis through deductive reasoning, we were aware of the crucial interaction among emotions and decisions (George, S. and Zane, B., 2016). If the purpose of emotions is not fully to motivate decisions (consistent with the meaning of its Latin root, emovere), at minimum, emotions strongly influence the nature of decisions people make. To explain the reason behind the breadth and compulsive tendency of addicted persons to make decisions which inevitably became destructive, we expanded on research on the limbic system (Goldstein, R. and Volkow, N., 2001; Koob, G. and Mason, M., 2016; Ceceli, A. et al, 2023; Feltenstein, M. et al, 2021) of the brain that had become severely maladapted.

In developing a model for the neural circuit that activated the irresistible compulsion to drink or use substances in severe addiction (DSM V), we believed that decisions corresponded with electrical activity within the brain. As such, a neural circuit could be represented initially as an electrical circuit and explored through this perspective (Illes, J., 2005; MacGregor, R., 1987; Singh, J. and Kapur, G., 2019; Cho, S. et al, 2021; Harrison, R. 2008). An NMOS transistor (Chin, A. et al, 2019) amplifier provided a working model:




Figure 3: An NMOS transistor acting as a binary circuit: when the voltage at G, V_{GS} exceeds the threshold voltage for the transistor semiconductor, V_T , electrical current travels through D to S

Depending on the material nature of the semiconductor (Chin, A. et al, 2019) and the polarity and strengths of the input, a small current at the emitter (D) can be amplified significantly at the output collector (S). In similar fashion we theorized that a small impulse to use substances could pass through a neural network analogue and become amplified to overpower all circuitry in the executive part of the brain. This would model the impulse to use substances at D, overpowering the inhibitory circuitry of the prefrontal cortex at G and producing an amplified output (to consume substances) at S. In neurobiological terms, the result would be powerlessness over the decision to use substances.

Given that this transistor-like amplifier circuit seemed so far beyond reach in the brain (most persons with addictions have tried many times to abstain and sought professional help often to no avail), our therapeutic target became the inputs (the prefrontal cortex, modelled by V_{GS} above). Should neurotypical function of the inputs be established, the internal circuit could potentially be brought under control. We also hoped that the actual physical composition of the semiconductor-like circuit at G would resultantly return to healthy.

From such reasoning did the union of neuroscience, 12 Step programs and counselling emerge. A concrete understanding of the neurobiological truth of Alcoholics Anonymous' Step One with counselling addressed at the inputs (thoughts, emotions and behaviors) offered a novel, potentially dramatically effective method of helping people afflicted by addiction.

Additional logical assumptions underscored what became the Step One approach: since we were dealing with the brain, the principle of neuroplasticity applied. Neuroplasticity has only become well accepted this century and refers to the modifiability of the brain (Innocenti, G., 2022). Changes in both structure and, consequently, function of the brain are inherent properties that occur throughout life with greater propensity for this in the young developing brain. Nonetheless, neuroplasticity occurs throughout the entire life span. It is not necessarily exclusively adaptive to external stimuli but can also be the cause of neurological and psychiatric pathologies (Innocenti, 2022).

Further logical hypothesis expansions were made prior to the rigorous research to explore our notions: given that addiction seemed to develop over time to eventually dominate behavior, neuroadaptations were most likely to be created within previously healthy circuitry. Addiction was not a separate neural circuit from the rest of the brain, rather it was fully integrated with malevolent neuroadaptations in previous healthy networks. Without definitive healthy neuroplastic changes, the overall state of addiction would continue to

exist. Of course, changes in structures and networks would be iterative and at times synergistic, leading to a non-linear progression. This fully matched our observations of persons over the course of recovery. The principles of neuroplasticity also offered the prospect of full recovery, as Alcoholics Anonymous had proclaimed in 1939 after only two years of its formation. A disorder could be reordered.

b) Basic Science Framework

With experience, observation and access to rigorous scientific articles available for our methodology, we explored models from several disciplines to form a framework for research. The American Psychiatric Association defines addiction as a complex condition – a brain disorder that is manifested by compulsive substance use despite harmful consequences. The limbic system itself is markedly complex both in structures and interconnectedness (Catani, M. et al., 2013; Volkow, N., Michaelides, M. and Baler, M., 2019), not to mention introducing neuroplastic changes (see Figure 4).

We chose to frame our exploration of the behavioural neuroscience of addiction through the lens of a complex dynamic system (Close, C., Frederick, D., Newell, J., 2002). A system is any collection of interacting elements for which there are cause-and-effect relationships among the variables. While a reductionist approach (Mazzocchi, F., 2012) is often used in complex systems and was an initial phase of our exploration, the systemic approach insists on an understanding of the interaction among variables in modelling. The specific system model under investigation was the transistor amplifier circuit, seen as having multiple, likely interconnected inputs and a semiconductor element with variable material composition. The term complexity refers to both the number of variables (inputs) and the interactions among all.

The descriptor, dynamic, in complex systems acknowledges that the variables and relationships among them change over time (neuroplasticity). In our case, changes may or may not be entirely adaptive to external environmental stimuli but are dependent on all other system elements. Fortunately, overall, there is relative stability of the system over days to months to years despite individual elements functioning in millisecond timeframes. With most neural network functioning at generally low frequencies (<100 Hz), current diagnostic methods in micro and macrobiology reveal considerable not only about structure but most importantly, function.

To date, we have identified numerous areas and major nuclei in the limbic system involved in addiction. Each of these have distinct subregions and multiple functions. Additionally, there are a number of potential physiological neuroadaptations at the neuronal level for

each neuron in the system. All areas and nuclei communicate with each other and information transmission integrity is subject to change although there is consistency in this aspect of neuroplasticity among all connections. Fortunately, the inter-element

communication is dominated by a number of discrete pathways simplifying understanding in this regard. This array of intercommunicating elements with a brief description of each is found in Tables 1., 2., and 3. and illustrated in Figures 4., 6., 7. and 9.

Table 1: Structures and Areas of the Limbic System Implicated in Addiction


Limbic Element	Brief Function(s) relevant to Addiction	Neuroadaptations in Addiction
Amygdala	Signals fear, anxiety, aggression, anger, male sexual impulse	Grows in size, hyperexcitable and hyperreactive
Hippocampus	Memory associated with emotion, also regulates cortisol in stress response	In active addiction, new information storage declines except regarding substances
Hypothalamus	Signals stress response from limbic system to the body	Maintains an elevated, less responsive baseline function
Lateral Habenula (LHb)	Important in aversion behavior, amplifies negative input and signals back to rest of system	Becomes increasingly sensitized to substance cessation
Mammillary Bodies	Involved in transmission of memory information, reward prediction	Function declines, especially with alcohol
Nucleus Accumbens (NAc)	Motivation for reward, reward prediction error, limbic-motor interface	Increasingly blunted DA response, greater influence over motor as PFC declines
Prefrontal Cortex (PFC)	Executive control over decision making, emotional regulation, working memory and calculation/planning, risk/benefit	Decline in all functions, appears (by deduction) to focus more on substances
Rostromedial Tegmentum (RMTg)	Reacts to negative DA in VTA, signals LHb, aversion	Responsive to substance cessation, temporarily relief if substances planned
Thalamus and Insula	Distributes external and interoceptive (Insula) stimuli to rest of system	Some literature points to preferential bias to transmit substance related information
Ventral Tegmental Area (VTA)	Signals want or need to NAc through DA efferents	Increasing response to substance related cues

Table 2: Communication Pathways Involved in Addiction

Communication Pathway	Major Elements Connected (note pathways are bidirectional)	Function
Mesolimbic pathway	VTA to NAc	Physiological DA response
Amygdalofugal Pathway	Amygdala to Hippocampus	Automatic fear response
Hypothalamic-Pituitary-Adrenal Axis	Rest of limbic system signals Hypothalamus	Stress response
Medial Forebrain Bundle	PFC to all elements of limbic system	Emotional regulation, executive decision making
Mesocortical Pathway	PFC to NAc	Reward prediction & analysis, executive decision input

Table 3: Mechanisms of neuroplasticity

Neuroadaptations at cellular level (neuroplasticity mechanisms)	
1.	Mitosis (cell duplication in younger people)
2.	Myelination or demyelination (affects transmission among neurons)
3.	Synaptic growth or recession
4.	Dendritic branching (recession or growth)
5.	DA receptor up or down-regulation
6.	DA transporter up or down regulation
7.	DA reuptake up or down regulation



Where: PFC represents Prefrontal Cortex, OFC is orbitofrontal cortex (within the PFC in our model), ACC is anterior cingulate gyrus, Thal is thalamus, Hippo is hippocampus, DS is dorsal striatum

Figure 4: From Volkow, N., Michaelides, M. and Baler, M. (2019). The Neuroscience of Drug Reward and Addiction

What seems a near impossible task to consolidate and understand is made manageable through, first, a reductionist approach identifying anatomy and function of the specific elements. Fortunately, there is general uniformity of basic neuronal structures and networks, as well as commonalities of thinking, emotional and behavioural patterns of addicted people. Most of our clients, virtual or residential, are able to grasp an adequate understanding of this material and self-insight within 45 days to establish sustainable recovery.

c) Specific Findings

We explored our model of an NMOS amplifier transistor as the decision circuit at the centre of

addiction, in the context of consensus surrounding the neurobiology of addiction being dominated by 'the dopamine hypothesis.' This was put forward convincingly by a pioneer of addiction neuroscience, George Koob of The Scripps Research Institute (Koob, G., 1992). This hypothesis has been the focus of significant research to present day and remains the most popular theory underlying the neuroscience of addiction. Addictive substances are classified as such because they all create a dopamine rise in a part of the midbrain (assuming the triune brain anatomically termed the nucleus accumbens (NAc).




Figure 5: The triune brain schematic

A rise in dopamine is temporally associated with the high of addictive substances. The dopamine system projecting to synapses in the NAc is still considered by most to be the center of reward, the reward system.

This view is unfortunately incomplete when one explores further the complexity of addiction. We also believe that the prevalence of theories which emphasize an uncontrolled reward system as a fundamental issue

that continues to maintain societal stigma of addiction. Through considering persons become so addicted to the reward of substances that they don't want to stop, implies continued choice on the part of addicted persons.

Through a reductionist framework, we deducted that it was an excess of dopaminergic neuron excitation, V_{GS} (compared to inhibitory neurons, V_I) that ultimately activated the circuit to use substances as modelled by the NMOS transistor (Figure 3.).

An exploration of the dopamine (DA) neuron reveals much when considered in the experiential context of addiction. In brief, chemicals including addictive substances bind to receptors on dendrites (they look like Christmas trees) on the cell body and create electrical charge which then creates an electrical current which travels down the axon. Axons are insulated with myelin and the electrical charge, on reaching the pre-synaptic terminal, activates dopamine transporters (DATs) (see Figure 7.). DATs then bring vesicles of dopamine (like bubbles containing DA) to the membrane and release DA into the synaptic cleft. In this space, DA then binds to dopamine receptors (DARs) of

the next neuron. In the binding process, Ca^{+} molecules are released. This leads to accumulation of positive charge in the post synaptic terminal and an electrical current which then travels down the axon of the ensuing neuron. This simple model explains how substances which activate one neuron then turn on the next neuron in the neural network, which could be, say, the experience of reward.

Recall that we are considering the midbrain which is accepted to be essential for survival through maintaining physiological stability referred to as salience. Clearly, addictive substances, which dramatically elevate DA levels and transmission, threaten salience. In a response to 'control' the dopamine response and maintain salience, through mRNA transcribed through genes in the nucleus, the neuron creates proteins which, at the synapse (Sulzer, D., 2011):

- Down-regulate DATs
- Down-regulate DARs; and,
- Up-regulate DA reuptake transporters




Figure 6: D2Rs (DA2 Receptors) in cocaine users. Harvard Medical School (2021). Understanding Addiction

As a result, over time, more of the substance is required to achieve the same electrical energy as occurred with the first experience. Additional adaptations which the neuron can perform are duplicating itself (in developing brains, mitosis) and change dendritic configuration. Advanced substance

use can also affect the integrity of the myelin sheath, giving rise to conductive loss and crosstalk (Ashton, C. & Duffie, D., 2022). Finally, the neuron can create new synaptic connections when so stimulated, a process referred to as Hebbian plasticity (Turriagano, G. and Nelson, S., 2000).




Figure 7: Schematic of the dopamine neuron and synapse

A few fundamental associations can be made through an understanding of neuroadaptations at the cellular level. Clearly, neurons can and do change through distinct mechanisms. Summarily, these help to demystify neuroplasticity in very simple terms. Second, a quick study of the neuron informs us about the science behind tolerance to substances which all addicted persons develop over time. Finally, this contributes to an understanding regarding the 'powerless' aspect of Step One and the inability to stop at one drink when driven to reach previous euphoria (trying to achieve the previous dopamine high in the face of resistant circuitry).

d) The Drive Towards Substances: The Nucleus Accumbens And Ventral Tegmental Area

Assuming the neural circuit to consume substances exists in the dopaminergic system within the nucleus accumbens, further proof of this mechanism as well as identification of inputs forms the next logical exploration. We were guided by much superb literature on the reward side of addiction (Volkow, N., Michaelides, M. and Baler, R., 2019), as well as partially conflicting literature surrounding the NAc in isolation (Floresco, S., 2015). There is good research confirming that the NAc is not simply a 'reward centre' responding proportionately to dopamine afferents. Rather, it is far more sophisticated and functionally aligned with the powerlessness aspect and narratives given and described as: 'there was always the curious mental phenomenon that parallel with our sound reasoning, there inevitably ran some insanely trivial excuse for taking the first drink (AA Basic Text, 1939).'

Returning to the triune brain model, the NAc is a deep structure within the survival-gearied midbrain. Executive thought and consciousness are generally accepted to exist in the frontal cortex. Floresco (2015) in his broad review states that the current emphasis on the NAc being the reward center has veered from the more general idea initially proposed by Mogenson (Mogenson et al., 1980) that this region serves as a limbic-motor interface. Emerging evidence further supports the NAc as having diffuse projections of axons to cortical motor areas (Salgado, S. and Kaplitt, M., 2015; Sawada, M. et al, 2015; Miyachi, S. et al, 2006). Additionally, while the focus is generally on the projections from the Ventral Tegmental Area (VTA), where the dopamine cell bodies reside, to the NAc, little is made of the bidirectionality of circuitry between the VTA and NAc.

Corkrum et al (2020) speak strongly to this as well as Floresco (2015) and Yang et al (2017). Logically, bidirectional communication better explains the actual functionality of reward driven behaviour as well as aversion (which will be explained later). While NAc dopamine synapse activation is seen in anticipation and reward experience, we suggest that the hedonic experience takes place through further projections to the

basal ganglia and ventral pallidum (there are few known neurotransmitter sites typically associated with the pleasant feeling of reward within the NAc itself) (Root, D. et al, 2015; Castro, D. et al (2015). With the VTA accepted as signalling 'want or need,' it makes most sense that major inputs to create this emanate from the NAc as Corkrum et al (2020) suggest.

We suggest a primary role for the NAc as a dopamine regulator to maintain salience, and a motor interface (mediated of course by the PFC). Dopamine itself in the mesolimbic system (VTA and NAc) then also acts as a motivator of action for which there is much good evidence (Wise, R., 2004). Significant for our understanding of addiction is the NAc function of measuring reward prediction error (RPE), essential to learning and memory (Yang et al, 2017). This argument for the multiple functions for the NAc and VTA will make further sense as we consider the role of avoidance (we term 'the anti-reward system') as equally significant as the positive reward system in addiction.

We know that as addiction develops, there is progressive decrease in reactivity of DA synapses in the NAc. Termed tolerance, there is increasingly less response in the NAc for the same dopamine signal from the VTA as the mesolimbic system strives to maintain salience. Additionally, in advanced addiction, numerous authors (Corkrum et al, 2020; Febo, M. et al, 2018; Koob, G. and Mason, M., 2016) suggest a deficit in extracellular dopamine creating what is functionally a hypodopaminergic state in advanced addiction. Returning to Step 1, this is a major contributor to craving as experienced as part of the powerless over addiction phenomenon.

People with severe addictions will admit being unable to perform even the most minor activities of daily living, let alone function at all in a work environment without substances. This is unfortunately true physiologically: in severe addiction, the hypodopaminergic state resulting from resistance which has developed in the neural circuitry along with potentially low basal extracellular dopamine levels has a devastating effect on motivation. The simplest things such as showering and getting dressed seem like a monumental task without topping up on substances.

While this may appear as a willful lack of care of self or a manifestation of self-loathing on the part of severely addicted persons, they simply do not have the motivational energy through the NAc. This terrible phenomenon is in addition to the relentless emotional and thought dysphoria persons experience. In a complete hijack of healthy functioning, the limbic system has now equated substances with survival, as only substances will provide the necessary dopamine response to feel functional.





e) *Failure to Inhibit the Drive to Substances: The Prefrontal Cortex*

The limbic system, and those elements specifically involved in addiction, is a complex dynamic system unto itself whose purpose is the maintenance of homeostasis. Homeostasis can be considered persistence of a living system and society in a state of specified and dynamic disequilibrium (Pross A., 2012). System wise, the interconnected limbic elements can be thought of as alternatively competing and cooperating with each other in active striving towards a persistent and specified orderliness (Turner, J., 2019). Viewed as an internal environment, the limbic system maintains homeostasis through adaptive interfaces (the various elements), which monitor external environments.

As a living, closed thermodynamic system unto itself, geared to maintaining the greater intracorporeal entity, energy is continuously required to survive in a dynamic environment. Knowledge and cognition are equally needed within the limbic system to construct cognitive representations of the environment in which it is embedded, with knowledge of what the contained environment should be, and continuously seek reward to maintain homeostasis in the face of ongoing perturbations to it (Feldman Barrett, L., 2006).

Implicit in this reward seeking function and processing external cues is the Nucleus Accumbens. As one of its critical functions, the NAc calculates reward prediction error (RPE) (Floresco, S., 2015) whereby positive RPE drives anticipation, motivation towards and memory of reward events. The primary neurotransmitter involved in RPE is, of course, dopamine. Resultant neurocircuitry of the midbrain in reward seeking and survival are motivation to activities, behaviours and external substances that result in dopamine increases in the mesolimbic system. Emotions are created through this process.

In simple terms, the elements of the limbic system react to internal and external stimuli, process it through intercommunication including the PFC to drive the mesolimbic system to drive decisions to maintain overall dopamine homeostasis, in an energy efficient manner as possible. Equally, the system creates negative RPE to direct diversion and avoidance. Proof of concept is highly supported by reviews of Deep Brain Stimulation (DBS) for addictions which is aimed at the Nucleus Accumbens (Chang et al, 2022). Electrical stimulation of the NAc through DBS acts as a system override of the rest of the contributions to the NAc of the limbic system.

Addictive substances and behaviours in vulnerable people all produce a positive rise in dopamine levels in the synapses of the NAc. This rise can be multiples of baseline levels and result in significantly elevated RPE initially on consumption. Regardless of input from the PFC, the midbrain interprets substances as contributing to positive valence

(Feldman Barrett, L., 2006). Despite societal beliefs influencing the PFC to inhibit substance use, the midbrain is susceptible to dominate the decision-making process and repeat the experience. In such a manner, addiction progresses to reach the widely accepted three-stage cycle of addiction as described by Koob (2001), Volkow (2018) and their associate researchers.

While acknowledging that participation and influence from other limbic structures is continuously present, focusing on the basics of the neuroscience involved in Step 1 does not detract from the veracity of this hypothesis. During the last decades, many studies have shown that addiction results in heterogeneous cognitive deficits (Pitel et al., 2023), including impairments of executive functions, memory, visuospatial abilities, difficulties in emotional processing, and theory of mind (ToM) abilities (Le Berre et al., 2019). ToM is defined as the capacity to infer mental states from others' social signals to predict their behaviors, desires, intentions, and beliefs. It is now well-known that attention, working memory, and executive functions rely notably on the prefrontal cortex (Maillard, A. et al, 2020).


Executive functions are cognitive abilities that control and regulate the emotional (survival) system to coordinate thoughts and actions toward a goal. Executive functions include mental flexibility, abstraction, planning, problem-solving, shifting of mental states, monitoring and updating of working memory representations, organization, rules deduction, reward delay discounting, and categorization (Maillard, A. et al, 2020).

There is a considerable and emerging literature concerning the role of the prefrontal cortex in emotional regulation, much of it focussing on the PFC and amygdala communication (Suzuki, Y. and Tanaka, S., 2021; Dixon, M. et al, 2017). Unfortunately, less research has been performed regarding the relationship between the PFC and the nucleus accumbens. The connection between the two is referred to as the mesocortical pathway. Within the mesocortical pathway, the anterior cingulate, prelimbic cortex, and orbital prefrontal cortex (OFC) have been accepted as being involved in the regulation of emotional responses, cognitive control, and executive function (Volkow et al. 2019).

Much of the literature points to neuroadaptations in the various aspects of the PFC as leading to persistent addictive behaviors, including the devaluation of natural rewards, diminished cognitive control, and hyper responsiveness to drug-associated stimuli (Koob, G. and Le Moal, M., 2001; Kalivas and Volkow, 2005; Carelli and West, 2014). Generally, the view continues that changes in the PFC contribute to escalation of substance use through mediating conditioned responses, drug craving, and loss of behavioral control (Feltenstein, M. et al, 2021).

In light of our observations of clients with severe substance use disorder over the first 30 to 45 days of recovery (abstinence), neuroimaging studies(Goldstein, R. and Volkow, N., 2011) and additional studies focusing

on the effects of chronic stress and substances on the PFC (Koob, G. and Schulkin, J., 2019), we offer an alternative view which is best observed in recovery.



Healthy Control After 2 years cocaine abuse

Figure 8: Harvard Medical School (2021). Understanding Addiction. OFC refers to orbitofrontal cortex

As shown in the scan above, neuroimaging studies reveal an emerging pattern of generalized prefrontal cortex (PFC) dysfunction in drug-addicted individuals. This worsening PFC hypofunction is coincident with more drug use, worse PFC-related task performance and greater likelihood of relapse. Widespread PFC activation is seen in drug-addicted individuals upon taking cocaine or other drugs and upon presentation of drug-related cues. We believe that this is compatible with dopamine surges from the NAc and VTA to the PFC, which then creates a transient focus on obtaining the drug of choice or otherwise. We additionally purport that a degree of PFC dysfunction precedes drug use, creating vulnerability for susceptible persons in developing substance use disorders.

Secondly, studies in neuropsychopharmacology demonstrate the molecular effect of stress hormones (Corticotropin Releasing Factor or CRF and cortisol) and substances together and alone in SUD (Wemm, S. and Sinha, R., 2019).

Accordingly, we assert that rather than PFC adaptations as exclusively pursuing substances (which is true to a small extent), the major issue is with the loss of PFC ability to regulate, and specifically inhibit, emotional and motivational drive from the midbrain. This is well observed as well as is partial recovery of the PFC in our clinical experience. Much of the behavioural impulsivity and extremes in emotions and compulsions seen in early recovery (ask anyone at AA about this phenomenon) is in large part due to loss of PFC inhibitory function.

In cognitive testing using the Creyos platform™, a robust assessment tool developed through combining neuroimaging with neuropsychological studies, clients uniformly scored well below average in virtually all domains regardless of their backgrounds. Fortunately, at the 30 day mark, cognitive scores improved to normal in general. Additionally, scores using Muse 2™neurofeedback meditation for 15 minutes daily improved substantially over the time in residence with us. This also supports PFC reactivation, a key goal of our program.

Interestingly and of tremendous clinical significance, Muse 2 scores declined greatly for all clients consistently over a two-day period at our center between 8 to 14 days from admission. Prior to Muse 2, the period from 8 to 14 days had been a time where clients would regularly insist on self-discharging, suffer severe anxiety at times and exhibit fully irrational thinking. We believe that this period refers to the time whereby the hypothalamus begins to resume normal activity (Ashton, C., 2014) and the resulting stress response is excessive until the hypothalamus achieves a new baseline activity over 2 days.

Emotional management including impulsivity is a challenge for both staff and clients, particularly in the first two weeks in residence. Also frustrating to clients is loss of working memory and inability to remember educational material, as well as orientation to place from one day to the next. Again uniformly, working memory returns dramatically for clients (much to their relief) around 18 to 20 days in the program.



To a large extent, this severe hypo functioning of the PFC explains powerlessness due to lack of inhibition over the NAc. Further research adds to understanding the magnitude of the drive to obtain substances and associated high risk behaviours. It is much more than a lack of inhibition and a notion which is slowly becoming more recognized in the scientific community (Ashton, GAB 2024). Recall the NMOS transistor amplifier model: in addition to acting as a binary circuit, the semiconductor is fully capable of amplifying the strength of the input signal. Strength of amplification depends on collateral inputs and the physical properties of the semiconductor.

Similarly, the decision to obtain and consume substances in persons with severe addiction often overwhelms any thought of consequences. This is the point where otherwise unconscionable acts are undertaken in substance use disorder, often resulting in severe, traumatic consequences as well as imbedding a seemingly hopeless level of shame.

In pursuit of understanding this third element creating powerlessness in severe addiction, we undertook to find the neurobiology which amplifies motivation to consume substances. Simple Pavlovian learned behavior is considered a fundamental process in the development of substance use disorder of which there is much literature (Day, J. and Carelli, R., 2007; Heinz, A. et al, 2019). On this basis, we explored the dopamine profile in the mesolimbic from steady state to being cued (small release of DA) to the full DA spike experience corresponding to reward.

It is well known in Pavlovian learning, this process becomes embedded such that reward is experienced in the mesolimbic system prior to the actual consumption and physiologic DA spike. This corresponds to narratives given by many clients that the anticipation of reward is often higher than the consumption experience itself. Additionally, if they do not experience the rewarding feeling anticipated, they become agitated and dysphoric. Complete omission of the reward following experiencing the high of anticipation results in further, often deep dysphoria. This phenomenon is articulated beautifully and termed reward prediction error (RPE) by Keiflin, R. and Janak, P. (2015).

RPE remains fundamental to the development of addiction. Assessment of a potential reward is predicted within the NAc through receiving information from other limbic areas and the cerebellum. This forms a prediction of the reward. What is physiologically the actual result of the behaviour or substance (i.e. the DA response) is measured as well by the NAc calculating reward prediction error. The magnitude of RPE (positive or negative) results in proportional glutamate projected to the hippocampus, as well as the VTA, facilitating encoding of memory. Simply put, the brain remembers things that are unexpectedly better than anticipated

(+RPE) to be able to repeat again. Equally, the brain remembers things which resulted in an experience worse than predicted (- RPE) to avoid repeating (avoidance learning).

Often termed the anti-reward system (Koob, G., 2015; Borsook, B. et al, 2016), this third element in powerlessness is as, if not more, influential as the reward seeking aspect in severe addiction. Negative DA states (Wise, R. and Robbie, M.; Keiflin, R. and Janak, P., 2015) are highly unpleasant and feel excessively fatiguing. There seems no healthy, fast way out of this state, barring external substances.

Reward omissions, as well as aversive outcomes, activate neurons in the lateral habenula (LHb) (Matsumoto, M. and Hikosaka, O., 2007), which indirectly inhibits DA neurons via activation of GABAergic neurons in the rostromedial tegmentum (RMTg). A role for involvement of the LHb in negative prediction errors has been established in animal studies (Keiflin, R. and Janak, P., 2015).

It is acknowledged that reduced activity in a large subset of dopaminergic projections to the nucleus accumbens is associated with aversion. Optogenetic inhibition of ventral tegmental area dopamine neurons produce avoidance of associated places and cues, an effect that depends on DA receptor signaling (Danjo et al., 2014). Inhibition of these neurons via DA receptor activation causes a conditioned place aversion and gives rise to negative affect (Liu et al. 2008). Aversive stimuli of various sensory modalities broadly inhibit dopamine release in the VTA and NAc (de Jong et al., 2018; Menegas et al., 2018).

This is considered a fundamental mechanism for aversion, a function necessary for survival. To put the neurobiology of addiction in perspective, recall our discussion on the mesolimbic pathway. This becomes so severely maladapted that substances are interpreted as necessary for survival. Following that logic, the severely addicted brain now sees absence of substances as a threat to survival and accordingly imposes its avoidance system to deter the person from stopping substance use.

At the prospect of abstinence, it is typical for addicted persons to respond with panic and often anger automatically as the avoidance defence manifests. Unexpected inability to obtain and consume substances is met with an often-prolonged state of severe dysphoria in severely addicted people. Superimposed on a hypodopaminergic state is sudden absence of substances following an anticipatory state, resulting in a negative dopamine spike (Keiflin, R. and Janak, P., 2015). This appears most relevant to the rostromedial tegmental nucleus (RMTg) aspect of the VTA, an area highly associated with registering and encoding aversion (Jhou, T. et al., 2009) which signals inhibitory control over DA neurons in the VTA.

The RMTg is located adjacent to the lateral habenula (LHb). There is a recently emerging literature surrounding both the RMTg and LHb which explains neurobiologically the terrible drive behind addiction (Suzuki, Y. and Tanaka, S., 2021). Simply put, in healthy persons, aversive stimuli activate lateral habenula neurons which, in turn, inhibit dopamine release in the VTA and mesolimbic pathway, thereby inhibiting positive action and creating aversion. As previously identified in this paper, two elements bear heavily on the science of powerless: reduced dopamine response (a chronic hypo dopaminergic state) and a faulty prefrontal cortex, unable to control impulses to consume substances.

The amplification of drive emanates from the LHb and a 180° hijack of the avoidance system. As addiction progresses, the dopaminergic mesolimbic system has been so fundamentally maladapted that it associates substances (natural rewards no longer are able to produce needed dopamine response) with survival; conversely cessation of substances is equated as a crisis to be avoided at all costs. This is particularly evidenced clinically in acute withdrawal, most noticeably in alcohol, opiate and benzodiazepine withdrawal. The pain, dysphoria and physical symptoms from activation of the anti-reward system become progressively severe to the point of becoming unbearable for persons with severe addiction.

With multiple episodes of full or partial acute withdrawal occurring over development of addiction, the anti-reward system becomes increasingly 'kindled' (Breese, G. et al., 2005), meaning increasingly sensitive and hyperreactive to negative stimuli. This is due in large part to the LHb becoming increasingly fierce in its response through progressive neuroadaptations due to stress and substances. While not completely understood (it is hypothesized that cell membrane changes are responsible (Authement, M. et al., 2018), progressive adaptations in the LHb create amplification of negative stimuli (Graziene, N. et al., 2018). This produces increasing distress with each attempt at stopping substances, an experience virtually all addicted people will attest to if they've had multiple attempts at quitting.

With responses to negative stimuli already magnified in other areas of the limbic system (the amygdala and hypothalamus particularly), this is then amplified by the LHb creating unbearable dysphoria. As well, with the kindling effect and dysphoria experienced with withdrawal episodes, this creates learned behavior to avoid abstinence in severely addicted people. Clinically, for persons in the contemplative stages for recovery, it is not uncommon for them to say they feel like they will die if they have to quit substances. For the caring professional, it is most helpful that this seeming complete irrationality is not driven by a compulsion to get high. Rather the anti-reward has created this level of thorough distortion neurobiologically: what appears as

ambivalence regarding quitting, is truly an excruciatingly frightening prospect for the addicted person.

While not necessarily described by the DSM 5, it our firm belief that, over the course of development of substance use, the anti-reward system becomes dominant over reward and presents a daunting challenge for caregivers to support the client in overcoming this challenge. Typical for addiction, the level of discomfort anticipated by addicted people in quitting substances is often higher than the experience of withdrawal itself. Given this shift to the major obstacle now being anti-reward and with concurrent diminishing of the reward system, clients will now generally admit to using substances to function, to try to feel and act normal, and to not get physically and mentally sick.

II. DISCUSSION

Understanding severe addiction through a neurobiological lens is most helpful in informing caregivers of what is actually happening in the addicted persons world at that time. Given the advancement and failure of healthy decision-making processes, it is easy to become enrolled in trying to understand the chaotic situation addicted persons' lives have become, to make sense of this complexity. Understanding this neurobiology then gives evidence and a lens for the counselor to focus their best modalities to support the person out of ambivalence and make the decision to cease substances.

The first part of Step One of Alcoholics Anonymous, we admitted we were powerless over alcohol, now has a concrete neurobiological explanation and proof of veracity. Three separate main processes converge to not only making cessation seem impossible and unwanted, but also an overwhelming neural energy to consume:

1. Severe hypo functioning of the prefrontal cortex;
2. Resistance developed in the mesolimbic circuit to experiencing positive reward; and,
3. Amplification of negative valence by the anti-reward system.

The poorly functioning prefrontal cortex simply does not have the electrical inhibitory power to stop the consumption drive created by the reward and, later, the anti-reward system.

As most alcoholic and addicts will verify in substance use disorder, the immediate thought process prior to acquiring and consuming substances cannot be recalled in any detail. The midbrain has completely taken over conscious thought; most people with addictions when questioned in detail will speak of some unknown sort of force carrying them through the acquisition and consumption phase. At best, they will report some internal chatter (PFC) suggesting that it was not a good idea but did not have the force to overcome the strange drive.



Given the extent of neuroadaptations which have occurred, it then becomes easier to understand and validate the often-unbelievable narratives we hear in addictions counselling. As an example, I received a distressed call from a young professional man who had finished our program and returned home 3 weeks prior to the call. He told me that he was sitting at his kitchen table with an unopened bottle of spirits in front of him. He had no recollection of driving to the liquor store and making the purchase, but acknowledged that somehow, he must have. Recall that the NAc has connections to the motor cortex.

The man had found himself on returning home suddenly under extreme family stress which included involvement of the justice system. Stress hormones, particularly CRF and cortisol, impair prefrontal cortex functioning in similar manner to substances in severe addiction (Sequeira, M. and Gourley, S., 2021); Wojdala, A., Molins, F. and Serrano, M., 2020). In this man's case, stress had overwhelmed the PFC to such an extent that the NAc was able to completely hijack his actions: he had travelled and purchased the bottle without conscious awareness of what he was doing, let alone the potentially tragic consequences.




Figure 9: The Powerlessness Circuit. PFC: Prefrontal Cortex, showing disempowered inhibitory (X) influence over a. Mesolimbic (reward) pathway of VTA (ventral tegmental area), RMTg, (rostromedial tegmentum) projecting to Nucleus Accumbens, and b. Anti-reward system main parts being Amygdala, Hypothalamus, Hippocampus, and Lateral Habenula. Result is anti-reward and mesolimbic pathway in full control.

While a striking example of full hijack of the brain by the NAc, his experience is all too common in severe addiction and early recovery. Physiologic stress is always the cause underlying return to substances both in early and later recovery (Koob, G. and Schukin, J., 2019) and a tremendous impediment to recovery of neurotypical brain function. Counselling can be

exceptionally effective at reducing allostatic load through facilitating new perspectives and solutions.

Unless one is very experienced in the field or has had lived experience with addiction, the irrational self-destruction seen is very difficult to comprehend or validate. Moreover, as substance use disorder has also affected most of the rest of the limbic system structures, thoughts, perceptions, explanations and actions

become increasingly distorted. Emotions which are typically a challenge for everyone in life, now become neurobiologically distorted by the structures and system which creates them. For addicted persons, the false thought process becomes fully illogical in that what they firmly believe is keeping them alive is actually bringing on a premature and painful death.

Demystifying powerlessness in addiction is generally met with great relief by clients coming into recovery. It gives concrete answers to a seemingly incomprehensible constellation of symptoms, behaviors and actions that defy rational thought. Understanding the neurobiology of addiction can reduce the persistent self-shaming people in addiction and early recovery experience, as well as allowing clients to focus on the processes that definitively facilitate brain recovery.

Neurocounselling, Step One Therapy, is predominantly educational and focused on giving clients an understanding of what addiction truly is so that they can see its dominant influence on their life and reverse its course over time. Counselors trained in the behavioural neuroscience of addiction are able to effectively facilitate client self-insight into their past and lingering inability to make healthy decisions, knowing that addiction remains a potentially dominant force in early recovery. The educational process to facilitate the client arriving at a level of insight to accept the neurobiology of Step One as tangible, requires proficiency in near all the current counseling modalities. Even though insight is being developed, return to neurotypical brain function does take time and patient understanding is vital to maintaining the therapeutic relationship. Self-disclosure or examples are used often and cautiously to illustrate how the disorder functions. Alcoholics Anonymous employs self-disclosure as large part of its program to effectively illustrate common purpose and belonging among its members.

III. CONCLUSIONS

The intent of this paper has been to unify two different schools of thought for treating addictions through the common denominator of behavioural neuroscience. This provides a common link between the counselling profession and self-help groups, demonstrating their compatibility and potential synergy. For the counselor, understanding the truth of the first part of AAs Step One, 'We admitted we were powerless over alcohol', gives further support to this community program and enriches the knowledge base for employing the various counseling modalities. Seen through the lens of behavioural neuroscience, counselors can interpret client issues in a precise fashion without being impeded by trying to interpret what is believable or not. What we term neurocounselling has definitive aims and processes:

1. Understand the robust neurobiology and associated behaviour patterns of Step One;
2. Through self-disclosure, examples or reflections of client experience, facilitate the client's understanding of this disorder and its outcomes within their own lives;
3. With addiction framed as a neurobiological disorder developed over a lifetime, facilitate client understanding of the actual effect of circumstances and resultant behaviours over their lifetime to date (Dodge K. et al, 2009); and,
4. Support recovery of the prefrontal cortex and promote neuroadaptation of the reward system to healthy through positive life changes and experiences.

With the evidence presented herein, we are hopeful that this serves as evidence to unify approaches to care for persons with addiction through a new and more effective paradigm of neurocounselling. Our Step One Therapy approach encompasses a fundamental element of the Alcoholics Anonymous program, as well as its frequent use of self disclosure. Best practices in counseling modalities are fundamental to be able to maintain a positive therapeutic process and teach the underlying neuroscience. Understanding neuroplasticity gives tremendous hope to persons seeking recovery in that healthy functioning, unencumbered by addiction, can be the result over time. With reducing physiologic stress and allostatic load being a concurrent goal of therapy, counselors now have a far more focused approach than sifting through a constellation of issues.

Part 2 of this series will address the second part of AAs Step One: that our lives have become unmanageable. As written in 1939, this suggests that addiction is a progressive disorder to reach a point where life becomes unmanageable by the addicted person. Through combining neurobiology, genetics and environmental stresses we argue that addiction is a progressive disorder that occurs over a lifetime. Fundamentally, we see addiction as a progressive disorder of decision making. Seemingly rational choices made, often well before becoming addicted, set the life circumstances and internal conflicts necessary to cause disorder progression. Decisions become increasingly unhealthy as do life circumstances to the point where addiction fully dominates brain function and affected persons are unable to see a path out of it.

While we admit that we have given a very brief summary of very robust neuroscience, exploration of the aggregate references herein supports proof of concept. Ultimately proof of concept exists for the vast majority of clients who have engaged in our program. Fortunately, in the five years of refining this method clinically, we are now able to fully expect that most people we see will experience sustained recovery and reclamation of lives that were once on the precipice of tragedy.

ACKNOWLEDGEMENTS

We would like to thank the anonymous peer reviewers who provided many insights to improve the message and quality of this paper. We additionally thank Liam Carrie, MD, who assisted in research into the science and application of this approach, as well as reviewing this writing.

REFERENCES RÉFÉRENCES REFERENCIAS

1. A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic Dopamine System. *Neuron* 101.
2. Alcoholics Anonymous World Services (2001). Alcoholics Anonymous Fourth Edition (The Big Book). New York City.
3. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). <https://doi.org/10.1176/appi.books.9780890425596>.
4. American Psychiatric Association. What is Substance Use Disorder. Available at: Psychiatry.org - What Is a Substance Use Disorder?
5. Ashton, C. and Duffie, D. (2014). Comprehensive Literature of Best Practises in Residential Care. Government of New Brunswick.
6. Ashton, C. and Duffie, D. (2015). A Clinical Pathway through Substance Use Disorder, First Episode Psychosis to Post Traumatic Stress Disorder on the Basis of HPA Axis Dysfunction. *Journal of Diabetes, Metabolic Disorders & Control*, Vol 2.
7. Ashton, C. and Duffie, D. (2022). Crossed wires: the Hall effect in substance use disorder. *MOJ Addiction Medicine & Therapy*.
8. Authement, M. et al (2018). A role for corticotropin-releasing factor signaling in the lateral habenula and its modulation by early-life stress. *Science Signaling*, Vol. 11.
9. Borsook, B. et al (2016). Reward deficiency and anti-reward in pain chronification. *Neuroscience & Biobehavioral Reviews*, Vol 68.
10. Breese, G. et al (2005). Conceptual framework for the etiology of alcoholism: a "kindling"/stress hypothesis. *Psychopharmacology*, Vol 178
11. Canadian Counselling and Psychotherapy Association (CCPA). The Profession and Regulation. Available at: The Profession & Regulation - Canadian Counselling and Psychotherapy Association.
12. Carelli, R. and West, E. (2014). When a good taste turns bad: Neural mechanisms underlying the emergence of negative affect and associated natural reward devaluation by cocaine. *Neuropharmacology*, Vol 76
13. Castro, D. et al (2015). Lateral hypothalamus, nucleus accumbens, and ventral pallidum roles in eating and hunger: interactions between homeostatic and reward circuitry. *Frontiers in Systems Neuroscience*, Vol 9
14. Catani, M. et al (2013). A revised limbic system model for memory, emotion and behaviour. *Neuroscience and Biobehavioral Reviews* 37
15. Ceceli, A. et al (2023). Altered prefrontal signaling during inhibitory control in a salient drug context in human cocaine addiction. *Cerebral Cortex*, Vol 33
16. Chang, R. et al (2022). Deep Brain Stimulation in Drug Addiction Treatment: Research Progress and Perspective. *Frontiers in Psychiatry*, Sec. *Neuroimaging*, Vol 13
17. Cho, S. et al (2021). Synaptic transistors with human brain-like fJ energy consumption via double oxide semiconductor engineering for neuromorphic electronics. *Journal of Materials Chemistry*
18. Chin, A. et al (2019). Technologies Toward Three-Dimensional Brain-Mimicking IC Architecture. 2019 *Electron Devices Technology and Manufacturing Conference (EDTM)*
19. Close, C., Frederick, D., Newell, J. (2002). Modeling and Analysis of Dynamic Systems, Third Edition. John Wiley and Sons, New York
20. Corkrum, M. et al (2020). Dopamine-Evoked Synaptic Regulation in the Nucleus Accumbens Requires Astrocyte Activity. *Neuron*, Vol 105
21. Danjo et al. (2014). Aversive behavior induced by optogenetic inactivation of ventral tegmental area dopamine neurons is mediated by dopamine D2 receptors in the nucleus accumbens. *PNAS* Vol 111.
22. Day, J. and Carelli, R. (2007). The Nucleus Accumbens and Pavlovian Reward Learning. *Neuroscientist* 13(2).
23. Dixon, M. et al (2017). Emotion and the prefrontal cortex: An integrative review. *APAPsyNet*.
24. Dodge K. et al (2009). A dynamic cascade model of the development of substance-use onset. *Monogr Soc Res Child Dev*, Vol 74.
25. Febo, M. et al (2017). Dopamine homeostasis: brain functional connectivity in reward deficiency syndrome. *Frontiers in Bioscience*, Vol 22.
26. Feldman Barrett, L. (2006). Valence is a basic building block of emotional life. *Journal of Research in Personality*, Vol 40.
27. Feltenstein, M. et al (2021). Neural Substrates and Circuits of Drug Addiction. *Cold Spring Harb Perspect Med.*, Vol 11.
28. Floresco, S. (2015) The Nucleus Accumbens: An Interface Between Cognition, Emotion, and Action. *Annu. Rev. Psychol.* 66:25.
29. Fordham, B. et al (2021). The evidence for cognitive behavioural therapy in any condition, population or context: a meta-review of systematic reviews and panoramic meta-analysis. *Psychological Medicine* 51.

30. George, S. and Zane, B. (2016). Affect, emotion, and decision making. *Organizational Behavior and Human Decision Processes*, Vol 136.

31. Goldstein, R. and Volkow, N. (2011). Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*.

32. Graziene, N. et al (2018). A Focus on Reward Prediction and the Lateral Habenula: Functional Alterations and the Behavioral Outcomes Induced by Drugs of Abuse. *Frontiers in Synaptic Neuroscience*, Vol 10.

33. Harrison, R. (2008). The Design of Integrated Circuits to Observe Brain Activity. *Proceedings of the IEEE*, Vol 96.

34. Heinz, A. et al (2019). Addiction as Learned Behavior Patterns. *Journal of Clinical Medicine*, 8(8)

35. Iles, J. (2005). Simple models of stimulation of neurones in the brain by electric fields. *Progress in Biophysics and Molecular Biology* 87.

36. Innocenti, G. (2022). Chapter 1 - Defining neuroplasticity. *Handbook of Clinical Neurology*, Vol 184.

37. Jhou, T. et al. (2009). The Rostromedial Tegmental Nucleus (RMTg), a GABAergic Afferent to Midbrain Dopamine Neurons, Encodes Aversive Stimuli and Inhibits Motor Responses. *Neuron* 61.

38. Kalivas, P. and Volkow, N. (2005). The Neural Basis of Addiction: A Pathology of Motivation and Choice. *American Journal of Psychiatry*, Vol 162

39. Keiflin, R. and Janak, P. (2015). Dopamine Prediction Errors in Reward Learning and Addiction: From Theory to Neural Circuitry. *Neuron* 88

40. Koob, G. (1992). Dopamine, addiction and reward. *Seminars in Neuroscience*, Vol 4, Issue 2

41. Koob, G. (2015), The dark side of emotion: the addiction perspective. *Eur J Pharmacol.* Vol 753

42. Koob, G. and Le Moal, M. (2001). Drug Addiction, Dysregulation of Reward, and Allostasis. *Neuropsychopharmacology*, Vol 24.

43. Koob, G. and Le Moal, M. (2008). Addiction and the Brain Antireward System. *Annual Review of Psychology*, Vol 59.

44. Koob, G. and Mason, M. (2016) Existing and Future Drugs for the Treatment of the Dark Side of Addiction. *Annu. Rev. Pharmacol. Toxicol.* Vol 2.

45. Koob, G. and Schulkin, J. (2019). Addiction and stress: An allostatic view. *Neuroscience & Biobehavioral Reviews*, Vol 106.

46. Koob, G. and Schulkin, J. (2019). Addiction and stress: An allostatic view. *Neuroscience & Biobehavioral Reviews*, Vol 106.

47. Kwako, L. et al (2016). Addictions Neuroclinical Assessment: A Neuroscience-Based Framework for Addictive Disorders. *Biological Psychiatry*, Vol 80

48. Le Berre et al. (2019). Emotional processing and social cognition in alcohol use disorder. *Neuropsychology*, 33(6).

49. Liu et al. (2008). Enhanced selective memory consolidation following post-learning pleasant and aversive arousal. *Neurobiology of Learning and Memory*, Vol 89

50. MacGregor, R. (1987). Neural and Brain Modeling. Academic Press Inc. San Diego.

51. Maillard, A. et al (2020). Neuropsychological deficits in alcohol use disorder: impact on treatment. *Cognition and Addiction*, Ch. 8, Elsevier.

52. Matsumoto, M. and Hikosaka, O. (2007). Lateral habenula as a source of negative reward signals in dopamine neurons. *Nature* vol 447.

53. Mazzocchi, F. (2012). Complexity and the reductionism–holism debate in systems biology. *WIREs Syst Biol Med*.

54. Menegas et al., (2018). Dopamine neurons projecting to the posterior striatum reinforce avoidance of threatening stimuli. *Nature Neuroscience*, 21(10).

55. Miyachi, S. et al, (2006). Somatotopically arranged inputs from putamen and subthalamic nucleus to primary motor cortex. *Neuroscience Research*, Vol 56.

56. Mogenson, G., Jones, D. and Yim, C. (1980). From motivation to action: functional interface between the limbic system and the motor system. *Progress in neurobiology*.

57. Pitel, A. et al (2023). Impaired Global Precedence Effect in Severe Alcohol Use Disorder and Korsakoff's Syndrome: A Pilot Exploration through a Global/Local Visual Paradigm. *J. Clin. Med.* 12(11).

58. PROJECT MATCH RESEARCH GROUP (1998). Matching patients with alcohol disorders to treatments: Clinical implications from Project MATCH. *Journal of Mental Health*, Vol 7.

59. Pross A. (2012). What is life? How chemistry becomes biology. London, UK: Oxford University Press.

60. Root, D. et al (2015). The ventral pallidum: Subregion-specific functional anatomy and roles in motivated behaviors. *Progress in Neurobiology*, 130

61. Salgado, S. and Kaplitt, M. (2015). The Nucleus Accumbens: A Comprehensive Review. *Stereotactic and Functional Neurosurgery* 93.

62. Sawada, M. et al (2015). Function of the nucleus accumbens in motor control during recovery after spinal cord injury. *Science*, Vol 350.

63. Sequeira, M. and Gourley, S. (2021). The Stressed Orbitofrontal Cortex. *Behavioral Neuroscience*, Vol 135.

64. Singh, J. and Kapur, G. (2019). Study of Adaptive CMOS Circuits Extracting Feature of Neuroplasticity in Human Brain. *2019 International Conference on Signalling*.



65. Sulzer, D. (2011). How Addictive Drugs Disrupt Presynaptic Dopamine Neurotransmission. *Neuron* 69.
66. Suzuki, Y. and Tanaka, S. (2021). Functions of the ventromedial prefrontal cortex in emotion regulation under stress. *Nature Scientific Reports* vol 11.
67. Turner, J. (2019). Homeostasis as a fundamental principle for a coherent theory of brains. *Phil. Trans. R. Soc. B* 374.
68. Turrigiano, G. and Nelson, S. (2000). Hebb and homeostasis in neuronal plasticity. *Current Opinion in Neurobiology*, Vol 10.
69. Verdejo-Garcia, A. et al (2019). A Roadmap for Integrating Neuroscience Into Addiction Treatment: A Consensus of the Neuroscience Interest Group of the International Society of Addiction Medicine. *Frontiers in Psychiatry*, Vol 10.
70. Volkow, N., Michaelides, M. and Baler, R. (2019). The Neuroscience of Drug Reward and Addiction. *Physiological Reviews*, Vol 99, Issue 4.
71. Walters, G. (2002). Lessons Learned From Project MATCH. *Addictive Disorders & Their Treatment*, 1(4)
72. Wemm, S. and Sinha, R. (2019). Drug-induced stress responses and addiction risk and relapse. *Neurobiology of Stress*, Vol 10.
73. Wise, R. (2004). Dopamine, learning and motivation. *Nature Reviews Neuroscience*, vol 5.
74. Wise, R. and Robbie, M. (2020). Dopamine and Addiction. *Annual Review of Psychology*, Vol 71.
75. Wojdala, A., Molins, F. and Serrano, M. (2020). Stress and drug addiction: an up-to-date perspective from 2020. *Adicciones*, Vol 32, No 4.
76. Yang, H. et al. (2018). Nucleus Accumbens Subnuclei Regulate Motivated Behaviour via Direct Inhibition and Disinhibition of VTA Dopamine Subpopulations. *Neuron* 97.


GLOBAL JOURNALS GUIDELINES HANDBOOK 2025

WWW.GLOBALJOURNALS.ORG

MEMBERSHIPS

FELLOWS/ASSOCIATES OF MEDICAL RESEARCH COUNCIL

FMRC/AMRC MEMBERSHIPS



INTRODUCTION

FMRC/AMRC is the most prestigious membership of Global Journals accredited by Open Association of Research Society, U.S.A (OARS). The credentials of Fellow and Associate designations signify that the researcher has gained the knowledge of the fundamental and high-level concepts, and is a subject matter expert, proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice. The credentials are designated only to the researchers, scientists, and professionals that have been selected by a rigorous process by our Editorial Board and Management Board.

Associates of FMRC/AMRC are scientists and researchers from around the world are working on projects/researches that have huge potentials. Members support Global Journals' mission to advance technology for humanity and the profession.

FMRC

FELLOW OF MEDICAL RESEARCH COUNCIL

FELLOW OF MEDICAL RESEARCH COUNCIL is the most prestigious membership of Global Journals. It is an award and membership granted to individuals that the Open Association of Research Society judges to have made a 'substantial contribution to the improvement of computer science, technology, and electronics engineering.

The primary objective is to recognize the leaders in research and scientific fields of the current era with a global perspective and to create a channel between them and other researchers for better exposure and knowledge sharing. Members are most eminent scientists, engineers, and technologists from all across the world. Fellows are elected for life through a peer review process on the basis of excellence in the respective domain. There is no limit on the number of new nominations made in any year. Each year, the Open Association of Research Society elect up to 12 new Fellow Members.



BENEFITS

TO THE INSTITUTION

GET LETTER OF APPRECIATION

Global Journals sends a letter of appreciation of author to the Dean or CEO of the University or Company of which author is a part, signed by editor in chief or chief author.



EXCLUSIVE NETWORK

GET ACCESS TO A CLOSED NETWORK

A FMRC member gets access to a closed network of Tier 1 researchers and scientists with direct communication channel through our website. Fellows can reach out to other members or researchers directly. They should also be open to reaching out by other.

Career

Credibility

Exclusive

Reputation



CERTIFICATE

CERTIFICATE, LOR AND LASER-MOMENTO

Fellows receive a printed copy of a certificate signed by our Chief Author that may be used for academic purposes and a personal recommendation letter to the dean of member's university.

Career

Credibility

Exclusive

Reputation



DESIGNATION

GET HONORED TITLE OF MEMBERSHIP

Fellows can use the honored title of membership. The "FMRC" is an honored title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FMRC or William Walldroff, M.S., FMRC.

Career

Credibility

Exclusive

Reputation

RECOGNITION ON THE PLATFORM

BETTER VISIBILITY AND CITATION

All the Fellow members of FMRC get a badge of "Leading Member of Global Journals" on the Research Community that distinguishes them from others. Additionally, the profile is also partially maintained by our team for better visibility and citation. All fellows get a dedicated page on the website with their biography.

Career

Credibility

Reputation



FUTURE WORK

GET DISCOUNTS ON THE FUTURE PUBLICATIONS

Fellows receive discounts on the future publications with Global Journals up to 60%. Through our recommendation programs, members also receive discounts on publications made with OARS affiliated organizations.

Career

Financial



GJ INTERNAL ACCOUNT UNLIMITED FORWARD OF EMAILS

Fellows get secure and fast GJ work emails with unlimited storage of emails that they may use them as their primary email. For example, john [AT] globaljournals [DOT] org.

Career

Credibility

Reputation



PREMIUM TOOLS

ACCESS TO ALL THE PREMIUM TOOLS

To take future researches to the zenith, fellows receive access to all the premium tools that Global Journals have to offer along with the partnership with some of the best marketing leading tools out there.

Financial

CONFERENCES & EVENTS

ORGANIZE SEMINAR/CONFERENCE

Fellows are authorized to organize symposium/seminar/conference on behalf of Global Journal Incorporation (USA). They can also participate in the same organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent. Additionally, they get free research conferences (and others) alerts.

Career

Credibility

Financial

EARLY INVITATIONS

EARLY INVITATIONS TO ALL THE SYMPOSIUMS, SEMINARS, CONFERENCES

All fellows receive the early invitations to all the symposiums, seminars, conferences and webinars hosted by Global Journals in their subject.

Exclusive





PUBLISHING ARTICLES & BOOKS

EARN 60% OF SALES PROCEEDS

Fellows can publish articles (limited) without any fees. Also, they can earn up to 70% of sales proceeds from the sale of reference/review books/literature/publishing of research paper. The FMRC member can decide its price and we can help in making the right decision.

Exclusive

Financial

REVIEWERS

GET A REMUNERATION OF 15% OF AUTHOR FEES

Fellow members are eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get a remuneration of 15% of author fees, taken from the author of a respective paper.

Financial

ACCESS TO EDITORIAL BOARD

BECOME A MEMBER OF THE EDITORIAL BOARD

Fellows and Associates may join as a member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer.

Career

Credibility

Exclusive

Reputation

AND MUCH MORE

GET ACCESS TO SCIENTIFIC MUSEUMS AND OBSERVATORIES ACROSS THE GLOBE

All members get access to 5 selected scientific museums and observatories across the globe. All researches published with Global Journals will be kept under deep archival facilities across regions for future protections and disaster recovery. They get 10 GB free secure cloud access for storing research files.



ASSOCIATE OF MEDICAL RESEARCH COUNCIL

ASSOCIATE OF MEDICAL RESEARCH COUNCIL is the membership of Global Journals awarded to individuals that the Open Association of Research Society judges to have made a 'substantial contribution to the improvement of computer science, technology, and electronics engineering.

The primary objective is to recognize the leaders in research and scientific fields of the current era with a global perspective and to create a channel between them and other researchers for better exposure and knowledge sharing. Members are most eminent scientists, engineers, and technologists from all across the world. Associate membership can later be promoted to Fellow Membership. Associates are elected for life through a peer review process on the basis of excellence in the respective domain. There is no limit on the number of new nominations made in any year. Each year, the Open Association of Research Society elect up to 12 new Associate Members.



BENEFITS

TO THE INSTITUTION

GET LETTER OF APPRECIATION

Global Journals sends a letter of appreciation of author to the Dean or CEO of the University or Company of which author is a part, signed by editor in chief or chief author.



EXCLUSIVE NETWORK

GET ACCESS TO A CLOSED NETWORK

A AMRC member gets access to a closed network of Tier 2 researchers and scientists with direct communication channel through our website. Associates can reach out to other members or researchers directly. They should also be open to reaching out by other.

Career

Credibility

Exclusive

Reputation



CERTIFICATE

CERTIFICATE, LOR AND LASER-MOMENTO

Associates receive a printed copy of a certificate signed by our Chief Author that may be used for academic purposes and a personal recommendation letter to the dean of member's university.

Career

Credibility

Exclusive

Reputation



DESIGNATION

GET HONORED TITLE OF MEMBERSHIP

Associates can use the honored title of membership. The "AMRC" is an honored title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., AMRC or William Walldroff, M.S., AMRC.

Career

Credibility

Exclusive

Reputation

RECOGNITION ON THE PLATFORM

BETTER VISIBILITY AND CITATION

All the Associate members of AMRC get a badge of "Leading Member of Global Journals" on the Research Community that distinguishes them from others. Additionally, the profile is also partially maintained by our team for better visibility and citation.

Career

Credibility

Reputation



FUTURE WORK

GET DISCOUNTS ON THE FUTURE PUBLICATIONS

Associates receive discounts on future publications with Global Journals up to 30%. Through our recommendation programs, members also receive discounts on publications made with OARS affiliated organizations.

Career

Financial



GJ ACCOUNT

UNLIMITED FORWARD OF EMAILS

Associates get secure and fast GJ work emails with 5GB forward of emails that they may use them as their primary email. For example, john [AT] globaljournals [DOT] org.

Career

Credibility

Reputation



PREMIUM TOOLS

ACCESS TO ALL THE PREMIUM TOOLS

To take future researches to the zenith, fellows receive access to almost all the premium tools that Global Journals have to offer along with the partnership with some of the best marketing leading tools out there.

Financial

CONFERENCES & EVENTS

ORGANIZE SEMINAR/CONFERENCE

Associates are authorized to organize symposium/seminar/conference on behalf of Global Journal Incorporation (USA). They can also participate in the same organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent. Additionally, they get free research conferences (and others) alerts.

Career

Credibility

Financial

EARLY INVITATIONS

EARLY INVITATIONS TO ALL THE SYMPOSIUMS, SEMINARS, CONFERENCES

All associates receive the early invitations to all the symposiums, seminars, conferences and webinars hosted by Global Journals in their subject.

Exclusive





PUBLISHING ARTICLES & BOOKS

EARN 60% OF SALES PROCEEDS

Associates can publish articles (limited) without any fees. Also, they can earn up to 30-40% of sales proceeds from the sale of reference/review books/literature/publishing of research paper

Exclusive

Financial

REVIEWERS

GET A REMUNERATION OF 15% OF AUTHOR FEES

Associate members are eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get a remuneration of 15% of author fees, taken from the author of a respective paper.

Financial

AND MUCH MORE

GET ACCESS TO SCIENTIFIC MUSEUMS AND OBSERVATORIES ACROSS THE GLOBE

All members get access to 2 selected scientific museums and observatories across the globe. All researches published with Global Journals will be kept under deep archival facilities across regions for future protections and disaster recovery. They get 5 GB free secure cloud access for storing research files.



ASSOCIATE	FELLOW	RESEARCH GROUP	BASIC
<p>\$4800 lifetime designation</p> <p>Certificate, LoR and Momento 2 discounted publishing/year</p> <p>Gradation of Research 10 research contacts/day</p> <p>1 GB Cloud Storage</p> <p>GJ Community Access</p>	<p>\$6800 lifetime designation</p> <p>Certificate, LoR and Momento</p> <p>Unlimited discounted publishing/year</p> <p>Gradation of Research Unlimited research contacts/day</p> <p>5 GB Cloud Storage</p> <p>Online Presense Assistance</p> <p>GJ Community Access</p>	<p>\$12500.00 organizational</p> <p>Certificates, LoRs and Momentos</p> <p>Unlimited free publishing/year</p> <p>Gradation of Research Unlimited research contacts/day</p> <p>Unlimited Cloud Storage</p> <p>Online Presense Assistance</p> <p>GJ Community Access</p>	<p>APC per article</p> <p>GJ Community Access</p>

PREFERRED AUTHOR GUIDELINES

We accept the manuscript submissions in any standard (generic) format.

We typeset manuscripts using advanced typesetting tools like Adobe In Design, CorelDraw, TeXnicCenter, and TeXStudio. We usually recommend authors submit their research using any standard format they are comfortable with, and let Global Journals do the rest.

Alternatively, you can download our basic template from <https://globaljournals.org/Template>

Authors should submit their complete paper/article, including text illustrations, graphics, conclusions, artwork, and tables. Authors who are not able to submit manuscript using the form above can email the manuscript department at submit@globaljournals.org or get in touch with chiefeditor@globaljournals.org if they wish to send the abstract before submission.

BEFORE AND DURING SUBMISSION

Authors must ensure the information provided during the submission of a paper is authentic. Please go through the following checklist before submitting:

1. Authors must go through the complete author guideline and understand and *agree to Global Journals' ethics and code of conduct*, along with author responsibilities.
2. Authors must accept the privacy policy, terms, and conditions of Global Journals.
3. Ensure corresponding author's email address and postal address are accurate and reachable.
4. Manuscript to be submitted must include keywords, an abstract, a paper title, co-author(s') names and details (email address, name, phone number, and institution), figures and illustrations in vector format including appropriate captions, tables, including titles and footnotes, a conclusion, results, acknowledgments and references.
5. Authors should submit paper in a ZIP archive if any supplementary files are required along with the paper.
6. Proper permissions must be acquired for the use of any copyrighted material.
7. Manuscript submitted *must not have been submitted or published elsewhere* and all authors must be aware of the submission.

Declaration of Conflicts of Interest

It is required for authors to declare all financial, institutional, and personal relationships with other individuals and organizations that could influence (bias) their research.

POLICY ON PLAGIARISM

Plagiarism is not acceptable in Global Journals submissions at all.

Plagiarized content will not be considered for publication. We reserve the right to inform authors' institutions about plagiarism detected either before or after publication. If plagiarism is identified, we will follow COPE guidelines:

Authors are solely responsible for all the plagiarism that is found. The author must not fabricate, falsify or plagiarize existing research data. The following, if copied, will be considered plagiarism:

- Words (language)
- Ideas
- Findings
- Writings
- Diagrams
- Graphs
- Illustrations
- Lectures



- Printed material
- Graphic representations
- Computer programs
- Electronic material
- Any other original work

AUTHORSHIP POLICIES

Global Journals follows the definition of authorship set up by the Open Association of Research Society, USA. According to its guidelines, authorship criteria must be based on:

1. Substantial contributions to the conception and acquisition of data, analysis, and interpretation of findings.
2. Drafting the paper and revising it critically regarding important academic content.
3. Final approval of the version of the paper to be published.

Changes in Authorship

The corresponding author should mention the name and complete details of all co-authors during submission and in manuscript. We support addition, rearrangement, manipulation, and deletions in authors list till the early view publication of the journal. We expect that corresponding author will notify all co-authors of submission. We follow COPE guidelines for changes in authorship.

Copyright

During submission of the manuscript, the author is confirming an exclusive license agreement with Global Journals which gives Global Journals the authority to reproduce, reuse, and republish authors' research. We also believe in flexible copyright terms where copyright may remain with authors/employers/institutions as well. Contact your editor after acceptance to choose your copyright policy. You may follow this form for copyright transfers.

Appealing Decisions

Unless specified in the notification, the Editorial Board's decision on publication of the paper is final and cannot be appealed before making the major change in the manuscript.

Acknowledgments

Contributors to the research other than authors credited should be mentioned in Acknowledgments. The source of funding for the research can be included. Suppliers of resources may be mentioned along with their addresses.

Declaration of funding sources

Global Journals is in partnership with various universities, laboratories, and other institutions worldwide in the research domain. Authors are requested to disclose their source of funding during every stage of their research, such as making analysis, performing laboratory operations, computing data, and using institutional resources, from writing an article to its submission. This will also help authors to get reimbursements by requesting an open access publication letter from Global Journals and submitting to the respective funding source.

PREPARING YOUR MANUSCRIPT

Authors can submit papers and articles in an acceptable file format: MS Word (doc, docx), LaTeX (.tex, .zip or .rar including all of your files), Adobe PDF (.pdf), rich text format (.rtf), simple text document (.txt), Open Document Text (.odt), and Apple Pages (.pages). Our professional layout editors will format the entire paper according to our official guidelines. This is one of the highlights of publishing with Global Journals—authors should not be concerned about the formatting of their paper. Global Journals accepts articles and manuscripts in every major language, be it Spanish, Chinese, Japanese, Portuguese, Russian, French, German, Dutch, Italian, Greek, or any other national language, but the title, subtitle, and abstract should be in English. This will facilitate indexing and the pre-peer review process.

The following is the official style and template developed for publication of a research paper. Authors are not required to follow this style during the submission of the paper. It is just for reference purposes.



Manuscript Style Instruction (Optional)

- Microsoft Word Document Setting Instructions.
- Font type of all text should be Swis721 Lt BT.
- Page size: 8.27" x 11", left margin: 0.65, right margin: 0.65, bottom margin: 0.75.
- Paper title should be in one column of font size 24.
- Author name in font size of 11 in one column.
- Abstract: font size 9 with the word "Abstract" in bold italics.
- Main text: font size 10 with two justified columns.
- Two columns with equal column width of 3.38 and spacing of 0.2.
- First character must be three lines drop-capped.
- The paragraph before spacing of 1 pt and after of 0 pt.
- Line spacing of 1 pt.
- Large images must be in one column.
- The names of first main headings (Heading 1) must be in Roman font, capital letters, and font size of 10.
- The names of second main headings (Heading 2) must not include numbers and must be in italics with a font size of 10.

Structure and Format of Manuscript

The recommended size of an original research paper is under 15,000 words and review papers under 7,000 words. Research articles should be less than 10,000 words. Research papers are usually longer than review papers. Review papers are reports of significant research (typically less than 7,000 words, including tables, figures, and references)

A research paper must include:

- a) A title which should be relevant to the theme of the paper.
- b) A summary, known as an abstract (less than 150 words), containing the major results and conclusions.
- c) Up to 10 keywords that precisely identify the paper's subject, purpose, and focus.
- d) An introduction, giving fundamental background objectives.
- e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition, sources of information must be given, and numerical methods must be specified by reference.
- f) Results which should be presented concisely by well-designed tables and figures.
- g) Suitable statistical data should also be given.
- h) All data must have been gathered with attention to numerical detail in the planning stage.

Design has been recognized to be essential to experiments for a considerable time, and the editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned unrefereed.

- i) Discussion should cover implications and consequences and not just recapitulate the results; conclusions should also be summarized.
- j) There should be brief acknowledgments.
- k) There ought to be references in the conventional format. Global Journals recommends APA format.

Authors should carefully consider the preparation of papers to ensure that they communicate effectively. Papers are much more likely to be accepted if they are carefully designed and laid out, contain few or no errors, are summarizing, and follow instructions. They will also be published with much fewer delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and suggestions to improve brevity.



FORMAT STRUCTURE

It is necessary that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

All manuscripts submitted to Global Journals should include:

Title

The title page must carry an informative title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) where the work was carried out.

Author details

The full postal address of any related author(s) must be specified.

Abstract

The abstract is the foundation of the research paper. It should be clear and concise and must contain the objective of the paper and inferences drawn. It is advised to not include big mathematical equations or complicated jargon.

Many researchers searching for information online will use search engines such as Google, Yahoo or others. By optimizing your paper for search engines, you will amplify the chance of someone finding it. In turn, this will make it more likely to be viewed and cited in further works. Global Journals has compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Keywords

A major lynchpin of research work for the writing of research papers is the keyword search, which one will employ to find both library and internet resources. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining, and indexing.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy: planning of a list of possible keywords and phrases to try.

Choice of the main keywords is the first tool of writing a research paper. Research paper writing is an art. Keyword search should be as strategic as possible.

One should start brainstorming lists of potential keywords before even beginning searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in a research paper?" Then consider synonyms for the important words.

It may take the discovery of only one important paper to steer in the right keyword direction because, in most databases, the keywords under which a research paper is abstracted are listed with the paper.

Numerical Methods

Numerical methods used should be transparent and, where appropriate, supported by references.

Abbreviations

Authors must list all the abbreviations used in the paper at the end of the paper or in a separate table before using them.

Formulas and equations

Authors are advised to submit any mathematical equation using either MathJax, KaTeX, or LaTeX, or in a very high-quality image.

Tables, Figures, and Figure Legends

Tables: Tables should be cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g., Table 4, a self-explanatory caption, and be on a separate sheet. Authors must submit tables in an editable format and not as images. References to these tables (if any) must be mentioned accurately.



Figures

Figures are supposed to be submitted as separate files. Always include a citation in the text for each figure using Arabic numbers, e.g., Fig. 4. Artwork must be submitted online in vector electronic form or by emailing it.

PREPARATION OF ELECTRONIC FIGURES FOR PUBLICATION

Although low-quality images are sufficient for review purposes, print publication requires high-quality images to prevent the final product being blurred or fuzzy. Submit (possibly by e-mail) EPS (line art) or TIFF (halftone/ photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Avoid using pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings). Please give the data for figures in black and white or submit a Color Work Agreement form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution at final image size ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs): >350 dpi; figures containing both halftone and line images: >650 dpi.

Color charges: Authors are advised to pay the full cost for the reproduction of their color artwork. Hence, please note that if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a Color Work Agreement form before your paper can be published. Also, you can email your editor to remove the color fee after acceptance of the paper.

TIPS FOR WRITING A GOOD QUALITY MEDICAL RESEARCH PAPER

1. Choosing the topic: In most cases, the topic is selected by the interests of the author, but it can also be suggested by the guides. You can have several topics, and then judge which you are most comfortable with. This may be done by asking several questions of yourself, like "Will I be able to carry out a search in this area? Will I find all necessary resources to accomplish the search? Will I be able to find all information in this field area?" If the answer to this type of question is "yes," then you ought to choose that topic. In most cases, you may have to conduct surveys and visit several places. Also, you might have to do a lot of work to find all the rises and falls of the various data on that subject. Sometimes, detailed information plays a vital role, instead of short information. Evaluators are human: The first thing to remember is that evaluators are also human beings. They are not only meant for rejecting a paper. They are here to evaluate your paper. So present your best aspect.

2. Think like evaluators: If you are in confusion or getting demotivated because your paper may not be accepted by the evaluators, then think, and try to evaluate your paper like an evaluator. Try to understand what an evaluator wants in your research paper, and you will automatically have your answer. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

3. Ask your guides: If you are having any difficulty with your research, then do not hesitate to share your difficulty with your guide (if you have one). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work, then ask your supervisor to help you with an alternative. He or she might also provide you with a list of essential readings.

4. Use of computer is recommended: As you are doing research in the field of medical research then this point is quite obvious. Use right software: Always use good quality software packages. If you are not capable of judging good software, then you can lose the quality of your paper unknowingly. There are various programs available to help you which you can get through the internet.

5. Use the internet for help: An excellent start for your paper is using Google. It is a wondrous search engine, where you can have your doubts resolved. You may also read some answers for the frequent question of how to write your research paper or find a model research paper. You can download books from the internet. If you have all the required books, place importance on reading, selecting, and analyzing the specified information. Then sketch out your research paper. Use big pictures: You may use encyclopedias like Wikipedia to get pictures with the best resolution. At Global Journals, you should strictly follow here.



6. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right? It is a good habit which helps to not lose your continuity. You should always use bookmarks while searching on the internet also, which will make your search easier.

7. Revise what you wrote: When you write anything, always read it, summarize it, and then finalize it.

8. Make every effort: Make every effort to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in the introduction—what is the need for a particular research paper. Polish your work with good writing skills and always give an evaluator what he wants. Make backups: When you are going to do any important thing like making a research paper, you should always have backup copies of it either on your computer or on paper. This protects you from losing any portion of your important data.

9. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several unnecessary diagrams will degrade the quality of your paper by creating a hodgepodge. So always try to include diagrams which were made by you to improve the readability of your paper. Use of direct quotes: When you do research relevant to literature, history, or current affairs, then use of quotes becomes essential, but if the study is relevant to science, use of quotes is not preferable.

10. Use proper verb tense: Use proper verb tenses in your paper. Use past tense to present those events that have happened. Use present tense to indicate events that are going on. Use future tense to indicate events that will happen in the future. Use of wrong tenses will confuse the evaluator. Avoid sentences that are incomplete.

11. Pick a good study spot: Always try to pick a spot for your research which is quiet. Not every spot is good for studying.

12. Know what you know: Always try to know what you know by making objectives, otherwise you will be confused and unable to achieve your target.

13. Use good grammar: Always use good grammar and words that will have a positive impact on the evaluator; use of good vocabulary does not mean using tough words which the evaluator has to find in a dictionary. Do not fragment sentences. Eliminate one-word sentences. Do not ever use a big word when a smaller one would suffice.

Verbs have to be in agreement with their subjects. In a research paper, do not start sentences with conjunctions or finish them with prepositions. When writing formally, it is advisable to never split an infinitive because someone will (wrongly) complain. Avoid clichés like a disease. Always shun irritating alliteration. Use language which is simple and straightforward. Put together a neat summary.

14. Arrangement of information: Each section of the main body should start with an opening sentence, and there should be a changeover at the end of the section. Give only valid and powerful arguments for your topic. You may also maintain your arguments with records.


15. Never start at the last minute: Always allow enough time for research work. Leaving everything to the last minute will degrade your paper and spoil your work.

16. Multitasking in research is not good: Doing several things at the same time is a bad habit in the case of research activity. Research is an area where everything has a particular time slot. Divide your research work into parts, and do a particular part in a particular time slot.

17. Never copy others' work: Never copy others' work and give it your name because if the evaluator has seen it anywhere, you will be in trouble. Take proper rest and food: No matter how many hours you spend on your research activity, if you are not taking care of your health, then all your efforts will have been in vain. For quality research, take proper rest and food.

18. Go to seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.

19. Refresh your mind after intervals: Try to give your mind a rest by listening to soft music or sleeping in intervals. This will also improve your memory. Acquire colleagues: Always try to acquire colleagues. No matter how sharp you are, if you acquire colleagues, they can give you ideas which will be helpful to your research.



20. Think technically: Always think technically. If anything happens, search for its reasons, benefits, and demerits. Think and then print: When you go to print your paper, check that tables are not split, headings are not detached from their descriptions, and page sequence is maintained.

21. Adding unnecessary information: Do not add unnecessary information like "I have used MS Excel to draw graphs." Irrelevant and inappropriate material is superfluous. Foreign terminology and phrases are not apropos. One should never take a broad view. Analogy is like feathers on a snake. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Never oversimplify: When adding material to your research paper, never go for oversimplification; this will definitely irritate the evaluator. Be specific. Never use rhythmic redundancies. Contractions shouldn't be used in a research paper. Comparisons are as terrible as clichés. Give up ampersands, abbreviations, and so on. Remove commas that are not necessary. Parenthetical words should be between brackets or commas. Understatement is always the best way to put forward earth-shaking thoughts. Give a detailed literary review.

22. Report concluded results: Use concluded results. From raw data, filter the results, and then conclude your studies based on measurements and observations taken. An appropriate number of decimal places should be used. Parenthetical remarks are prohibited here. Proofread carefully at the final stage. At the end, give an outline to your arguments. Spot perspectives of further study of the subject. Justify your conclusion at the bottom sufficiently, which will probably include examples.

23. Upon conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print for the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects of your research.

INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form which is presented in the guidelines using the template.
- Please note the criteria peer reviewers will use for grading the final paper.

Final points:

One purpose of organizing a research paper is to let people interpret your efforts selectively. The journal requires the following sections, submitted in the order listed, with each section starting on a new page:

The introduction: This will be compiled from reference material and reflect the design processes or outline of basis that directed you to make a study. As you carry out the process of study, the method and process section will be constructed like that. The results segment will show related statistics in nearly sequential order and direct reviewers to similar intellectual paths throughout the data that you gathered to carry out your study.

The discussion section:

This will provide understanding of the data and projections as to the implications of the results. The use of good quality references throughout the paper will give the effort trustworthiness by representing an alertness to prior workings.

Writing a research paper is not an easy job, no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record-keeping are the only means to make straightforward progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear: Adhere to recommended page limits.



Mistakes to avoid:

- Insertion of a title at the foot of a page with subsequent text on the next page.
- Separating a table, chart, or figure—confine each to a single page.
- Submitting a manuscript with pages out of sequence.
- In every section of your document, use standard writing style, including articles ("a" and "the").
- Keep paying attention to the topic of the paper.
- Use paragraphs to split each significant point (excluding the abstract).
- Align the primary line of each section.
- Present your points in sound order.
- Use present tense to report well-accepted matters.
- Use past tense to describe specific results.
- Do not use familiar wording; don't address the reviewer directly. Don't use slang or superlatives.
- Avoid use of extra pictures—include only those figures essential to presenting results.

Title page:

Choose a revealing title. It should be short and include the name(s) and address(es) of all authors. It should not have acronyms or abbreviations or exceed two printed lines.

Abstract: This summary should be two hundred words or less. It should clearly and briefly explain the key findings reported in the manuscript and must have precise statistics. It should not have acronyms or abbreviations. It should be logical in itself. Do not cite references at this point.

An abstract is a brief, distinct paragraph summary of finished work or work in development. In a minute or less, a reviewer can be taught the foundation behind the study, common approaches to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Use comprehensive sentences, and do not sacrifice readability for brevity; you can maintain it succinctly by phrasing sentences so that they provide more than a lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study with the subsequent elements in any summary. Try to limit the initial two items to no more than one line each.

Reason for writing the article—theory, overall issue, purpose.


- Fundamental goal.
- To-the-point depiction of the research.
- Consequences, including definite statistics—if the consequences are quantitative in nature, account for this; results of any numerical analysis should be reported. Significant conclusions or questions that emerge from the research.

Approach:

- Single section and succinct.
- An outline of the job done is always written in past tense.
- Concentrate on shortening results—limit background information to a verdict or two.
- Exact spelling, clarity of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else.

Introduction:

The introduction should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable of comprehending and calculating the purpose of your study without having to refer to other works. The basis for the study should be offered. Give the most important references, but avoid making a comprehensive appraisal of the topic. Describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will give no attention to your results. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here.



The following approach can create a valuable beginning:

- Explain the value (significance) of the study.
- Defend the model—why did you employ this particular system or method? What is its compensation? Remark upon its appropriateness from an abstract point of view as well as pointing out sensible reasons for using it.
- Present a justification. State your particular theory(-ies) or aim(s), and describe the logic that led you to choose them.
- Briefly explain the study's tentative purpose and how it meets the declared objectives.

Approach:

Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done. Sort out your thoughts; manufacture one key point for every section. If you make the four points listed above, you will need at least four paragraphs. Present surrounding information only when it is necessary to support a situation. The reviewer does not desire to read everything you know about a topic. Shape the theory specifically—do not take a broad view.

As always, give awareness to spelling, simplicity, and correctness of sentences and phrases.

Procedures (methods and materials):

This part is supposed to be the easiest to carve if you have good skills. A soundly written procedures segment allows a capable scientist to replicate your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order, but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt to give the least amount of information that would permit another capable scientist to replicate your outcome, but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section.

When a technique is used that has been well-described in another section, mention the specific item describing the way, but draw the basic principle while stating the situation. The purpose is to show all particular resources and broad procedures so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step-by-step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

Materials may be reported in part of a section or else they may be recognized along with your measures.

Methods:

- Report the method and not the particulars of each process that engaged the same methodology.
- Describe the method entirely.
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures.
- Simplify—detail how procedures were completed, not how they were performed on a particular day.
- If well-known procedures were used, account for the procedure by name, possibly with a reference, and that's all.

Approach:

It is embarrassing to use vigorous voice when documenting methods without using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result, when writing up the methods, most authors use third person passive voice.

Use standard style in this and every other part of the paper—avoid familiar lists, and use full sentences.

What to keep away from:

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings—save it for the argument.
- Leave out information that is immaterial to a third party.



Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part as entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Use statistics and tables, if suitable, to present consequences most efficiently.

You must clearly differentiate material which would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matters should not be submitted at all except if requested by the instructor.

Content:

- Sum up your conclusions in text and demonstrate them, if suitable, with figures and tables.
- In the manuscript, explain each of your consequences, and point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation of an exacting study.
- Explain results of control experiments and give remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or manuscript.

What to stay away from:

- Do not discuss or infer your outcome, report surrounding information, or try to explain anything.
- Do not include raw data or intermediate calculations in a research manuscript.
- Do not present similar data more than once.
- A manuscript should complement any figures or tables, not duplicate information.
- Never confuse figures with tables—there is a difference.

Approach:

As always, use past tense when you submit your results, and put the whole thing in a reasonable order.

Put figures and tables, appropriately numbered, in order at the end of the report.

If you desire, you may place your figures and tables properly within the text of your results section.

Figures and tables:

If you put figures and tables at the end of some details, make certain that they are visibly distinguished from any attached appendix materials, such as raw facts. Whatever the position, each table must be titled, numbered one after the other, and include a heading. All figures and tables must be divided from the text.

Discussion:

The discussion is expected to be the trickiest segment to write. A lot of papers submitted to the journal are discarded based on problems with the discussion. There is no rule for how long an argument should be.

Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implications of the study. The purpose here is to offer an understanding of your results and support all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of results should be fully described.

Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact, you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved the prospect, and let it drop at that. Make a decision as to whether each premise is supported or discarded or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."



Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work.

- You may propose future guidelines, such as how an experiment might be personalized to accomplish a new idea.
- Give details of all of your remarks as much as possible, focusing on mechanisms.
- Make a decision as to whether the tentative design sufficiently addressed the theory and whether or not it was correctly restricted. Try to present substitute explanations if they are sensible alternatives.
- One piece of research will not counter an overall question, so maintain the large picture in mind. Where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

When you refer to information, differentiate data generated by your own studies from other available information. Present work done by specific persons (including you) in past tense.

Describe generally acknowledged facts and main beliefs in present tense.

THE ADMINISTRATION RULES

Administration Rules to Be Strictly Followed before Submitting Your Research Paper to Global Journals Inc.

Please read the following rules and regulations carefully before submitting your research paper to Global Journals Inc. to avoid rejection.

Segment draft and final research paper: You have to strictly follow the template of a research paper, failing which your paper may get rejected. You are expected to write each part of the paper wholly on your own. The peer reviewers need to identify your own perspective of the concepts in your own terms. Please do not extract straight from any other source, and do not rephrase someone else's analysis. Do not allow anyone else to proofread your manuscript.

Written material: You may discuss this with your guides and key sources. Do not copy anyone else's paper, even if this is only imitation, otherwise it will be rejected on the grounds of plagiarism, which is illegal. Various methods to avoid plagiarism are strictly applied by us to every paper, and, if found guilty, you may be blacklisted, which could affect your career adversely. To guard yourself and others from possible illegal use, please do not permit anyone to use or even read your paper and file.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILED)
BY GLOBAL JOURNALS

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals.

Topics	Grades		
	A-B	C-D	E-F
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

INDEX

A

Acetylated · 4, 2, 4

B

Bradykinesia · 2

C

Chromatin · 2, 5

D

Disorientation · 2
Donepezil · 1

E

Epigenetic · 2, 5

G

Glutamate · 3, 4

H

Hallucinogenic · 2
Homeostasis · 3, 4

L

Lysosomal · 2, 3

N

Neurodegenerative · 1, 2, 4
Neutralization · 4

R

Randomized · 3
Rivastigmine · 1

S

Serotonergic · 3, 2, 3
Streptavidin · 2



save our planet



Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.MedicalResearchJournal.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals