

GLOBAL JOURNAL

OF MEDICAL RESEARCH: D

Radiology, Diagnostic Imaging and Instrumentation

Characterization of Ascitis

Epileptic Sudanese Children

Highlights

Diagnosis and Management

Neurosurgical and Radiologic

Discovering Thoughts, Inventing Future

VOLUME 13

ISSUE 3

VERSION 10



GLOBAL JOURNAL OF MEDICAL RESEARCH: D
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION

GLOBAL JOURNAL OF MEDICAL RESEARCH: D
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
VOLUME 13 ISSUE 3 (VER. 1.0)

OPEN ASSOCIATION OF RESEARCH SOCIETY

© Global Journal of Medical Research . 2013.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website <http://globaljournals.us/terms-and-condition/menu-id-1463/>

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)

Sponsors: *Open Association of Research Society*
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please
email us at local@globaljournals.org

eContacts

Press Inquiries: press@globaljournals.org
Investor Inquiries: investers@globaljournals.org
Technical Support: technology@globaljournals.org
Media & Releases: media@globaljournals.org

Pricing (Including by Air Parcel Charges):

For Authors:

22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

INTEGRATED EDITORIAL BOARD
(COMPUTER SCIENCE, ENGINEERING, MEDICAL, MANAGEMENT, NATURAL
SCIENCE, SOCIAL SCIENCE)

John A. Hamilton, "Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software
Engineering
Director, Information Assurance
Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at
Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D. and M.S. Syracuse University,
Syracuse, New York
M.S. and B.S. Bogazici University,
Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A. Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range
Nutritionist
Ph.D. Edinburgh University - Animal
Nutrition
M.S. Aberdeen University - Animal
Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing &
Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems
Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinipiac
BS, Jilin Institute of Technology; MA, MS,
PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley
& Istanbul University

Dr. Bart Lambrecht

Director of Research in Accounting and Finance
Professor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD
(Cambridge)

Dr. Carlos García Pont

Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management),
Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE,
University of Navarra
Degree in Industrial Engineering,
Universitat Politècnica de Catalunya

Dr. Fotini Labropulu

Mathematics - Luther College
University of Regina
Ph.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windsor

Dr. Lynn Lim

Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD,
FHEA

Dr. Mihaly Mezei

ASSOCIATE PROFESSOR
Department of Structural and Chemical
Biology, Mount Sinai School of Medical
Center
Ph.D., Eötvös Loránd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram

Department of Accounting and Finance
Lancaster University Management School
Ph.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño

Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso

Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and
Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.

Director, EP Laboratories, Philadelphia VA
Medical Center
Cardiovascular Medicine - Cardiac
Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng

MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular
Medicine
Davee Department of Neurology and Clinical
Neuroscience
Northwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli

Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic
Radiology
M.D., State University of New York at
Buffalo, School of Medicine and
Biomedical Sciences

Dr. Roberto Sanchez

Associate Professor
Department of Structural and Chemical
Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun

Professor of Earth and Atmospheric
SciencesPurdue University Director
National Center for Typhoon and
Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences,
National Central University, Chung-Li,
TaiwanUniversity Chair Professor
Institute of Environmental Engineering,
National Chiao Tung University, Hsin-
chu, Taiwan.Ph.D., MS The University of
Chicago, Geophysical Sciences
BS National Taiwan University,
Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick

M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and
Hypertension Division (PMC)
Penn Medicine, University of
Pennsylvania
Presbyterian Medical Center,
Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of
Internal Medicine

Dr. Bassey Benjamin Esu

B.Sc. Marketing; MBA Marketing; Ph.D
Marketing
Lecturer, Department of Marketing,
University of Calabar
Tourism Consultant, Cross River State
Tourism Development Department
Co-ordinator , Sustainable Tourism
Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.

IEEE Senior Member
Chairperson, Department of Computer
Science
AUST - American University of Science &
Technology
Alfred Naccash Avenue – Ashrafieh

PRESIDENT EDITOR (HON.)

Dr. George Perry, (Neuroscientist)

Dean and Professor, College of Sciences

Denham Harman Research Award (American Aging Association)

ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization

AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences

University of Texas at San Antonio

Postdoctoral Fellow (Department of Cell Biology)

Baylor College of Medicine

Houston, Texas, United States

CHIEF AUTHOR (HON.)

Dr. R.K. Dixit

M.Sc., Ph.D., FICCT

Chief Author, India

Email: authorind@computerresearch.org

DEAN & EDITOR-IN-CHIEF (HON.)

Vivek Dubey(HON.)

MS (Industrial Engineering),

MS (Mechanical Engineering)

University of Wisconsin, FICCT

Editor-in-Chief, USA

editorusa@computerresearch.org

Sangita Dixit

M.Sc., FICCT

Dean & Chancellor (Asia Pacific)

deanind@computerresearch.org

Suyash Dixit

(B.E., Computer Science Engineering), FICCTT

President, Web Administration and

Development , CEO at IOSRD

COO at GAOR & OSS

Er. Suyog Dixit

(M. Tech), BE (HONS. in CSE), FICCT

SAP Certified Consultant

CEO at IOSRD, GAOR & OSS

Technical Dean, Global Journals Inc. (US)

Website: www.suyogdixit.com

Email: suyog@suyogdixit.com

Pritesh Rajvaidya

(MS) Computer Science Department

California State University

BE (Computer Science), FICCT

Technical Dean, USA

Email: pritesh@computerresearch.org

Luis Galárraga

J!Research Project Leader

Saarbrücken, Germany

CONTENTS OF THE VOLUME

- i. Copyright Notice
- ii. Editorial Board Members
- iii. Chief Author and Dean
- iv. Table of Contents
- v. From the Chief Editor's Desk
- vi. Research and Review Papers

- 1. Characterization of Ascitis using Echo Texture . **1-8**
- 2. Height Difference Between Right and Left Hemi Diaphragm in Normal Sudanese Adults using Postero -Anterior Computed Chest Radiography. **9-15**
- 3. Study of Weekly Paclitaxel with Concurrent Radiotherapy V/S Weekly Cisplatin with Concurrent Radiotherapy in Advanced Cervix Carcinoma. **17-22**
- 4. Perforated Duodenal Diverticulum: Diagnosis and Management. **23 -25**
- 5. Characterization of Hippocampus in Epileptic Sudanese Children using MRI. **27 -35**
- 6. Right Internal Carotid Agenesis: A Rare Case Anatomical, Neurosurgical and Radiological Perspective. **37 -39**

- vii. Auxiliary Memberships
- viii. Process of Submission of Research Paper
- ix. Preferred Author Guidelines
- x. Index



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Characterization of Ascitis using Echo Texture

By Mohammed Saeed AlNumeiri, Caroline Edward Ayad,
Boushra Hussein Ahmed & Elsafi Ahmed Abdulla

Sudan University of Science and Technology, Sudan

Abstract- Objectives: The Purpose of this study is to characterize the asites fluid using echo texture and to compare the findings with laboratory results in order to predict the ascitis type by imaging methods for ascitic patients.

Methods: 53 Patients who underwent Sonography for ascitis between February 2012 and January 2013 were included in the study. The study was implemented in Wad- Medani Military Hospital, Alia Specialist Hospital- Wad-Madani. Wad-medani Teaching Hospital, National Cancer Institute- Wad-Madani and Omdurman Military Hospital. Gray scale Sonography of the abdomen was performed using Aloka SSD- 500 and Honda SSD-500 with frequency (3.5 MHz) convex probe. The scanning was done on longitudinal and transverse scan for abdomen and pelvis.

Keywords: ascitis, ultrasound, laboratory tests.

GJMR-D Classification : NLMC Code: WF 975, WN 1



Strictly as per the compliance and regulations of:



Characterization of Ascitis using Echo Texture

Mohammed Saeed AlNumeiri ^α, Caroline Edward Ayad ^σ, Boushra Hussein Ahmed ^ρ & Elsafi Ahmed Abdulla ^ω

Abstract- Objectives: The Purpose of this study is to characterize the asites fluid using echo texture and to compare the findings with laboratory results in order to predict the ascitis type by imaging methods for ascitic patients.

Methods: 53 Patients who underwent Sonography for ascitis between February 2012 and January 2013 were included in the study. The study was implemented in Wad-Medani Military Hospital, Alia Specialist Hospital- Wad-Madani. Wad-medani Teaching Hospital, National Cancer Institute- Wad-Madani and Omdurman Military Hospital. Gray scale Sonography of the abdomen was performed using Aloka SSD-500 and Honda SSD-500 with frequency (3.5 MHz) convex probe. The scanning was done on longitudinal and transverse scan for abdomen and pelvis. The patients ascitis echo texture had been evaluated and the findings were correlated with Ascitis Paracentesis results, including total protein values, total albumin, Serum Ascitis Albumin Gradient (SAAG), ascitis type, liver and spleen texture as well as the Ultrasound findings.

Results: During this period, 53 patients (31males, 22females) underwent Sonography for ascitis, 5(9.4%) had abdomen tuberculosis, 1(1.9%) acute renal failure, 2(3.8%) Ca breast, 1(1.9%) Ca prostate, 1(1.9%) Ca colon, 2(3.8%) Ca ovaries, 3(5.7%) Congestive heart disease, 6(11.3%) Chronic renal failure, 1(1.9%) Cystadenocarcinoma, 2(3.8%) hepato cellular carcinoma, 1(1.9%) Hepatitis, 23(43.4%) Liver Cirrhosis, 1(1.9%) lymphoma, 1(1.9%) portal hypertension, 2(3.8%) Pelvis inflammatory diseases, 1(1.9%) Urinary bladder Schistosomiasis. The correlation between the Ascitis Echo Texture with total protein values, total albumin, SAAG, ascitis type, ultrasound findings were found to be significant at p value 0.05. liver echo texture has significant relation with Ascitis Echo Texture and laboratory results, but no impact of the spleen texture in the ascites character was noted.

Conclusion: Ultrasound is valuable in the investigation of ascetic patients. It is recommended to replace other methods of laboratory investigation as it can predict the ascitis type from its echo texture, as well as ruling the underlying causes.

Keywords: ascitis, ultrasound, laboratory tests.

I. INTRODUCTION

Ascites is defined as the accumulation of fluids in the peritoneal cavity. [1] Two important factors that control the balance exchange of fluids in the

body, these including the plasma colloid osmotic pressure and the portal venous pressure and its disturbance may cause Ascites. [2, 3, 4]

The most common cause of Ascites from Western Europe and North America is Cirrhosis, which records about 80% of the cases. The other common causes are Malignancy (10%), Cardiac Failure (5%) and Tuberculosis and other causes. [5, 6]

The differentiation between Ascitis causes is important for diagnostic and therapeutic purposes [7] Diagnostic abdominal paracentesis is considered as an essential investigation for differential diagnoses in all patients with clinically noticed Ascites. [8]

Clinically; Ascites is detected by the presence of flank dullness to percussion, [2]. In recent years Ultrasound has been proven an accurate and reliable method of detecting abdominal fluid, determining its site [9] but questions have been raised about its capability to differentiate the Ascitis of transudates concept from inflammatory or malignant exudates. Assessment of the Ascites is necessary in monitoring the progress of the disease and in selecting appropriate methods of treatment. [10]

Radiologic evaluation of the abdomen was helpful in confirming the presence of fluid, Computed Axial Tomography are considered quite sensitive in the detection of ascitic fluid. Physical examination findings are consistent with the diagnosis of an intra-abdominal fluid accumulation; but its sensitivity yields varying results. [2]

To evaluate the reflection of Ascitis Echo Texture related to the fluid composition; we have correlated the Ultrasound findings, with patient history, clinical findings of proven Ascites cases and laboratory findings including Total Protein, Total Albumin and the Serum Ascites Albumin Gradient (SAAG).

II. MATERIALS AND METHODS

This study was implemented in Wad-Medani Military Hospital, Alia Specialist Hospital- Wad-Madani. Wad-Medani Teaching Hospital, National Cancer Institute- Wad-Madani and Omdurman Military Hospital during the period from 2012 to 2013.

Population and samples size: (53) patients in both genders who were clinically diagnosed proved to have abdominal ascites.

Author α: Diagnostic Radiology, Omdurman Military Hospital, Khartoum-Sudan. e-mail: mahmmednumeeri@yahoo.com

Author σ ω: Associate Professor, PhD Diagnostic Radiology, College of Medical Radiological Science, Sudan University Of Science and Technology. e-mail: carolineayad@yahoo.com, dr.elsafi@gmail.com

Author ρ: Professor, PhD Diagnostic Medical Ultrasound, AlRabat University. e-mail: Bushrahuseen@yahoo.com

Instrumentation:

Aloka SSD-500 with frequency (3.5 MHz) convex probe, and Honda SSD-500 with frequency (3.5 MHz) convex probe and the facility of printing and computerized reporting system are used.

Data were collected, executed by direct ultrasound examination, the collected data were analyzed using statistical method, ANOVA independent values, mean and standard deviation, for ethical Issue: permission of department and patients arise at the area of the study were taken.

Techniques for doing examination:**Scanning technique of abdominal and pelvic ultrasound.**

For ultrasound examination, each patient was scanned, in international scanning guidelines and protocols and Ascites was determined by Ultrasound Brightness Mode (B-Mode).

The scanning protocol for Sonography of the peritoneal cavity varies, depending on the patient's clinical symptoms, and the differential clinical diagnosis, No patients preparation was done. Patients were scanned in supine position in right lateral decubitus and in left lateral decubitus.

Transverse section: Transducer was positioned directly under the Xiphisternum and angled cephalic, Transducer beam was swept through the liver, and then was moved distally to umbilicus, applying graded compression, observing lesser sac (around the pancreas) and transverse mesocolon. The transducer was positioned directly below the right costal margin and the scan was repeated, observing all peritoneal spaces. Transducer was placed directly below the left costal margin; transducer was moved in the transverse plane, throughout the remainder of the abdomen to cover the entire abdomen.

Longitudinal section: Transducer was placed below the Xiphisternum, tipped cephalic: Swept across the abdomen from left to right, moved along right costal margin, angling towards the right lateral border of the

liver repeated process to the lateral border of the spleen. Longitudinal scans were performed throughout the remaining abdomen, observing all spaces.

Grading of the findings:**Ultrasound Findings:**

The Liver and spleen echo texture had been evaluated as Fine texture when smooth homogeneous contour of the parenchyma is present, and it was given grade 0 which stands for normal. Coarse texture when irregular contour of the parenchyma and was given grade 1. Diffuse disease is irregular heterogeneous contour with fibrotic changes of parenchyma and was given grade 2.

Sonographic appearance and classification of the ascites was done by giving grades to classify the Ascitis Echo Texture as Clear Ascites and was given grade 0 and Turbid Ascites grade 1

Laboratory investigation after Paracenteses :

Normal Total protein range (6-8)g/dl, Normal serum albumin range (3.5-6.5) g/dl, The Ascites Serum Ascites Albumin Gradient (SAAG) the Serum Albumin Concentration minus the ascitic fluid albumin concentration. Transudate (low in total protein concentration) < 30g/dl, Excaudate (high in total protein concentration) > 30g/dl

III. RESULTS

The sample include 53 patients 31 (58.5%) were male and 22 (41.5%) were females with ascitis arises from different causes, Clinically 32 (60.4%) of the sample have a history of liver disease, 8 (15.1%) have history of renal diseases 8 (15.1%) chest pain, 3 (5.7%) are with hypertension and the residue acts 2 (3.8%) they came with different signs and symptoms including Nausea 3 (5.7%), Anorexia 7 (13.2%), Lower limb edema 13 (35.8%), shortness of breathing 22 (41.5%). All were significantly related to Ultrasound findings at p -value 0.05.

Table 1 : Ultrasound Findings/Diagnosis in patients with ascitis

Ultrasound Findings /Diagnose	Frequency	Percentages
Abdomen Tuberculosis	5	9.4%
Acute Renal Failure	1	1.9%
Breast Cancer	2	3.8%
Prostate Cancer	1	1.9%
Colon Cancer	1	1.9%
Ovarian cancer	2	3.8%
Congestive Heart Diseases (CHD)	2	3.8%
Congestive Heart Diseases+ Chronic Renal Failure	1	1.9%
Chronic Renal Failure	6	11.3%

Cystoadenocarcinoma	1	1.9%
Hepato Cellular Carcinoma(HCC)	2	3.8%
Hepatitis	1	1.9%
Liver Cirrhosis	23	43.4%
Lymphoma	1	1.9%
Portal Hyper Tension(PHTN)	1	1.9%
Pelvic Inflammatory Diseases(PID)	2	3.8%
Urinary Bladder Schistosomiasis	1	1.9%
Total	53	100.0%

Table 2 : Association between ultrasound findings /diagnosis** /classifications& (SAAG Classes (*P-Value* = 0.000*))

Diagnose**	SAAG Classes			Total
	High gradient	Normal	Low gradient	
Inflammatory	7.0(46.7%)	6.0(50.0%)	3.0(11.5%)	16.0(30.2%)
Tumors	6.0(40.0%)	4.0(33.3%)	0.0(0.0%)	10.0(18.9%)
Cardiac & PHT	1.0(6.7%)	1.0(8.3%)	2.0(7.7%)	4.0(7.5%)
Liver cirrhosis	1.0(6.7%)	1.0(8.3%)	21.0(80.8%)	23.0(43.4%)
Total	15.0(100.0%)	12.0(100.0%)	26.0(100.0%)	53.0(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.** The ultrasound findings /diagnosis were classified into inflammatory , tumors, cardiac diseases and portal Hyper tension (PHT) and liver cirrhosis

Table 3 : ultrasound findings/Diagnosis classifications, Total protein and Albumin values

Diagnose	Total protein	Albumin
Inflammatory	8.2 ± 0.9	2.8 ± 0.7
Tumors	8.5 ± 0.7	3.2 ± 0.7
Cardiac & PHT	7.1 ± 1.3	2.7 ± 1.0
Liver cirrhosis	6.3 ± 0.9	2.0 ± 0.4
P-value	0.000*	0.000*

Values are expressed as Mean ± SD; * Significant at *P*-value < 0.05.

Table 4 : Association between (ultrasound findings /diagnosis classifications) & (ascitis echo texture) (*P-Value* = 0.000*)

Diagnose	Ascitis Echo Texture		Total
	Clear	Turbid	
Inflammatory	10.0(25.0%)	6.0(46.2%)	16.0(30.2%)
Tumors	3.0(7.5%)	7.0(53.8%)	10.0(18.9%)
Cardiac & PHT	4.0(10.0%)	0.0(0.0%)	4.0(7.5%)
Liver cirrhosis	23.0(57.5%)	0.0(0.0%)	23.0(43.4%)
Total	40.0(100.0%)	13.0(100.0%)	53.0(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 5 : Association between (Ascites type) & (Sonographic Appearance) (*P-Value* = 0.000*)

<i>Sonographic Appearance</i>	Ascites type		Total
	Transudate	Excaudate	
Clear	34.0(85.0%)	6.0(15.0%)	40.0(100.0%)
Turbid	1.0(7.7%)	12.0(92.3%)	13.0(100.0%)
Total	35.0(66.0%)	18.0(34.0%)	53.0(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 6 : Ascites type, Total Protein, Albumin value, Ascitis Serum albumin gradient (SAAG), Sonographic Appearance

Ascites type	Total Protein	Albumin value	SAAG Value	Sonographic Appearance
Transudation	6.7 ± 1.1	2.3 ± 0.7	1.7 ± 0.6	0.02 ± 0.2
Exudates	8.5 ± 0.9	2.9 ± 0.8	0.3 ± 0.6	0.7 ± 0.5
P-value	0.000*	0.009*	0.000*	0.000*

Values are expressed as Mean ± SD; * Significant at P-value < 0.05.

Table 7 : Association between ultrasound findings /Diagnosis classifications & Asites type (*P-Value* =0.000*)

Diagnose		Asites type		Total
		Transudate	Excaudate	
Inflammatory	Count	7.0(20.0%)	9.0(50.0%)	16.0(30.2%)
Tumors	Count	1.0(2.9%)	9.0(50.0%)	10.0(18.9%)
Cardiac & PHT	Count	4.0(11.4%)	0.0(0.0%)	4.0(7.5%)
Liver cirrhosis	Count	23.0(65.7%)	0.0(0.0%)	23.0(43.4%)
Total	Count	35.0(100.0%)	18.0(100.0%)	53.0(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 8 : Association between Liver texture & Ascitis Echo Texture (*P-Value* = 0.005*)

Liver texture	Ascitis Echo Texture		Total
	Clear	Turbid	
Fine	13(32.5%)	11(84.6%)	24(45.3%)
Coarse	10(25.0%)	1(7.7%)	11(20.8%)
Diffuse	17(42.5%)	1(7.7%)	18(34.0%)
Total	40(100.0%)	13(100.0%)	53(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 9 : Association between Liver texture &SAAG Class (*P-Value* = 0.000*)

Liver texture	SAAG Class			Total
	High gradient	Normal	Low gradient	
Fine	10(66.7%)	11(91.7%)	3(11.5%)	24(45.3%)
Coars e	2(13.3%)	0(0.0%)	9(34.6%)	11(20.8%)
Diffuse	3(20.0%)	1(8.3%)	14(53.8%)	18(34.0%)
Total	15(100.0%)	12(100%)	26(100%)	53(100.%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 10 : Association between Liver texture & Asites type (*P-Value* = 0.003*)

Liver texture	Asitestype		Total
	Transudate	Excaudate	
Fine	10(28.6%)	14(77.8%)	24(45.3%)
Coarse	9(25.7%)	2(11.1%)	11(20.8%)
Diffuse	16(45.7%)	2(11.1%)	18(34.0%)
Total	35(100.0%)	18(100.0%)	53(100.0%)

*Normal p-value is < 0.05; therefore the statistical difference is highly significant.

Table 11 : Association between Spleen texture & Ascitis Echo texture (*P-Value* = 0.326)

Spleen texture	Ascitis Echo texture		Total
	Clear	Turbid	
Fine	29(72.5%)	12(92.3%)	41(77.4%)
Coarse	10(25.0%)	1(7.7%)	11(20.8%)
Diffuse	1(2.5%)	0(0.0%)	1(1.9%)
Total	40(100.0%)	13(100.0%)	53(100.0%)

Table 12 : Association between Spleen texture & SAAG Class (*P-Value* = 0.336)

Spleen texture	SAAG Class			Total
	High gradient	Normal	Low gradient	
Fine	13(86.7%)	11(91.7%)	17(65.4%)	41(77.4%)
Coars e	21(3.3%)	1(8.3%)	8(30.8%)	11(20.8%)
Diffuse	0(0.0%)	0(0.0%)	1(3.8%)	1(1.9%)
Total	15(100.0%)	12(100.0%)	26(100.0%)	53(100.0%)

Table 13 : Association between Spleen texture & Asites type (*P-Value* = 0.101)

Spleen texture	Asites type		Total
	Transudate	Excaudate	
Fine	2468.6%	1794.4%	4177.4%
Coarse	1028.6%	15.6%	1120.8%
Diffuse	12.9%	00.0%	11.9%
Total	35100.0%	18100.0%	53100.0%

Table 14 : Liver texture with Total Protein & total albumin

Liver Texture	Total Protein	Total Albumin
Fine	8.2 ± 0.8	3.0 ± 0.7
Coarse	7.0 ± 1.1	2.4 ± 0.5
Diffuse	6.3 ± 1.2	2.0 ± 0.5
P-value	0.000*	0.000*

Values are expressed as Mean ± SD; * Significant at P-value < 0.05.

Table 15 : Spleen texture with Total Protein & total albumin

Spleen Texture	Total Protein	Total Albumin
Fine	7.6 ± 1.3	2.7 ± 0.8
Coarse	6.4 ± 1.2	1.9 ± 0.3
Diffuse	6.8 ± 0.0	2.1 ± 0.0
P-value	0.028	0.005

Values are expressed as Mean ± SD; * Significant at P-value < 0.05.

IV. DISCUSSION

The evaluation of Ascites is one of the earliest uses of abdominal ultrasound, since this method is suitable for fluids characterization [11] the ultrasound findings and diagnosis as frequency and percentage were presented in table [1].

The clinical signs and symptoms, causes hint that may be due to pressure on the internal organs, veins, diaphragm and sodium and water retention [12]. Association between ultrasound findings/diagnosis and clinical findings were found to be significant at *P-Value* (0.001). These clinical history associated with the ascites criteria may narrow the line for diagnoses and may predict the underlying cause of the diseases. The knowledge about the patients' previous history of diseases is important to be correlated with ultrasound findings as it was mentioned in the guidelines on the ascites management. [6]

Ultrasound has great value in diagnosing ascites causes. The study showed that the causes are either from presence of malignancies, inflammatory, or cardiac in origin where the highest score of the ascites causes was found in patients affected by liver cirrhosis

23(43.4%) table[1] another studies had mentioned similar causes of ascites. [5,6]

For studying the character of ascites; SAAG, Total Albumin, Total Protein values have been evaluated and correlated with ultrasound findings. The study showed a significant relation with the variables at *P-Value* (0.000)table. [2,3]

15 out of 53(28.3%) were of high Serum Ascitis Albumen Gradient (SAAG), 12(22.6%) were normal and 26(49.1%) were of low Serum Ascitis Albumen Gradient (SAAG). The study showed that the cases of congestive heart diseases and portal hypertension as well as liver cirrhosis have low SAAG values table [2]. In Mauer and Mazzone, [13] assessment of ascites show that SAAG best reflected elevated hydrostatic pressure, contributing significantly to the development of portal hypertensive ascites, even in patients with a low serum albumin concentration as well as SAAG have value in distinguishing congestive heart failure and malignant ascites.

The Total Protein and Total Albumin, were found to be significantly correlated with the ultrasound findings at *P-Value* (0.000). The high value was found in patients

with tumors and low values in patients with congestive heart failure, portal hypertension, and liver cirrhosis table [3]. Similarly in the study of Rector and Reynolds [14] patients with chronic heart failure and cirrhosis ascites had lower fluid total protein concentrations.

After ultrasound scanning, the ascites echo texture has been evaluated table [4] and were classified into 2 main appearances: 40 out of 53 (75.5%) were clear and 13 out of 53 (24.5%) were turbid. The cause of changing the echo texture of the ascites fluid collections may be either due to debris appears as multiple fine echoes in the cases of inflammation, or the coarse network of echoes generated from the fibrous strings. The clear appearance of most ascitic fluids reflects the etiology as cirrhosis. [15] similarly in our study the cases of cirrhosis acts 23 (43.4%) with clear ascites and no turbid ascites is noted, but in the inflammatory cases which acts 16 (30.2%) patients; 10 cases were found to be clear and 6 cases were turbid. Turbidity, may be due to presence of neutrophils or malignancy due to triglyceride seen in the fluid. [16] as our study includes 5 cases of abdominal TB (9.4%), 1 case of Hepatitis (1.9%), 2 cases of pelvic inflammatory diseases (3.8%) and many cases of malignancies in the breast, prostate, colon, bladder, ovaries, hepatocellular carcinoma (HCC), and lymphoma. A number of different proteins have been present in the malignant cases and should be evaluated for their usefulness in diagnosing malignancy-related ascites in combination with other parameters such as SAAG, that may change the character of the ascites. [17,18]

The importance of differentiation between ascites type; is that malignant ascites is of critical predictive significance with survival. [19]

The clear sonographic texture of ascites were found to be transudate in origin 34.0 (85.0%) and exudates 6.0 (15.0%) where turbid texture were found to be transudate 1.0 (7.7%) and exudates in 12.0 (92.3%) table. [5]

The ascites type had also been evaluated and was found to be related significantly at P -value < 0.05 . With total protein values, Total Albumin value, Ascites Serum albumin gradient (SAAG) and sonographic ascites texture as turbid and clear table. [6]

The causes of ascites formation as transudate or exudate do not always follow the expected pattern [20] it was therefore proposed that the transudate or exudate concept for the classification of ascites should be terminated. [21] Additional tests and methods should be considered.

The Ascites Total Protein (TP) attempting to reflect the different processes of fluid formation, studies supported that SAAG act as a marker of portal hypertension this approach has been adopted in the British and the American Guidelines. [6]

The demonstration of ascites has been aided by Ultrasound, as it can scan through many anatomic

planes and it can detect as little as 10 ml of free fluid [22,23] therefore, uses of Ultrasound was found to be useful as it can characterize the ascites and find out the causes.

The clinical findings do not reflect the changes in the echo texture in both the liver and spleen. We consider that ultrasound is valuable in the investigation of patients who are difficult to examine clinically, abdominal ultrasound scan should be done in order to evaluate the appearance of the liver, pancreas, and lymph nodes as well as the presence of splenomegaly, which may be a sign of portal hypertension. [6]

Association between liver Echo Texture with Total Protein, Total Albumin, SAAG, ascites type, ascites echo texture was found to be significant at p -value < 0.05 table [8,9,10,14] but the spleen echo texture have significant relations only with Total Protein and Total Albumin values but no relationship was noted significantly with the other examined parameters, as seen in table. [11,12,13]

We believe that Ultrasound should take place in the investigation of abdomen and pelvis for ascites, it is also valuable in the investigation of ascitic patients with large or small ascites amount of fluids and patients who are difficult to be examined clinically. It is recommended to replace other methods of laboratory investigation as it can predict the ascites type from its echo texture, as well as ruling the underlying causes.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Runyon BA. 1989, Ascites, ascitic fluid infection and hepato renal syndrome. In: Beker S, ed. Hepatology for the Clinician: A Problem Oriented Approach. New York: Alan R Liss Inc.; 105–29.
2. Cattau EL, Benjamin SB, Knuff TE, Castell DO. 1982, The accuracy of the physical examination in the diagnosis of suspected ascites. JAMA; 247: 1164–6.
3. Goldberg BB, Goodman GA, Clearfield HR. 1970, Evaluation of ascites by ultrasound. Radiology; 96:15–22.
4. Sherlock S. Ascites. In: Sherlock S, ed. 1981, Diseases of the Liver and Biliary System. 6th edn. Oxford: Blackwell Scientific Publications Ltd.; 119–30.
5. Runyon BA. 2004, Management of adult patients with ascites due to cirrhosis, AASLD Practice Guideline. Hepatology; 39:1–16.
6. Moore KP, Aithal GP. 2006, Guidelines on the management of ascites in cirrhosis. Gut; 55(Suppl. VI):vi1–12.
7. Anita R. Bijoor And T. Venkatesh, 2001, Value Of Ascitic Fluid Cholesterol And Serum - Ascites Albumin Gradient In Differentiating Cirrhotic And Malignancy Related Ascites Indian Journal Of Clinical Biochemistry; 16(1), 106.109.

8. Hoefs JC. 1990, Diagnostic paracentesis: a potent clinical tool. *Gastroenterology*;98:230–6.
9. Goldberg BB: 1976, Ultrasonic evaluation of intraperitoneal fluid. *JAMA* 235:2427-2430.
10. Szkodziak P. R, Wozniak S, Czuczwar P, Kludka-Sternik M, Paszkowski M, Paszkowski T. Ascites index – a new method of ultrasound evaluation of ascites volume in patients with ovarian cancer, 10–14 October 2010, Prague, Czech Republic Poster abstract.s
11. Goldberg BB, Goodman GA, Clearfield HR: 1970, Evaluation of ascites by ultrasound. *Radiology* 96 : 15-22.
12. Shah R et al; Ascites, Medscape, May 2009.
13. Mauer K, Manzione NC. 1988, Usefulness of serum-ascites albumin difference in separating transudative from exudative ascites; another look. *Dig Dis Sci*;33:1208–12.
14. Rector WG, Reynolds TB. 1984, Superiority of the serum-ascites albumin difference over the ascites total protein concentration in separation of transudative and exudative ascites. *Am J Med*; 77: 83–5.
15. Mchutchison JG. 1997, Differential diagnosis of ascites. *Semin Liver Dis*;17:191–202.
16. Runyon BA, Akriviadis EA, Keyser AJ. 1991, the opacity of portal hypertension-related ascites correlates with the fluid's triglyceride concentration. *Am J Clin Pathol*;96:142–3.
17. Villamil FG, Sorroche PB, Aziz HF, Lopez PM, Oyhamburu JM. 1990, Ascitic fluid alpha 1-antitrypsin. *Dig Dis Sci*;35:11105–9.
18. Kountouras J, Boura P, Tsapas G, Charisis K, Magoula I, Tsakiri I. 1993, Value of ascitic fluid ferritin in the differential diagnosis of malignant ascites. *Anticancer Res*;13:2441–5.
19. Sears D, Hajdu SI. 1987, the cytologic diagnosis of malignant neoplasms in pleural and peritoneal effusions. *Acta Cytol*;31:85–97.
20. Hoefs JC. 1983, Serum protein concentration and portal pressure determine the ascitic fluid protein concentration in patients with chronic liver disease. *J Lab Clin Med*;102:260–73.
21. Tarn A C and Lapworth R, 2010, Biochemical analysis of ascitic (peritoneal) fluid: what should we measure? *Ann Clin Biochem*; 47: 397–407.
22. Forsby J, Henriksson L: 1984, Detectability of intraperitoneal fluid by ultrasonography: an experimental investigation. *Acta Radiol [Diagn] (Stockh)*; 25: 375-378.
23. Dinkel E, Lehnart R, Troger J et al: 1984, Sonographic evidence of intraperitoneal fluid: an experimental study and its clinical implications. *Pediatr Radiol*; 14: 299-303.



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Height Difference between right and left hemi Diaphragm in Normal Sudanese Adults using Postero-Anterior Computed Chest Radiography

By Hassan Ali Abdelrahman Salih, Caroline Edward Ayad,
Elsafi Ahmed Abdalla, Ikhlas Abdelaziz Hassan & Abdelmoneim Saeed Ahmed

Sudan University of Science and Technology, Sudan

Abstract- Diaphragm position and shape on chest radiographs are routinely used as indicators for diseases. Previous population studies were done for normal diaphragm position and shape.

The study was done to determine the distance between right and left hemi diaphragm in normal Sudanese adults using digital radiography.

The study sample was 100 subjects, 50(50%) were males and 50(50%) were females, with different ages ranged from 18 – 82 years old; all subjects were with normal abdominal ultrasound and normal chest radiographs. The study was done in Fedail hospital in the period from January up to March 2013. The measurements were done on ideal postero anterior (PA) chest radiograph by measuring the difference between the distance from the highest point of right hemi diaphragm and highest point of the left hemi diaphragm. The result showed that the right hemi diaphragm was higher than the left in 98% of total sample and the left was higher in 1% and they were in the same height in 1%.

Keywords: chest x-ray, hemi diaphragm, thoracic, digital radiography.

GJMR-D Classification : NLMC Code: WN 180, WF 800



Strictly as per the compliance and regulations of:



Height Difference between right and left hemi Diaphragm in Normal Sudanese Adults using Postero-Anterior Computed Chest Radiography

Hassan Ali Abdelrahman Salih ^α, Caroline Edward Ayad ^ο, Elsafi Ahmed Abdalla ^ρ
Ikhlaz Abdelaziz Hassan ^ω & Abdelmoneim Saeed Ahmed [¥]

Abstract- Diaphragm position and shape on chest radiographs are routinely used as indicators for diseases. Previous population studies were done for normal diaphragm position and shape.

The study was done to determine the distance between right and left hemi diaphragm in normal Sudanese adults using digital radiography.

The study sample was 100 subjects, 50(50%) were males and 50(50%) were females, with different ages ranged from 18 – 82 years old; all subjects were with normal abdominal ultrasound and normal chest radiographs. The study was done in Fedail hospital in the period from January up to March 2013. The measurements were done on ideal postero anterior (PA) chest radiograph by measuring the difference between the distance from the highest point of right hemi diaphragm and highest point of the left hemi diaphragm. The result showed that the right hemi diaphragm was higher than the left in 98% of total sample and the left was higher in 1% and they were in the same height in 1%.

The right was higher in range of 1cm to 2.5cm in 79% and the Male diaphragm height difference ranged between 0.6-2.8 and Female ranged from 0.5-2.9cm. The study found that there are linear relation between distance difference of right and left hemi diaphragm with subjects' age, body mass index (BMI), height and weight.

Computed radiography was useful in measurements of diaphragm height difference for Sudanese as it is accurate and easy.

Keywords: chest x-ray, hemi diaphragm, thoracic, digital radiography.

I. INTRODUCTION

The diaphragm is a curved muscle fibrous sheet that separates the thoracic from abdominal cavity, the position of the domes or cupola of the diaphragm are extremely variable as they differ on body built. [1-3]

Authors α ο ω: MSc Diagnostic Radiology, Sudan University of Science and Technology, College of Medical Radiological Science.
e-mails: HassanAli@yahoo.com, carolineayad@yahoo.com, dralsaffi@gmail.com

Author ρ: PhD Diagnostic Medical Ultrasound, Sudan University of Science and Technology, College of Medical Radiological Science.
e-mail: Eklassabdelaziz@yahoo.com

Author ¥: Diagnostic Radiology, National College of Medical and Technical Studies. e-mail: abdelmoneimsaeed@yahoo.com

Although the use of computerized Tomography has greatly increased over the past decades, chest radiography remains the most frequently performed imaging examination [4]. A good understanding of normal anatomy and variations is essential for the interpretation of postero anterior (PA) chest radiographs include fissure lines [5, 6, 7, 8, 9], and vascular structures [10], aortic nipple [11, 12, 13] is important.

When interpreting chest radiographs, the position and shape of the diaphragm are routinely used to evaluate whether the lungs are underinflated, as with inadequate inspiration effort or restrictive lung disease, or overinflated, as in emphysema or other obstructive lung diseases. Conventional teaching that the normal position of the right hemi diaphragm projects at about the anterior sixth rib [14, 15] appears to originate from a single study.[16]

A more precise characterization of the normal variation in diaphragm position and shape, based on easily recognized anatomic landmarks, and analysis of factors that might contribute to this variation, such as age and weight, may provide a more consistent origin for this review.[17]

There have been few reports describing the normal radiographic anatomy and variations of the thoracic structures in different population [13]. Most of the data regarding these normal structures have been provided from the Europe and USA. [5–12]

Therefore we reviewed normal PA chest radiographs and analyzed the radiographic anatomy among Sudanese. To our knowledge no information is available in the open literature about Sudanese population. Therefore, we conducted this study in order to measure the height difference between right and left hemi diaphragm on PA chest Digital radiographs obtained from Sudanese subjects with normal chest and abdomen, and to identify the relationship of diaphragm position to Sudanese demographic data including age gender, weight, height and BMI.

II. MATERIALS AND METHODS

a) Patients

The sample was 100 Sudanese Subjects (50 males and 50 female) with age range from 18-82 years.

The chest radiographs were obtained as a part of routine work-up or follow-up study for non-chest diseases, all patients were diagnosed as normal chest x-ray and the abdominal ultrasound was diagnosed as normal; Subjects with plural effusion, lung diseases, scoliosis, sarcoidosis, collagen vascular disease, pulmonary hypertension, large cardio thoracic ratio or, large volume of the abdominal contents, ascites, liver cirrhosis, liver or spleen enlargement disease were excluded.

The x-ray machine: major x-ray machines, Shimadzu were used.-Manufacturer: Shimadzu, Japan. Manufacturing date: Jan 2011. Focal spot: Small (0.6) mm / Large (1.2) mm. Maximum kV: 150 kVp. Maximum mA: 500mA. Year of Installation: 2011.

Processing machine: Images processed by using computed radiography, Fuji film (FCR CAPSULLA) model XL II.

Imaging technique: Firstly the procedure was explained to the patients (pts). Then the pts were asked to change his/her clothes with the hospital gown. The patients were centered in the erect position in the stand bucky as for ideal postero-anterior chest radiograph and asked to take a deep breath and hold it.

Postero-anterior chest radiograph technique: A 35.43cm or 35.35cm cassette is selected, depending on the size of the patient. The patient is positioned facing the cassette, with chin extended and centered to the middle of the top of the cassette. The feet are placed slightly apart so that the patient is able to remain steady. The median sagittal plane is adjusted at right angles to the middle of the cassette. The shoulders are rotated forward and pressed downward in contact with cassette.

This is achieved by placing the dorsal aspect of the hands behind and below the hips, with the elbows brought forward, or by allowing the arms to encircle the cassette. The horizontal central beam is directed at right angles to the cassette at the level of thoracic vertebra which is consigned with lung midpoint. The surface marking of T7 spinous process can be assessed by using the inferior angle of the scapula. Exposure is made in full normal arrested inspiration.[18]

The exposure made at inspiration. Images were processed using Fuji computed radiography system (FCR). Images were manipulated, adjusted, and then stored in the computer.

b) Method of measurement

The subjects were asked for their ages and the weight had been registered using weights measuring device, the heights were measured by a meter. After ideal chest radiograph acquisition and processing then measurements were done. A straight line was drawn at the maximum height of each hemi diaphragm. The lines are parallel to each other. Then the distance between the both lines was measured.

III. RESULTS

The following figures and tables presented the data obtained from 100 normal subjects after measuring the distance between the right and left hemi diaphragm, the variables taken were age, height (m), weight (Kg), Body Mass Index (BMI).Correlations were taken by presenting trend line and resultant equations in the graphs, this was done for males who act 50% of the sample and the females who act 50% of the sample as well as the Total sample (100%).

Table 1 : mean and standard deviation of the variables (age, Hemi diaphragm distance, weight, height, BMI for male, Female and total sample

Variables	Male (mean± STDV)	Female (mean± STDV)	Total Sample (mean± STDV)
Ages (years)	39.98 ±16.8	44.04±16.6	42.01±16.7
Hemi Diaphragms			
Distance	1.6±0.6	1.42±0.7	1.5±0.6

Weight (Kg)	74.5±13.2	69.92±13.2	72.2±13.3
Height (M)	1.7±0.1	1.64±0.1	1.7±0.1
(BMI)	24.3±4.0	25.65±4.8	24.9±4.5

Table 2 : Hemi Diaphragms distance Classes for the Total Sample, Male and Female, Frequency and Percentages

Hemi Diaphragms distance (Total sample)	Frequency And Percentage%	Diaphragm distance (Males)	Frequency And Percentage%	Diaphragm distance (Females)	Frequency And Percentage%
-0.5—0.5	5(5%)	0.6-1.0	11(22%)	-0.5-0.5	5(10%)
0.6-1.1	23(23%)	1.4-1.8	21(42%)	0.6-1.1	10(20%)
1.2-1.7	38(38%)	1.9-2.3	11(22%)	1.2-1.7	22(44%)
1.8-2.2	22(22%)	2.4-2.8	6(12%)	1.8-2.3	9(18%)
2.3-2.8	11(11%)	>2.8	1(2%)	2.4-2.9	4(8%)
>2.8	1(1%)	-	-	-	-

Table 3 : Height Classes in Meters (M) For the Total Sample, Male and Female, Frequency and Percentages

Total sample Height(M)	Frequency And Percentage%	Males Height (M)	Frequency And Percentage%	Females Height (M)	Frequency And Percentage%
1.37-1.47	1(1%)	1.37-1.47	1(2%)	1.55-1.59	11(22%)
1.48-1.58	8(8%)	1.48-1.58	1(2%)	1.60-1.64	15(30%)
1.59-1.69	38(38%)	1.59-1.69	7(14%)	1.65-1.69	12(24%)
1.70-1.80	42(42%)	1.70-1.80	30(60%)	1.70-1.74	11(22%)
1.81-1.91	10(10%)	1.81-1.90	10(20%)	1.75-1.79	1(2%)
>1.91	1(1%)	>1.90	1(2%)	-	-

Table 4 : Weight Classes in kilograms (kg) For the Total Sample, Male and Female, Frequency and Percentages

Total sample Weight(Kg)	Frequency and Percentage%	Males Weight (Kg)	Frequency and Percentage%	Females Weight (Kg)	Frequency and Percentage%
44-54	6(6%)	44-54	2(4%)	48-58	9(18%)
55-65	26(26%)	55-65	12(24%)	59-69	18(36%)
66-76	36(36%)	66-76	15(30%)	70-80	16(32%)
77-87	19(19%)	77-87	12(24%)	81-91	3(6%)
88-98	10(10%)	88-98	8(16%)	92-102	2(4%)
>98	3(3%)	99-109	1(2%)	>102	2(4%)

Table 5 : Body /Mass Index (BMI) Classes for the Total Sample, Male and Female, Frequency and Percentages

Total sample (BMI)	Frequency and Percentage%	Male (BMI)	Frequency and Percentage%	Female (BMI)	Frequency and Percentage%
18.1-19.6	9(9%)	18.1-21.1	9(18%)	18.5--23.1	19(38%)
20.6-22.2	24(24%)	22.1-25.1	22(44%)	24.1-28.8	22(44%)
23.2-24.8	22(22%)	26.1-29.1	13(26%)	29.8-34.5	7(14%)
25.8-26.3	12(12%)	30.1-33.1	5(10%)	35.5-40.2	1(2%)
27.3-28.8	18(18%)	34.1-37.1	1(2%)	>40	1(2%)
>28.8	15(15%)	-	-	-	-




Figure 1 : [A] Scatter Plot Diagramme shows linear relationship between the diaphragm heights in males with age; the relation shows that the height decreased by 4.8 as the age increased starting from 47.7. [B] shows linear relationship between the diaphragm heights in males with BMI; the relation shows that the BMI decreased by 0.9




Figure 2 : [A] Scatter Plot Diagramme shows linear relationship between the diaphragm heights in males with Height; the relation shows that the height increased by 0.009 as the diaphragm distance increased starting from 1.72. [B] shows linear relationship between the diaphragm heights in males with Weight; the relation shows that the diaphragm height decreased as the weight increased.




Figure 3 : [A] Scatter Plot Diagramme shows linear relationship between the diaphragm heights in Females with age; the relation shows that the height increased by 0.6 as the age increased starting from 43.2. [B] shows linear relationship between the diaphragm heights in Females with BMI; the relation shows that the height increased by 1.4 as the BMI increased starting from 23.7




Figure 4 : Scatter Plot Diagramme shows linear relationship between the diaphragm heights in Females with Weight; the relation shows that the height increased by 2.6 as the Weight increased starting from 66.2.[B] shows linear relationship between the diaphragm heights in Females with Height; the relation shows that the Diaphragm height decreased by 0.01 as the Female Height increased starting from 1.7 meter.




Figure 5 : Scatter Plot Diagram shows linear relationship between the diaphragm heights in Total Sample with age; the relation shows that the height decreased by 2.3 as the age increased starting from 45.5.[B] shows linear relationship between the diaphragm height and BMI, The diaphragm height increase as the BMI increased.

IV. DISCUSSION

Measurements were performed on ideal PA chest radiograph, measurement points was clearly identified. Table 1 shows the total sample means and standard deviations of the variables, the total sample age mean was 42.01 and ± 16.73 , the total sample weight mean was 72.2 ± 13.34 , the total sample height mean was 1.6926 ± 0.09 , the total sample body mass index (BMI) was 24.957 ± 4.45 and the total sample diaphragm distance mean was 1.522 ± 0.64 .

The mean age for male was 39.98 it was lower than mean age for females which was 44.04. The mean of male weight was 74.48 and for female was 69.92, the height of males was greater by 0.1m, the BMI mean for male was 24.26 and for female 25.65, the BMI of female was greater than male because male height was greater. The total sample age range was from 18 to 85, for male the range was from 18 to 82, while for female the range was from 19 to 85. Age classes for total sample, frequency and percentage were studied.

For the total sample the right hemi diaphragm was higher than the left hemi diaphragm in 98 sample which was act 98% of the total sample, the left hemi diaphragm was higher in 1 sample which was 1% of the total sample and they was in the same height in also 1 sample which was 1% of the total sample.

In males samples the right was higher in 50 sample (100%), while in females the right was higher in 48 samples (96%), the left was higher in 1 sample (2%) and they was in the same height In 1 sample (2%).For the total sample 79(79%) of the right hemi diaphragm was higher in range of 1.0 to 2.5 cm, 7(7%) were in range of 2.5 to 3.0 cm, 12 were in range of 0.1 to 0.9, 1 (1%) the right and the left were in the same height and in 1 sample the left was higher. Of 1 person with a high diaphragm was overweight; It is possible that he took an improperly deep inspiration. There were two instances where the diaphragm in addition to being low was also flat, but there were no cardiovascular or lung disease this is similar to what had been mentioned previously. [19]

For the males the distance between right and left hemi diaphragm is decreased as the age, BMI and weight increased subsequently , while it increased as the height increased For the females the distance between right and left hemi diaphragm increased as the age, BMI and weight increased as ,while it decreased as the height increased For the total, the distance between right and left hemi diaphragm decreased as the age increased while the distance increased as the BMI, weight and height increased .

Simile study showed that the averages the diaphragm was slightly higher in females, in heavy persons and after the age of 40 years. [19]

V. CONCLUSION

The study found that the right hemi diaphragm is higher than the left in 98% of sample, the left hemi diaphragm may be higher than the right but it consider significant and can be at the same height.

There is a correlation between hemi diaphragms distance and sample age, BMI, weight and height; the distance decrease with increase of age, the distance increase as increase of BMI, the distance increase as increase of weight and the distance increase as increase of height.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Alfred P.Fishman-2005One hundred year of Chronis pulmonary disease, American journal of Respiratory and critical Care Medicine; Volume 171, pageno: 941-948.
2. Grays anatomy 39th edition, page nos-1081, 1082.
3. Vincentcoltin,Nathalie Freymond Jean Cabane and Jean Fancois Cordier-2011.Combined pulmonary fibrosis and emphysema syndrome in a patient age 28 years with sever systemic sclerosis the journal of Rheumatology; Volume 38,2082-2083.
4. Abiru H, Ashizawa K, Hashmi R, Hayashi K, 2005Normal radiographic anatomy of thoracic structures: analysis of 1000 chest radiographs in Japanese population The British Journal of Radiology, 78 ,398–404.
5. Felson B. 1973Chest roentgenology. Philadelphia, PA: WB Saunders.
6. Proto AV, Ball JB Jr. 1983The superolateral major fissures. AJR Am J Roentgenol; 140:431–7.
7. Proto AV. 1984The chest radiograph: anatomic considerations.Clin Chest Med; 5:213–46.
8. Friedman E. 1966Further observations on the vertical fissure line. Am J Roentgenol Radium Ther Nucl Med; 97:171–3.
9. Godwin JD, Tarver RD. 1985Accessory fissure of the lung. AJR Am J Roentgenol; 144:39–47.
10. Proto AV, Chaliff MI. 1986Apical opacity: a normal finding on posteroanterior chest radiographs. Radiology; 161:429–32.
11. Ball JB Jr, Proto AV. 1982The variable appearance of the left superior intercostal vein. Radiology; 144:445–52.
12. Friedman AC, Chambers E, Sprayregen S. The normal and abnormal left superior intercostals vein. AJR Am J Roentgenol 1978; 131:599–602.
13. Takahashi K, Shinozaki T, Hyodo H, Ogawa C, Ohsawa T. 1994Focal obliteration of the descending aortic interface on normal frontal chest radiographs: correlation with CT findings. Radiology; 191:685–90.
14. Fraser RS, Mu" ller NL, Colman N, et al. 1999The respiratory muscle and chest wall. In: Fraser RS, ed. Fraser and Pare's diagnosis of diseases of the chest. Philadelphia, PA: W.B. Saunders,; 246–268.
15. Armstrong P. The normal chest. In: Armstrong P, ed. 2000imaging of diseases of the chest. London, UK: Mosby,; 21–62.
16. Lennon EA, Simon G. The height of the diaphragm in the chest radiograph of normal adults. Br J Radiol 1965; 38:937– 943.
17. Thitiporn Suwatanapongched, David S. Gierada; Richard M. Slone, Thomas K. Pilgram,; and Peter G. Tuteur, Variation in Diaphragm Position and Shape in Adults With Normal pulmonary Function* www.chestjournal.org CHEST / 123 / 6 / JUNE, 2003 2019-2027.
18. Stewart. A, Sloane. C, Hoadley. G, Adrian .D and Chrissie .W .A, CLARK'S positioning in radiography. 12th. Hodder Arnold, London, 2005, (206-207).
19. Lennon E. A., G. Simon, 1965.The Height Of The Diaphragm In The Chest Radiograph Of Normal Adults Brit. J. Radiol, 38, 937-943.



This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Study of Weekly Paclitaxel with Concurrent Radiotherapy V/S Weekly Cisplatin with Concurrent Radiotherapy in Advanced Cervix Carcinoma

By Dr. Prashant Rajendra Kumbhaj, Dr. Rameshwaram Sharma,
Dr. Aseem Rai Bhatnagar & Dr. Peeyush Kumar Saini

Department of Radiotherapy & Oncology SMS Medical College & Attached hospitals, India

Abstract- Purpose: To evaluate the Response and Efficacy of weekly paclitaxel with concurrent radiotherapy in comparison to weekly cisplatin with concurrent radiotherapy in advanced cervical cancer patients.

Methods: Patients coming to the department of radiotherapy S.M.S Hospital were included in the study. 50 patients of advanced squamous cell carcinoma of cervix FIGO stage III & IV were selected. In the study and control group, patients were given 5 weeks of external beam radiotherapy followed by brachytherapy. External beam irradiation each of 200cGy/fr, on linear accelerator by parallel opposed – anterior and posterior fields was given. External beam irradiation was followed by Intracavitary brachytherapy 700cGy/week for 3 weeks each with High Dose Rate Iridium 192 source. Patients received 50 Gy by External beam irradiation (200cGy/fr, total 25 Fr) and 21 Gy by Intracavitary brachytherapy (7Gy/fr, 3fr). In control group, patients were treated by concurrent weekly cisplatin with external beam radiotherapy. In study group patients were treated by concurrent weekly paclitaxel with external beam radiotherapy.

GJMR-D Classification : NLMC Code: WP 470, WR 660



Strictly as per the compliance and regulations of:



Study of Weekly Paclitaxel with Concurrent Radiotherapy V/S Weekly Cisplatin with Concurrent Radiotherapy in Advanced Cervix Carcinoma

Dr. Prashant Rajendra Kumbhaj ^α, Dr. Rameshwaram Sharma ^σ, Dr. Aseem Rai Bhatnagar ^ρ & Dr. Peeyush Kumar Saini ^ω

Abstract- Purpose: To evaluate the Response and Efficacy of weekly paclitaxel with concurrent radiotherapy in comparison to weekly cisplatin with concurrent radiotherapy in advanced cervical cancer patients.

Methods: Patients coming to the department of radiotherapy S.M.S Hospital were included in the study. 50 patients of advanced squamous cell carcinoma of cervix FIGO stage III & IV were selected. In the study and control group, patients were given 5 weeks of external beam radiotherapy followed by brachytherapy. External beam irradiation each of 200cGy/fr, on linear accelerator by parallel opposed – anterior and posterior fields was given. External beam irradiation was followed by Intracavitary brachytherapy 700cGy/week for 3 weeks each with High Dose Rate Iridium 192 source. Patients received 50 Gy by External beam irradiation (200cGy/fr, total 25 Fr) and 21 Gy by Intracavitary brachytherapy (7Gy/fr, 3fr). In control group, patients were treated by concurrent weekly cisplatin with external beam radiotherapy. In study group patients were treated by concurrent weekly paclitaxel with external beam radiotherapy.

Results: At end of the completion of treatment 76% in the study group and 72% in the control group were found to have complete response to the treatment, 16% and 20% were found to have partial response in the study and control group respectively. Statistical analysis was done by chi square test. After 6 months of completing the treatment the result was equivalent in both the groups, 84% had complete response and 8% had partial response. Though the difference in tumour response was not statistically significant ($\chi^2 = .13$, $p = 0.94$), the rate of reduction in tumor size was found faster in study group at the end of the treatment, complete response was little more in the study group. The vomiting during the treatment is statistically significantly more in the control group ($\chi^2 = 23.548$, $p < 0.0001$). Total leukocyte count was significantly decreased more in study group compared to control group ($\chi^2 = 9.8$, $p = 0.0106$).

Conclusions: The overall response with the use of paclitaxel, which is the study arm, are equal to those with cisplatin. Though not significant, but tumor response was more in paclitaxel arm compared to cisplatin arm.

Authors α ρ ω: Department of Radiotherapy & Oncology SMS Medical College & Attached hospitals, Jaipur, Rajasthan India.
e-mails: drprashantkumbhaj@yahoo.com, piyushasini622@gmail.com, aseemrai123@gmail.com

Author σ: Department of pathology, Government medical college, Surat, Gujarat India. e-mail: cancertherapy@yahoo.co.in

I. INTRODUCTION

Cancer is the third biggest killer in India. India recorded 9.8 lakh new cases of cancer in the year 2010, an increase of about 80,000 new cases as compared to 2009. In India, four lakhs die of cancer every year. There is data that shows 25 lakh cancer patients in the country at any point in time. By 2015, the number of new cases in India is expected to cross 15 lakhs (1). Among Indian males, lung, head and neck cancer is the most common and among women, breast and cervical cancer is the most frequent. Carcinoma of the Uterine Cervix is the most common malignancy to affect females of developing countries. It ranks second in incidence among females, after Cancer Breast. Worldwide, approximately 5 lakh new cases are identified each year and about 2.33 lakh patients die of the disease (2). In developing countries, it accounts for about 3.4 lakh new cases & 1.6 lakh deaths every year (3). In India about 1.82 lakh new cases & 80,000 deaths every year from this disease (4). An estimated 1,82,027 new cases and 77,096 deaths due to cervical cancer occurred in India in 2010, contributing to 29% and 30% of the global burden of cervical cancer incidence and mortality (5).

Patients with Cervical Cancer usually present with locally advanced disease that is clinically confined to the pelvis (FIGO stage IIB, III & IV) in which surgery has higher morbidity. Radiotherapy plays a major role in management of these patients. The limitation of radiotherapy in controlling pelvic diseases for locally advanced cervical cancers is that radiation doses required to treat large tumours in the setting of poor tumour oxygenation exceeds the limit of toxicity in normal tissue. This was the main reason for treatment failure supporting by the fact that about 70% of relapses have pelvic failure as the first sites (6). Many strategies have been made trying to improve outcomes in locally advanced diseases such as uses of hypoxic cell sensitizers, hyperbaric oxygen, neutron therapy, and hyper-fractionation. However, results of those mentioned were found limited or unsuccessful (7). After a 1999

National Cancer Institute (NCI) Clinical alert was issued, chemoradiotherapy has become widely used in treating women with cervical cancer. Cisplatin is considered the most active cytotoxic agent and the drug of choice for concurrent chemoradiation. Meta-analysis studies have also demonstrated improved local control rates and survival with cisplatin-based chemotherapy concurrent to radiation therapy (RT). Although many prospective studies had shown that CTRT with cisplatin-based chemotherapy clearly improve the outcome of patients with carcinoma of the cervix, many patients treated on these protocols continue to fail in the pelvis and at distant sites [8]. In addition, one intergroup study using weekly concurrent cisplatin with radiotherapy for patients with carcinoma of the cervix could not demonstrate a beneficial effect of CTRT over standard RT alone [9]. This non-superiority finding was attributed to many factors like possible enrollment of patients with paraaortic lymph nodes, and an imbalance among randomization groups for known prognostic factors such as anemia [10]. These facts have lead many groups to investigate other drugs for CTRT like paclitaxel in an attempt to improve on what can be achieved by concurrent cisplatin [11]. Paclitaxel is a taxane chemotherapy drug that was found to have significant activity in solid tumors especially epithelial ovarian cancer, lung, and breast cancer. Preclinical studies have shown a radiosensitizing effect of paclitaxel in human cervical cancer cell lines. (12)

At Our Centre, the treatment for locally advanced cervical cancer is concurrent chemoradiotherapy; the protocol is cisplatin 50 mg/m² once a week for 5 weeks concurrent with pelvic radiation. Radiation is administered with cobalt-60 and HDR Brachytherapy. Paclitaxel is also radiosensitizing agent. We conducted study to investigate the efficacy and response of Paclitaxel with concurrent radiotherapy versus Cisplatin with concurrent radiotherapy.

II. METHOD AND MATERIALS

Patients coming to the department of radiotherapy & cancer treatment centre, SMS medical college & hospital jaipur were included in the study. After proper evaluation of the Biochemical and Hematopoietic profile, 50 patients were randomly divided into study and control group.

III. INCLUSION CRITERIA

1. Patients with newly diagnosed cases of Squamous Cell Carcinoma Cervix FIGO stage-III & IV were taken for study and control group.
2. Patients with advanced & inoperable cervix cancer.
3. Age between 18 to 70 years.
4. ECOG (eastern cooperative oncology group) performance status. 0 to 2.

IV. EXCLUSION CRITERIA

1. Patients previously treated with radiotherapy, Surgery, Chemotherapy.
2. Any histology other than squamous cell carcinoma.
3. Poor performance status ECOG 3 & 4.
4. Any uncontrolled intercurrent illness.
5. Pt who don't give consent for the study.
6. Class III & IV cardiac failure according to NYHA classification.
7. Pregnant & lactating women.

In the study and control group, patients were given 5 weeks of external beam radiotherapy followed by brachytherapy. External beam irradiation each of 200cGy/fr, on linear accelerator by parallel opposed – anterior and posterior fields was given. External beam irradiation was followed by Intracavitary brachytherapy 700cGy/week for 3 weeks each with High Dose Rate Iridium 192 source. Patients received 50 Gy by External beam irradiation (200cGy/fr, total 25 Fr) and 21 Gy by Intracavitary brachytherapy (7Gy/fr, 3fr). In control group, patients were treated by concurrent weekly cisplatin with external beam radiotherapy. In study group patients were treated by concurrent weekly paclitaxel with external beam radiotherapy.

Response was assessed as per the RECIST Criteria 1.1(13). The results of study group were analyzed & compared with control group in terms of various aspects like side effects, tumor response, & local disease status. The data thus collected were analyzed by using Chi-square test for co-relation.

V. RESULTS

Treatment Response at The end of Study

Table a shows treatment response of disease of patients using RECIST version 1.1 criteria while on treatment and subsequent follow up.

After 2nd week of initiating treatment, 9 (36%) patients of study group and 8 (32%) patients of control group had partial response, 16 (64%) patients of study group and 17 (68%) patients of control group had stable disease ($\chi^2 = 0.089$, $p = 0.76$).

After 4th week of initiating treatment, 2(8%) patients of study group and 1(4%) patients of control had complete response, 18 (72%) patients of both study & control group had partial response and 5 (20%) patients of study group and 6 (24%) patients of control group had stable disease ($\chi^2 = .42$, $p = 0.80$).

After 5th week of initiating treatment, 4(16%) of control group and 5(20%) of study group had complete response, 17 (68%) patients of control group and study group had partial response 4(16%) patients of control group and 3 (12%) patients of study group had stable disease ($\chi^2 = .25$, $p = 0.88$).

At end of treatment, 18(72%) of control & 19(76%) of study group had complete response, 5 (20%) patients of control group and 4(16%) patients of study group had partial response 2 (8%) patients of study group and control group had stable disease ($\chi^2 = .13$, $p = 0.94$).

At 1st month of follow up, 19 (76%) patients of control group and 20 (80%) patients of study group had complete response 4 (16%) patients of control group and 3(12%) patients of study group had partial response and 2 (8%) patients of study group and control group had stable disease ($\chi^2 = .16$, $p = 0.91$).

At 3rd month of follow up, 21 (84%) patients of study group and control group had complete response, 3 (12%) patients of study group and control group had partial response and 1 (4%) patients of study and control group had stable disease ($\chi^2 = 0$, $p = 1$).

At 6th month of follow up, 21 (84%) patients of study group and control group had complete response, 4 (16%) patients of study group and control group had partial response and none patients of study group and control group had stable disease ($\chi^2 = 0.14$, $p = 0.699$).

Table a : Treatment Response Evaluation For Disease Control

TREATMENT RESPONSE	STUDY GROUP					CONTROL GROUP					STATISTICS	
	Complete Response	Partial Response	No Change	Progressive Disease	TOTAL	Complete Response	Partial Response	No Change	Progressive Disease	TOTAL	χ^2	P

2 nd week	0	9	16	0	25	0	8	1 7	0	25	0. 0 8	0.76
4 th week	2	18	5	0	25	1	18	6	0	25	0. 4 2	0.80
5 th week	5	17	3	0	25	4	17	4	0	25	0. 2 5	0.88
End of Treat ment	19	4	2	0	25	18	5	2	0	25	0. 1 3	0.94
1 st month	20	3	2	0	25	19	4	2	0	25	0. 1 6	0.91
3 rd month	21	3	1	0	25	21	3	1	0	25	0 0	1
6 th month	21	4	0	0	25	21	4	0	0	25	0. 1 4	0.69 9

Acute Toxicity During Treatment

The incidence of nausea during treatment in both the groups are summarised in the table b. As per RTOG criteria Table b shows that in the study group 10 (40%) of the patients had Grade 1 Reactions as compared to 6 (24%) in the control group whereas 2(8%) in the study group and 7 (28%) in the control

group had Grade 2 Reaction and the incidence of Grade 3 was 2 (8%) and 5 (20%) in the study group and control group respectively($\chi^2=5.9$, $p=0.11$).

The vomiting during the treatment is statistically significantly more in the control group. 22(84%) patient in study group had Grade 0-1 vomiting and only 5(20%) had in study group, whereas only 3(12%) had grade 2-3

vomitting in study group, comparing to 11(44%) had grade 2 and 9(36%) had grade 3 vomiting in control group ($\chi^2=23.548, p= <0.0001$).

The haematological toxicity is represented in the table b in three sub heads- haemoglobin, Total Leucocytes count and Absolute Neutrophil count. 4 (16%) and 5(20%) had grade 1, 21(84%) and 20(80%) had grade 2 haemoglobin level in the study group and the control group ($p=1$).

Total leucocyte count in the table b shows that study group had more grade 2 and 3 toxicity {6(24%) and 4(16%)} comparing to control group having 2(8%) grade 2 toxicity & 1(4%) having grade 3 toxicity. 8(32%) in study group and 19(76%) group had grade 0 toxicity, 7(28%) and 3(12%) had grade 1 toxicity in study and control group respectively. This difference is statistically significant, showing more toxicity in the study group compared to control group ($\chi^2=9.8, p=0.0106$), absolute neutrophil count shows the same pattern of toxicity profile. 10(40%) and 20(80%) had grade 0, 8(32%) and 3(12%) had grade 1, 4(16%) and 1(4%) had grade 2 toxicity in the study and control group respectively. 3 (12%) developed grade 3 toxicity in the study group and 1(4%) in control group. This difference was statistically significant more in the study group compared to control group ($\chi^2=8.4, p=0.03$).

There was no nephrotoxicity found in 24(96%) in study group and 19(76%) in the control group, while 1(4%) and 4(16%) were found to have grade 1 toxicity in the study and control group respectively and 1(4%) was found to have grade 2 toxicity in control group ($p=.11$)

Table b

VI. DISCUSSION

The clinical feasibility of concurrent RT and paclitaxel was tested in phase I trials and a maximum tolerated dose (MTD) of 50 mg/m² per week concurrently with radiation therapy was established. In addition, the clinical efficacy of paclitaxel has been tested in phase II and III studies for metastatic and recurrent cervical cancer with objective response rates ranging between 36 and 47%. In all these studies paclitaxel was used in conjunction with either cisplatin (4/7 studies) or carboplatin (3/7 studies) but was never used alone for CRT. The majority of these studies was phase I (4/7 studies), with one study being a combined phase I/II study conducted by the GOG [60]. The number of patients enrolled in these studies varied between 8 and 35 patients and the rates of progression free survival ranged between 39 and 88%. The dose limiting toxicity was primarily neutropenia in 4 studies [14,15] or diarrhea (14).

The present study was carried out on 50 histopathologically confirmed newly diagnosed cases of Squamous cell Carcinoma cervix Stage IB to IVA. These cases were registered for treatment in department of

Radiotherapy at S.M.S hospital jaipur from December 2011 to June 2012.. In this study there was no statistically significant toxicity between the study group and control group for acute skin reaction, nausea during treatment, acute diarrhea, haemoglobin changes during treatment and nephrotoxicity. The statistically significant neutropenia was found in study group and vomiting in control group.

The tumour response in this study was evaluated in the 2nd, 4th, 5th week and at the end of treatment. After the completion, patients were evaluated every month till 6th month. Reduction in the tumour size was seen comparatively more in the study group and rendering them fit for brachytherapy earlier than the control group. In the 4th week of treatment 8% in the study group and 4% in the control group had complete response and in the 5th week 20% in the study group and 16% in the control group had complete response.

At end of the completion of treatment 76% in the study group and 72% in the control group were found to have complete response to the treatment, 16% and 20% were found to have partial response in the study and control group respectively. After 6 months of completing the treatment the results were equivalent in both the groups, 84% had complete response and 8% had partial response. Though the difference in tumour response was not statistically significant, the rate of reduction in tumour size was found faster in study group at the end of the treatment complete response was little more in the study group. So paclitaxel can be used as a radiosensitizer concurrently with radiotherapy in advanced inoperable patients in whom renal functions are not normal due to (1)hydronephrosis caused by advance disease,(2)Chronic renal disease.

VII. CONCLUSIONS

Our study provides a direct comparison between cisplatin and paclitaxel used as weekly concurrent chemotherapy with definitive radiation for advanced carcinoma of the cervix. Our data indicate that the overall response with the use of paclitaxel, which is the study arm, are equal to those with cisplatin. Though not significant, but tumor response was more in paclitaxel arm compared to cisplatin arm. However, the results were encouraging and it shall require larger number of patients and longer follow up in order to arrive at a concrete conclusion as far as disease free survival, cause specific survival, pelvic control rate, and long term sequel or complications are concerned.

VIII. BIBLIOGRAPHY

1. Kounteya Sinha, TNN Nov 10, 2011, 04.40AM IST (TOI) Hoskins WC, Perez CA, Young RC, eds. Principles and practice of Gynecologic oncology, 4th ed. Philadelphia: Lippincott Williams & Wilkins, 2005.

2. Kvale G, Heuch I, Ursin G. Reproductive factor and risk of cancer of the uterine corpus: A prospective study. *Cancer Res* 48: 6217, 1988.
3. Sankaranarayanan R, Esmy PO, Raj Kumar Rand Muwonge R: Visual Screening and Cervical Cancer. *The National Medical Journal of India*. Volume – 20, Number 5, 370: 398 -406, 2007.
4. The Times of India 2012-10-21: Dr. Surendra S. Shastri Principles and Practice of Gynecologic Oncology, 4e Asian Pacific Journal of Cancer Prevention, Vol 11, 2010
5. 7.Asian Pacific Journal of Cancer Prevention, Vol 11, 2010.
6. Keys HM, Bundy BN, Stehman FB. et al. Cisplatin, radiation and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for Bulky stage IB cervical carcinoma. *N Engl J Med*. 1999; 340:1154–61.doi: 10.1056/NEJM199904153401503.
7. Pearcey R, Brundage M, Drouin P. et al. Phase III trial comparing radical radiotherapy with and without cisplatin chemotherapy in patients with advanced squamous cell cancer of the cervix. *J Clin Oncol*. 2002;20(4):96672doi:10.1200/JCO.20.4.966. 7.
8. Rose PG. Concurrent chemoradiation for locally advanced carcinoma of the cervix: where are we in 2006? *Ann Oncol*. 2006; 17(Suppl 10):x224–9. doi: 1093/annonc/mdl264.
9. Chen MD, Paley PJ, Potish RA. et al. Phase I trial of Taxol as radiation sensitizer with Cisplatin in advanced cervical cancer. *Gynecol On-col*. 1997; 67:131–6. doi:1006/gyno.1997.4851
10. DiSilvestro PA, Walker JL, Morrison A. et al. Gynecologic Oncology Group. Radiation therapy with concomitant paclitaxel and cisp-latin chemotherapy in cervical carcinoma limited to the pelvis: a phase I/II study of the Gynecologic Oncology Group. *Gynecol Oncol* . 2006;103(3): 1038–42. doi: 1016/j. ygyno. 2006.06.017.
11. New response evaluation criteria in solid tumours.



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Perforated Duodenal Diverticulum: Diagnosis and Management

By Zhou Yaw LOO & Asiri Arachchi

Abstract- Duodenal diverticula occur in up to 20% of the general population and are generally asymptomatic. Complications such as perforation are exceedingly rare, with only 101 cases reported in the literature.¹ Early diagnosis is a challenge and clear guideline for the management of this rare entity is still lacking. We present a case of a 50 year-old man who was admitted to our institution with a perforated duodenal diverticulum. The diagnosis was achieved by contrast CT imaging of his abdomen and he was subsequently managed conservatively with nasogastric decompression and broad-spectrum antibiotics.

GJMR-D Classification : NLMC Code: WS 141



Strictly as per the compliance and regulations of:



Perforated Duodenal Diverticulum: Diagnosis and Management

Zhou Yaw LOO^a & Asiri Arachchi^c

Abstract- Duodenal diverticula occur in up to 20% of the general population and are generally asymptomatic. Complications such as perforation are exceedingly rare, with only 101 cases reported in the literature.¹ Early diagnosis is a challenge and clear guideline for the management of this rare entity is still lacking. We present a case of a 50 year-old man who was admitted to our institution with a perforated duodenal diverticulum. The diagnosis was achieved by contrast CT imaging of his abdomen and he was subsequently managed conservatively with nasogastric decompression and broad-spectrum antibiotics.

I. CASE REPORT

A 50-year-old man presented to our centre with a 1-day history of sudden-onset epigastric and right upper quadrant pain, associated with nausea and vomiting. There was no fever or any other infective symptoms. His past medical history was significant for Type 2 diabetes, hypertension and hypercholesterolaemia. On examination, he was normotensive, tachycardic and apyrexial. Localised epigastric tenderness with associated voluntary guarding was appreciated. Rectal examination did not reveal any presence of blood or melaena. Laboratory data showed a haemoglobin of 162 g/L and white cell count of $17.1 \times 10^9/L$. CT abdomen with contrast demonstrated a thickened duodenal wall with a 2-centimetre diverticulum in the region of the duodenal papilla (Figure 1). This was associated with peri-duodenal fat stranding and extraluminal air posterior and lateral to the duodenal wall (Figure 1 & 2). After initial fluid resuscitation, a nasogastric tube was inserted and he was commenced on broad-spectrum antibiotics. Total parental nutrition was initiated and a repeat CT few days later demonstrated significantly smaller para-duodenal collection. He was slowly progressed to soft diet and was discharged 12 days after initial presentation with a plan for gastroscopy in 6 weeks.

II. DISCUSSION

Duodenum is the second most common location to be affected by diverticula, after the colon.^{2,3,5} They are typically pseudo-aneurysm, as not all layers of the bowel wall are affected.^{4,5,6} About 75% of them are located in the second part of the duodenum, usually within 2.5cm of the ampulla of Vater. Most of them (88%)

are located along the pancreatic or mesenteric border. The incidence of duodenal diverticula has been reported between 1-5% in upper GI studies and up to 22% at autopsies.¹ Duodenal diverticula are mostly asymptomatic, however complications such as bleeding, inflammation and perforation can develop in 5% of patients.⁵ Perforation represents the rarest but most serious complication, with mortality of up to 13%.¹

Perforated duodenal diverticulum poses a diagnostic challenge as there is no pathognomonic clinical features. Of all the 101 cases reported in literature, only 13 of the cases achieved accurate pre-operative diagnosis.¹ The symptoms are mostly non-specific and could mimic conditions such as peptic ulcer disease, acute cholecystitis and pancreatitis.⁷ Majority of the patients presented with acute-onset epigastric or right upper quadrant pain associated with nausea and vomiting,^{3,4,5} while some experienced chronic abdominal pain with anorexia and malaise.³ The diagnosis is by exclusion and requires clinician to possess a high index of suspicion.

Contrast-enhanced CT is a valuable diagnostic modality due to its ability to detect small amount of extraluminal air and delineate the retroperitoneal anatomy.³ It can even demonstrate the presence of a diverticulum in some cases. Typical radiographic findings include thickened bowel wall, fat stranding and extraluminal air or fluid.^{4,5} However, these are not diagnostic as perforated viscus close to an asymptomatic duodenal diverticulum can give a similar appearance.^{2,7} Post-bulbar duodenal ulcer for instance, can produce the clinical and radiological presentation of a perforated duodenal diverticulum.² Upper gastrointestinal series may be employed as part of the investigation but previous literature fail to prove its effectiveness.^{2,3} This could be explained by cases of diverticulum with tight neck or the presence of a calculus.³ Forward-viewing endoscopy can be useful in excluding pathologies such as peptic ulcer, but it has been shown to have a low-predictive value of diagnosing duodenal diverticulum.⁵ Side-viewing duodenoscopy may be the tool of choice,⁵ with Leinoven et al and Jin et al finding duodenal diverticulum in 7.1% and 24.6% of all patients who underwent ERCP, respectively.^{8,9} One rationale behind this is that forward-

viewing endoscopy may not reliably assess the concavity of the descending duodenum close to the ampulla of Vater, where most of the diverticulum lies.⁵

In cases of perforated duodenal diverticulum, the treatment of choice seems to be transduodenal diverticulectomy with two-layer closure.¹ Bypass procedure such as gastrojejunostomy and tube duodenostomy may be required.² Due to the close proximity of distal common bile duct and the ampulla of Vater, the surgeon must proceed with utmost care to avoid iatrogenic injury.¹ Complications associated with repair of perforated diverticulum include duodenal fistula, intra-abdominal abscess, sepsis and pancreatitis, with duodenal fistula being the commonest (20%).¹ In addition, laparoscopic¹⁰ and endoscopic¹¹ treatment of perforation have also been described. Conservative management of perforation is a feasible option provided

that the patient has mild symptoms and has radiographic evidence of a contained leak.⁶ Treatment involves nasogastric decompression, broad-spectrum antibiotics with or without percutaneous drainage of subsequent abscess formation.⁴ There have been reports in literature of successful outcome with such treatment.^{1,4,6} However, should there be clinical deterioration, surgical intervention is obviously warranted.

In conclusion, perforation represents the rarest complication of duodenal diverticulum and early diagnosis is challenging. Optimal treatment of this condition has not been well-established and may be guided by the patient's clinical condition. We present a case of a perforated duodenal diverticulum diagnosed by contrast-enhanced CT and managed successfully with a conservative approach.



Figure 1 : A 2-centimetre diverticulum can be appreciated in the regional of duodenal papilla




Figure 2 : Axial view showing extraluminal gas and induration of abdominal fat around the duodenum

REFERENCES RÉFÉRENCES REFERENCIAS

1. Duarte B, Nagy K, Cintron J. Perforated duodenal diverticulum. Br J Surg 1992; 79: 877-881.
2. Huang R, Romano A, Stone M, et al. Diagnosis and treatment of a perforated duodenal diverticulum. Emerg Radiol 2007; 13: 285-287.
3. Sakurai Y, Miura H, Matsubara T, et al. Perforated duodenal diverticulum successfully diagnosed preoperatively with abdominal CT scan associated with upper gastrointestinal series. J Gastroenterol 2004; 39: 379-383.
4. Martinez-Cecilia D, Arjona-Sanchez A, Gomez-Alvarez M, et al. Conservative management of perforated duodenal diverticulum: A case report and review of the literature. World J Gastroenterol 2008; 14: 1949-1951.
5. Schnueriger B, Vorbuerger S, Banz V, et al. Diagnosis and management of the symptomatic duodenal diverticulum: A case series and a short review of the literature. J Gastrointest Surg 2008; 12: 1571-1576.
6. Marhin W, Amson B. Management of perforated duodenal diverticula. Can J Surg; 48: 79-80.
7. Ames J, Federle M, Pealer K. Perforated duodenal diverticulum: Clinical and imaging findings in eight patients. Abdom Imaging 2009; 34: 135-139.
8. Leivonen M, Halttunen J, Kivilaakso E. Duodenal diverticulum at endoscopic retrograde cholangiopancreatography, analysis of 123 patients. Hepatogastroenterology 1996; 43: 961-966.
9. Jin T, Ho G, Eun Y, et al. Incidence of juxtapapillary duodenal diverticula is higher in brown pigment stone but not in black pigment or cholesterol stone. Gastroint Endosc 2006; 63: 297.
10. Lee H, Hong J, Oh S, et al. Laparoscopic diverticulectomy for a perforated duodenal diverticulum: a case report. J Laparoendosc Adv Surg Tech A 2010; 20: 757-760.
11. Eeckhout G, Vanstiphout J, Van Pottelbergh I, et al. Endoscopic treatment of a perforated duodenal diverticulum. Endoscopy 2000; 32: 991-993.

This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Characterization of Hippocampus in Epileptic Sudanese Children using MRI

By Caroline Edward Ayad, Mohammed Abdalla Bashir Abdalla,
Mohammed .E.M. Garelnabi & Elsafi Ahmed Abdalla

Sudan University of Science and Technology, Sudan

Abstract- Seizures are the most noticeable clinical manifestations of epilepsy; measurement of the hippocampus on magnetic resonance (MR) images of the brain has been useful for the assessment of patients with seizures. Therefore this study aimed to diagnose subjects who had developed epilepsy by measuring the RT and LT hippocampus area and volume considering epilepsy duration, seizure frequency, EEG results, T2 signal intensity, treatment, and children ages as well as to correlate the MRI findings with EEG findings. This Study was conducted at Gezira University- MRI Diagnostic center –Khartoum –Sudan during the period from 2010 up to 2011 using Siemens 0.5 Tesla MRI machine. A total of 29 Sudanese children of both genders with clinical history of generalized seizures, and 17 normal children who were selected as control group at different ages were involved in this study. Children with head injury or organic lesion found on MR images were excluded.

Keywords: *epilepsy, siezers, hippocampus.*

GJMR-D Classification : *NLMC Code: WS 141, WN 180*



Strictly as per the compliance and regulations of:



Characterization of Hippocampus in Epileptic Sudanese Children using MRI

Caroline Edward Ayad ^α, Mohammed Abdalla Bashir Abdalla ^σ, Mohammed .E.M. Garelnabi ^ρ & Elsafi Ahmed Abdalla ^ω

Abstract- Seizures are the most noticeable clinical manifestations of epilepsy; measurement of the hippocampus on magnetic resonance (MR) images of the brain has been useful for the assessment of patients with seizures. Therefore this study aimed to diagnose subjects who had developed epilepsy by measuring the RT and LT hippocampus area and volume considering epilepsy duration, seizure frequency, EEG results, T2 signal intensity, treatment, and children ages as well as to correlate the MRI findings with EEG findings. This Study was conducted at Gezira University- MRI Diagnostic center –Khartoum –Sudan during the period from 2010up to2011using Siemens 0.5 Tesla MRI machine. A total of 29 Sudanese children of both genders with clinical history of generalized seizures, and 17 normal children who were selected as control group at different ages were involved in this study. Children with head injury or organic lesion found on MR images were excluded.

The hippocampus was characterized in the coronal MR images; the area and volume of the right and Left sides were found to be reduces significantly in the diseased sample comparing with the control group; there was relationship between patient's age and seizure duration and frequency as well as hippocampus area and volume. Significant differences were detected in the hippocampus area and volume in children who have got treatment and none treated children.T2 weighted images showed hyper signal intensity in the hippocampuses of the diseased sample. No significant differences were detected in males and females measurements. EEG failed in diagnoses of 12 (41.4%) cases of the sample, and new formulas were established for prediction of EEG findings regarding the hippocampus area and epilepsy duration.MRI has a great value in characterization hippocampus in epileptic Sudanese children.

Keywords: epilepsy, siezers, hippocampus.

1. INTRODUCTION

Epilepsy is a chronic brain disorder that is characterized by repeated spontaneous seizures, and increased risk of mortality.[Wilner AN et al.,2010] Epilepsy is one of the most common neurologic disorders encountered in clinical practice. Epilepsy is more common among the elderly, higher incidence in men than women.

[Hesdorffer DC et al., 2011, Banerjee PN et al .,2009], more common among racial or ethnic minorities or among low-income populations. [Banerjee PN et al., 2009].

Seizures are the most obvious clinical manifestations of epilepsy; it refers to the changes in physical findings that occur in relation to abnormal electrical activity within the brain [Fisher RS et al., 2005] The electroencephalogram (EEG), can measures changes in brain electrical activity, and identify brain regions where seizures are most likely to originate. More recently, imaging technologies such as magnetic resonance imaging (MRI), has become indispensable tool in evaluating patients with epilepsy and other seizure disorders, because of it's superior soft tissues contrast, and ability of viewing variety of planes, providing greater sensitivity and accuracy for the identification of underlying lesions in patients with epilepsy .According to guidelines from the American College of Radiology (ACR), MRI is considered appropriate for all epilepsy evaluations, and become the standard for brain imaging in patients with seizures.[www.acr.org 2011].

A systematic approach to the evaluation of seizure disorders is essential to ensure that patients receive appropriate treatment and to rule out the many other medical conditions that may resemble epilepsy [Noe KH et al., 2011].

Common MRI protocols used in the evaluation of epilepsy include volumetric T1-weighted sequencing; proton-density, T2-weighted and fluid-attenuated inversion recovery (FLAIR) sequences in oblique coronal and axial planes; and gradient echo sequences. [Duncan JS. ,2010].

Measuring hippocampus volume with MRI has provided important information about several neuropsychiatric disorders, smaller hippocampus volumes have been reported in epilepsy [Geuze et al., 2009].

Measurement of hippocampus volumes has been found to be the most accurate approach to evaluate the hippocampus pathology [Bronen RA.,1991]. This has found its greatest clinical application in the assessment of patients with seizures [Ashtari M et al., 1991, Jack CR.,1990]. Other factors may also help identify hippocampal sclerosis: quantitative evidence of increased T2 values; and visual assessment of the disruption of internal architecture, volume loss, and

Author α: Sudan University of Science and Technology-College of Medical Radiological Science. e-mail: carolineayad@yahoo.com

Authors σ ρ ω: Gezira University- MRI Diagnostic center, Khartoum – Sudan. e-mails: mohammedbasheer@yahoo.com, mohamedgarelnabi@yahoo.com, dr.elsaffi@gmail.com

intensity elevation [Jackson GD., 1990, Kuzniecky R., 1987].

This study aimed to diagnose subjects who had developed epilepsy by measuring the RT and LT hippocampus area and volume considering epilepsy duration, seizure frequency, EEG results, T2 signal, treatment, and children ages as well as to correlate the MRI findings with EEG findings.

II. MATERIAL AND METHODS

This Study was conducted at Gezira University-MRI Diagnostic center-Khartoum-Sudan during the period from 2010up to2011.

MRI machine: Siemens 0.5 Tesla

Patients: A total of 29 children in both genders with clinical history of generalized seizures, and 17 were normal as control group at different ages were involved in this study. The ethics and research committee approved the study.

Exclusion criteria: Subjects who were normal, with head injury or organic lesion found on MR images were excluded.

MRI technique and Hippocampus area and volume measurements

MRI Brain was done first axial T1, T2, FLAIR and sagittal images; it is used as screening for presence of

mass lesion or other abnormalities. Sagittal image was used as a guide to obtain perpendicular plane to temporal lobe. T1, T2 of coronal 3mm slice thickness, images perpendicular to the long axis of the temporal lobe were taken.

Volume measurement

The whole hippocampal volume was measured. Measurement of hippocampal body and tail included subicular complex, the hippocampus proper, the dentate gyrus, the alveus and hippocampal fimbria. ROI-manually traced with computer mouse, right and left sides were measured individually. The measures were done three times for each ROI and the average was taken. Summation of the average of each slice = the total area of hippocampus for each side.

Volume of the hippocampal complex (cm3)=total area of the hippocampal complex (cm2) X the slice thickness (cm)

III. RESULTS

The 29 subjects studied consist of 11(37.9%) males and 18(62.1%) females. The mean ages of the subjects were 14.79 ± 6.01 years. The mean age of the seizure onset was found to be 12.4 ± 5.7 and frequency of 6.6 ± 4.35 .

Table 1 : The mean and standard deviation of the hippocampus volume and area measurements for epileptic patients

	Area (RT) Cm ²	Volume (RT) cm ³	Area (LT) Cm ²	Volume (LT) cm ³
Mean	2.75	2.47	2.62	2.36
Std. Deviation	±0.23	±0.21	±0.21	±0.19

Table 2 : Shows the frequency of epilepsy duration in the sample

Epilepsy duration (minutes)	Frequency	Percent%
2	7	24.1
3	5	10.3
4	5	13.8
5	5	6.9
8	1	3.4

9	3	10.3
11	1	3.4
13	2	6.9
Total	29	100.0

Table 3 : Shows the frequency of EEG results

		Frequency	Percent%
EEG	Consigned with MRI (negative epileptic findings)	7	24.1
	Adverse findings with MRI	12	41.4
	Consigned with MRI (positive epileptic findings)	10	34.5
	Total	29	100.0

Table 4 : Shows the frequency of T2 signal classified as (iso signal intensity& hyper signal intensity)

		Frequency	Percent%
Signal intensity	Iso intense	7	24.1
	Hyper intensity	22	75.9
	Total	29	100.0

Table 5 : Shows frequency of not treated & treated patients

		Frequency	Percent%
Treatment	Not treated	23	79.3
	Treated	6	20.7
	Total	29	100.0

Table 6 : Shows correlation between seizure frequency & Treatment

Model	Un-standardized Coefficients		Standardized Coefficients	Sig.
	B	Std. Error	Beta	
(Constant)	-.099	.124		.432
Seizures Frequency	.047	.016	.492	.007

*This table showed that the seizures frequency decreased by .099 according to the treatment

Table 7 : Shows correlation among EEG, Epilepsy duration, LT Hippocampus area in both genders

gender	Model	Un-standardized Coefficients		Standardized Coefficients	Sig.
		B	Std. Error	Beta	
Male	(Constant)	-3.229	1.924		.105
	Area (LT) Cm ²	1.570	.730	.388	.041
Female	(Constant)	-4.125	1.812		.032
	Area (LT) Cm ²	1.726	.676	.427	.017
	Epilepsy Duration	.101	.043	.397	.025

+here also EEG results can be related to the area (LT) and epilepsy duration as follow:

$$EEG = [(\text{area (LT)} \times 1.726) + (\text{epilepsy duration} \times 0.101)] - 4.13$$

Table 8 : Shows correlation among EEG, Epilepsy duration, RT Hippocampus area in both genders

gender	Model	Un-standardized Coefficients		Standardized Coefficients	Sig.
		B	Std. Error	Beta	
Male	(Constant)	-2.336	1.026		.126
	Area (RT) Cm ²	1.679	.680	.372	.036
Female	(Constant)	-3.236	1.789		.029
	Area (RT) Cm ²	1.624	.457	.496	.028
	Epilepsy Duration	.101	.043	.397	.025

*here also EEG results can be related to the area (RT) and epilepsy duration as follow:

$$EEG = [(\text{area (RT)} \times 1.624) + (\text{epilepsy duration} \times 0.101)] - 3.24$$

Table 9 : Shows T-Test Difference between male & female measurement of the area and Volume as mean, Std. Deviation and P-value

gender		N	Mean	Std. Deviation	Sig. (2-tailed)
Area (RT) cm ²	Male	11	2.7182	.21442	.607
	Female	18	2.7650	.24677	.595
Volume (RT) cm ³	Male	11	2.4473	.19267	.616
	Female	18	2.4883	.22150	.604

Area (LT) cm ²	Male	11	2.6173	.19571	.906
	Female	18	2.6272	.22862	.902
Volume (LT) cm ³	Male	11	2.3564	.17614	.891
	Female	18	2.3667	.20594	.887

Table 10 : Shows the measured area and volume (comparison between none treated & treated measurement in mean & Std. Deviation and *P*-value

Treatment		N	Mean	Std. Deviation	Sig. (2-tailed)
Area (RT) cm ²	No treated	23	2.7322	.22682	.504
	Treated	6	2.8050	.26569	.558
Volume (RT) cm ³	No treated	23	2.4583	.20252	.473
	Treated	6	2.5283	.24128	.534
Area (LT) cm ²	No treated	23	2.6061	.20701	.400
	Treated	6	2.6900	.24331	.464
Volume (LT) cm ³	No treated	23	2.3474	.18663	.409
	Treated	6	2.4217	.21885	.471

Table 11 : Shows the T2 signal status in normal & abnormal patients as mean Std. Deviation and *P*-value

T2 Signal		N	Mean	Std. Deviation	Sig. (2-tailed)
Area (RT) Cm2	Normal	7	2.6186	.15291	.093
	Abnormal	22	2.7882	.24073	.043
Volume (RT) cm3	Normal	7	2.3586	.14053	.097
	Abnormal	22	2.5091	.21580	.048
Area (LT) Cm2	Normal	7	2.5200	.14059	.143
	Abnormal	22	2.6564	.22413	.074
Volume (LT) cm3	Normal	7	2.2700	.12490	.145
	Abnormal	22	2.3923	.20225	.073

Table 12 : Shows mean and standard Deviation of the epileptic subjects and control group for RT and LT area and volume measurement (p -value at 0.05)

	Area (sampl e)	Area (contro l) (RT)	Volume (sampl e) (RT)	Volume (contro l) (RT)	Area (sam ple) (LT)	Area (contro l) (LT)	Volume (sampl e) (LT)	Volume (contro l) (LT)
	cm ²	cm ²	cm ³	cm ³	cm ²	cm ²	cm ³	cm ³
Mean	2.7	2.9	2.47	3.01	2.62	2.8	2.4	3.21
STDV	±.23	±.26	±.21	±.21	±.21	±.32	±.19	±.21
p - value	0.0423		0.059		0.0563		0.034	




Figure 1 : The relation between the Seizure duration and the age




Figure 3 : The relation between the Seizure frequency and the RT hippocampus area




Figure 2 : The relation between the Seizure frequency and the age




Figure 4 : The relation between the Seizure frequency and the LT hippocampus area




Figure 5 : The relation between the Seizure frequency and the RT hippocampus volume




Figure 6 : The relation between the Seizure frequency and the LT hippocampus Volume

IV. DISCUSSION

This study reported the MR findings in a group of patients with documented temporal lobe epilepsy. The diagnosis of temporal lobe epilepsy was based on typical clinical seizure characteristic. In epilepsy researches and in temporal lobe epilepsy in particular, hippocampal volumetry with MRI were mainly used in the determination of hippocampal atrophy and hippocampal sclerosis. The clinical features of this study of 29 epileptic children with temporal lobe epilepsy were in line with four cross sectional previous studies (L.E. Betting et al., 2006; Idil Cavus et al., 2008; Mai Hanamiya et al., 2009; Douglas E et al., 2010). Females were affected more often than males (18 females, 11 males). This gender results were consistently seen in our studies, supporting the hypothesis that the cause could be a biological phenomenon rather than a research entity. Table 1 showed the measurements of hippocampus area and duration and table 2 showed the duration of seizures (minutes) in the sample. Abnormalities on MR images were detected in 22 (75.9%), hippocampal sclerosis being the most

common lesion. EEG confirmation was present in 10 (34.5%) regarding 7 (24.1%) had normal EEG and 12 cases were reported adverse results from the MRI findings as presented in table (3). Overall agreement between the two observers was fair, indicating that hippocampal sclerosis and other lesion can be detected reliably on MR images. The reported MR features of hippocampal sclerosis included hippocampal atrophy, increased signal intensity on T2, loss of internal architecture of the hippocampus, and loss of signal intensity on T1 weighted images. In our study only the first two were used as criteria for diagnosis of hippocampal sclerosis.

No lesion was detectable on MR images of 7 (24.1%) concordant with 7 (24.1%) whom EEG result was normal as the T2 signal intensity was found to be iso intense and patients were diagnosed as normal but the abnormal signal was detected in 22 (75.9%) of the sample showing hyper signal intensity and were diagnosed as abnormal this was presented in table (4). The epilepsy was usually mild and responsive to medication this was presented in table (5). 23 out of 29 (79.3%) were on no treatment at the time of study. 6 out of 29 (20.7%) had good seizure control with a single anti-epileptic drug. The seizures frequency decreased by 0.99 according to treatment and the relation is significant between the frequency of seizures and cases administrated with drugs at p value 0.007; this was presented in table (6).

EEG results can be related to the left hippocampus area and epilepsy duration as: $EEG = [(\text{area (LT)} \times 1.726) + (\text{epilepsy duration} \times 0.101)] - 4.13$, where EEG results can be related to the right hippocampus area and epilepsy duration as: $EEG = [(\text{area (RT)} \times 1.624) + (\text{epilepsy duration} \times 0.101)] - 3.24$ as presented in tables (7) and (8).

No significant differences were detected between males and females mean measurements for the hippocampus area and volume as presented in table (9).

The hippocampal volume for treated patient were RT 2.53 ± 0.24 , LT 2.42 ± 0.22 and for none treated were RT 2.46 ± 0.2 , LT 2.35 ± 0.19 respectively table (10). The difference at T test (P value = 0.05) was found to be insignificant for both RT & LT as to be for RT (0.47, 0.53).

Table (4) showed that 7 (24.1%) in the group of patients had normal hippocampus T2 signal, 22 (75.9%) showed high signal intensity on T2. The hippocampal volume of normal T2 signal were RT 2.36 ± 0.14 , LT 2.27 ± 0.12 were the volume for high signal intensity RT 2.51 ± 0.21 , LT 2.39 ± 0.2 respectively table (11). T-test showed that the difference at (P -value 0.05) was to be insignificant for the normal and abnormal T2 signal for hippocampus volume as (0.97, 0.15) for RT and (0.5, 0.73) for LT respectively, but the difference was to be

significant for the normal T2 signal of the RT hippocampus area when compared with the abnormal T2 signal for the same area.

Previous investigations using volumetry showed that hippocampal atrophy is a reliable marker of hippocampal sclerosis (L.E. Betting et al., 2006; Mai Hanamiya et al., 2009). In an investigation of patients with temporal lobe epilepsy, 22 (75.9%) patients with hippocampal sclerosis pathologically proven had atrophy in the MRI evaluation.

In another volumetric evaluation of patients with idiopathic generalized epilepsy, no hippocampal atrophy had been detected but the hippocampal hyperintense T2 signal was present (Mai Hanamiya et al., 2009).

A significant reduction had been detected in the right and left hippocampus area and volume of the epileptic children when compared with the control group as presented in table (12).

The reduced hippocampus volumes described herein were slightly different from the typical findings of hippocampal sclerosis. Ipsilateral reduction of the fornix is associated with hippocampal atrophy in most of the patients with unilateral hippocampal atrophy. Hippocampal hyperintense T2 signal was present in 22 (75.9%) similar findings were mentioned by (L.E. Betting et al., 2006).

The justification is that a disturbance in the blood and oxygen supply during epileptic attacks causes destruction of hippocampal nerve cells and that the sclerosis, or scarring, is the end result of this process (Margerison J. H., et al 2010). The tendency for the damage to occur preferentially in certain parts of the hippocampus, is attributed either to a special sensitivity of these areas to lack of oxygen, or to their exceptional need for oxygen through increased cellular activity at a time when it is liable to be in short supply. This discriminating susceptibility may be due to variations in the chemical make-up of its different parts, (Purpura and Gonzalez-Monteaudo, 1960), the researchers suggested that these were the causes of changing the signal intensity in T2 weighted images.

EEG failed in diagnosing of 12 patients where MRI images showed a significant reduction and changing in the T2 signal intensity.

The study also noted that as the children ages increased the seizure duration was also decreased by 0.11 starting from 6.4 minutes, but the frequency increased by 0.14 as the children ages increased, this was presented in figure (1) and figure (2).

As the RT and LT hippocampus areas increased; the seizures frequency was also increased at R^2 0.04 and 0.039, this was presented in figures (3, 4) as well as the RT and LT hippocampus volume; it increased at R^2 0.038, 0.039 respectively and this was presented in the figures 5 and 6.

Regarding the results, the routine electroencephalogram (EEG) is often helpful in diagnosing epilepsy because it can detect the abnormal electrical discharges in the brain that indicate epilepsy. However, the EEG is very often normal in patients with proven epilepsy, so it cannot be used alone to exclude epilepsy. MRI is an excellent imaging modality used to characterize the hippocampus by measuring its area and volume as well as to evaluate the signal intensity as these articles are very important to evaluate patients with neurological disorders.

REFERENCES RÉFÉRENCES REFERENCIAS

1. American College of Radiology. ACR Appropriateness Criteria: Seizures and Epilepsy. Available at: http://www.acr.org/secondmainmenucategories/quality_safety/app_criteria/pdf/expertpanelonneurollogicimaging/epilepsydoc3.aspx. Reviewed 2011; Accessed August 12, 2011.
2. Ashtari M, Barr WB, Schawl N, Bogerts B. Three-dimensional fast Low angle shot imaging and computerized volume measurement Of the hippocampus in patients with chronic epilepsy of the temporal Lobe. *Am J Neuroradiol* 1991;12:941–947.
3. Banerjee PN, Filippi D, Allen Hauser W. The descriptive epidemiology of epilepsy-a review. *Epilepsy Res.* 2009;85:31-45.
4. Bronen RA, Cheung G, Charles JT, et al. Imaging findings in Hippocampal sclerosis: correlation with pathology. *Am J Neuroradiol* 1991;12:933–940.
5. Douglas E. Crompton Ingrid E. Scheffer-Isabella Taylor, Mark J. Cook, Penelope A. McKelvie, Danya F. Vears, Kate M. Lawrence, Jacinta M. McMahon, Bronwyn E. Grinton, Anne M. McIntosh Samuel F. Familial mesial temporal lobe epilepsy: a benign epilepsy syndrome showing complex inheritance Berkovic, Oxford Journals. *Medicine Brain*, Volume 133, Issue 11, Pp. 3221-3231.
6. Duncan JS. Imaging in the surgical treatment of epilepsy. *Nat Rev Neurol.* 2010;6:537-550.
7. Fisher RS, van Emde Boas W, et al. Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia.* 2005; 46:470-472.
8. Geuze E, Vermetten E, de Kloet CS, Hijman R, et al. (2009). Neuropsychological performance is related to current social and occupational functioning in veterans with posttraumatic stress disorder. *Depression and Anxiety* 26(1):7-15.
9. Hesdorffer DC, Logroscino G, Benn EK, et al. Estimating risk for developing epilepsy: a population-based study in Rochester, Minnesota. *Neurology.* 2011;76:23-27.

10. Jack CR, Sharborough FW, Twomey CK, et al. Temporal lobe Seizures: lateralisation with MR volume measurements of the hippocampal Formation. *Radiology* 1990;175:423–429.
11. Jackson GD, Berkovic SF, Tress BM, Kalnins RM, Fabinyi GCA, Bladin PF. Hippocampal sclerosis can be reliably detected by Magnetic resonance imaging. *Neurology* 1990;40:1869–1875.
12. Jullie W. Pan, Anne Williamson, Idil Cavus, Hoby P. Hetherington, Hitten Zaveri, Ognen A. C. Petroff, and Dennis D. Spencer Neurometabolism in human epilepsy. *Epilepsia*. 2008; 49(0 3): 31–41.
13. Kuzniecky R, de la Sayette V, Ethier R, et al. Magnetic resonance Imaging in temporal lobe epilepsy: pathological correlation. *Ann Neurol* 1987;22:341–347.
14. Luiz Eduardo Betting a, Susana Barreto Mory a, I' scia Lopes-Cendes b, Li Min Li , Marilisa M. Guerreiro a, Carlos A.M. Guerreiro a, Fernando Cendes a, MRI volumetry shows increased anterior thalamic volumes in patients with absence seizures, *a Epilepsy & Behavior* 8 (2006) 575–580.
15. Margerison J. H. , Corsi F, J. A. N. *Epilepsy And The Temporal Lobes A Clinical, Electroencephalographic And Neuropathological Study Of The Brain In Epilepsy, With Particular Reference To The Temporal Lobes* Downloaded From [Http: // Brain.Oxfordjournals.Org/By Guest](http://Brain.Oxfordjournals.Org/By Guest) On August 19, 2012.
16. Noe KH. Seizures: diagnosis and management in the outpatient setting. *Semin Neurol*. 2011;31:54–64.9. National Institute of Neurologic Disorders and Stroke. Brain basics. Available at: [http: // www. ninds.nih.gov/disorders/brain_basics/know_your_brain.htm](http://www.ninds.nih.gov/disorders/brain_basics/know_your_brain.htm). Accessed August 16, 2011.
17. Porpora, D. P., And Gonzalez-Monteagudo, O. (1960) /. *Neuropath, exp. Neurol*, 19, 421.
18. Ryota Serino, Yoichi Ueta, *Mai Hanamiya*, Masayoshi Nomura, Ilan A. Kerman . Circadian rhythms: interactions with seizures and *epilepsy* (2009), *Autonomic Neuroscience* Vol. 148, Issue 1, Pages 83.
19. Wilner AN, Benbadis SR. Epilepsy therapy in 2010. *Therapy*. 2010;7:445–448.





This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH
RADIOLOGY, DIAGNOSTIC, IMAGING AND INSTRUMENTATION
Volume 13 Issue 3 Version 1.0 Year 2013
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN : 0975-5888

Right Internal Carotid Agenesis: A Rare Case Anatomical, Neurosurgical and Radiological Perspective

By Dr. Ashfaq Ul Hassan, Dr. S habinul Hassan, Prof Nisha Kaul &
Prof Ghulam

SKIMS Medical College Bemina, India

Abstract- We present a case of 35 year old female with Right Internal carotid artery agenesis . The Axial and coronal T2- weighted MRI demonstrated an absent right internal carotid artery flow void. A subsequent magnetic resonance angiography showed absence of the right internal carotid artery. This finding was confirmed by magnetic resonance angiography of the cervical vessels, and axial computed tomography angiography showed agenesis of the right carotid artery. The literature reports such finding in association with other anomalies such as transsphenoidal encephaloceles and circle of Willis aneurysms. These associations were not observed in the present case. The patient remained asymptomatic.

GJMR-D Classification : NLMC Code: WL 368



Strictly as per the compliance and regulations of:



© 2013. Dr. Ashfaq Ul Hassan, Dr. S habinul Hassan, Prof Nisha Kaul & Prof Ghulam. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Right Internal Carotid Agenesis: A Rare Case Anatomical, Neurosurgical and Radiological Perspective

Dr. Ashfaq Ul Hassan ^α, Dr. Shabinul Hassan ^σ, Prof Nisha Kaul ^ρ & Prof Ghulam ^ω

Abstract- We present a case of 35 year old female with Right Internal carotid artery agenesis. The Axial and coronal T2-weighted MRI demonstrated an absent right internal carotid artery flow void. A subsequent magnetic resonance angiography showed absence of the right internal carotid artery. This finding was confirmed by magnetic resonance angiography of the cervical vessels, and axial computed tomography angiography showed agenesis of the right carotid artery. The literature reports such finding in association with other anomalies such as transsphenoidal encephaloceles and circle of Willis aneurysms. These associations were not observed in the present case. The patient remained asymptomatic.

I. INTRODUCTION

In the Medical OPD of SKIMS, 50 patients were worked upon for evaluation. These patients had neurological symptoms like headache, giddiness, weakness in limbs, loss of consciousness, dementia. After subjecting them for further evaluation one patient was found to have Neck Angiography revealing small right common carotid artery and no Internal carotid artery. MR Angiography revealed normal left petrous and Cavernous internal carotid. Right Middle cerebral artery and right anterior cerebral artery are likely supplied by anterior communicating artery and posterior communicating artery. In addition bilateral hyper intensity lesions were seen in heads of Caudate nuclei.

II. DISCUSSION

The Cerebral circulation is divided into Anterior circulation and posterior circulation. The internal carotid arteries form the anterior circulation and the vertebral arteries form the posterior circulation. These two meet at the Circle of Willis and most of the blood supply is derived from the internal carotid system. Embryologically the internal carotid is formed by the portion of the third arch artery distal to the bud along with the original dorsal aorta cranial to the attachment of third arch artery.

By inherent nature Agenesis of Carotid arteries is a rarity with not so frequent cases reported worldwide. It is mostly asymptomatic because of the fact of compensation by Blood flow through Circle of Willis. Internal carotid agenesis occurs when one of the blood vessels which supply blood to the brain does not develop (agenesis). It is rare, occurring in less than 0.01% of people.¹ Usually there is a pair of internal carotid arteries, one on the left side and one on the right side; agenesis occurs on the left side three times more frequently than on the right side.²

However, if symptoms occur, they may include headache, blurred vision, paralysis of some of the nerves in the head (palsy), epilepsy, or muscle weakness on one side of the body (hemiparesis).² The main concern with internal carotid agenesis is the increased risk for enlargement of the other blood vessels (aneurysm), which can occur in up to 67% of people with this condition (compared to 2-4% of individuals without this condition).³ It usually presents unilaterally and rarely has been reported bilateral.⁴ Associated defects seen are Aneurysms in the Circle of Willis Especially the Anterior circulation and Encephaloceles, intracranial hemorrhages^{5, 6} and Arachnoid cysts.^{7,8}

According to Padget⁹, the internal carotid artery originates from the dorsal aorta and the third aortic arch at approximately the 3-mm embryonic stage, but complete development does not occur earlier than the 16–18 mm stage (40 days). So far, there is no exact explanation for developmental anomalies of the internal carotid artery, but all variations are thought to occur because of insults to the developing embryo. Keen¹⁰ suggested that mechanical insults to the developing embryo such as excessive folding of the embryo to one side, pressure effects, or restriction by amniotic bands may cause unilateral absence of internal carotid artery.

A 35 year old female who had dementia reported to a medical OPD in SKIMS Soura. All her Lab Parameters were within normal range. A CT/ MRI was asked to be done by a Neurologist which revealed an Absent cavernous carotid artery. Neck Angiography revealing small right common carotid artery and no Internal carotid artery. MR Angiography revealed normal left petrous and Cavernous internal carotid. Right Middle cerebral artery and right anterior cerebral artery

Author ^α: Lecturer Anatomy SKIMS Medical College Bemina.

e-mail: zahidasool12@rediffmail.com


Author ^σ: Dental Surgeon, SKIMS Soura.

Author ^ρ: Prof. Anatomy Santosh Medical College Delhi.

Author ^ω: Ex Prof and Head Anatomy SKIMS Medical College Bemina.

are likely supplied by anterior communicating artery and posterior communicating artery. In addition bilateral

hyperintensity lesions were seen in heads of Caudate nuclei.




Magnetic resonance angiography of the cervical vessels showing agenesis of the right carotid artery

III. CONCLUSION

There are multiple variations in the cerebral vasculature but rarely do patients present with such a

rare anomaly. Such cases are usually detected incidentally or in some cases where a clinician would not suspect such a diagnosis in the first instance.



REFERENCES RÉFÉRENCES REFERENCIAS

1. Afifi, AK, Goersky JC, Menezes A, Smoker WR, Bell WE, and Jacoby CG. Cerebral Hemiatrophy, Hypoplasia of Internal Carotid Artery, and Intracranial Aneurysm: A Rare Association Occuring in an Infant. *Arch Neurol*. 1987.
2. Florio F, Balzano S, Nardella M, Strizzi V, Cammisa M, Bozzini V, Catapano G, and D'Angelo V. Congenital Absence of the Internal Carotid Artery. *Vardiovasc Intervent Radiol*. 1999.
3. Lee JH, Oh CW, Lee SH, and Han DH. Aplasia of the internal carotid artery. *Acta Neruochin*. 2003.
4. Rumboldt Z, Castillo M, Solander S. Bilateral congenital absence of the internal carotid artery. *Eur Radiol*. 2003;13 Suppl 6:L130–2.
5. Blustajn J, Netchine I, Frédy D, et al. Dysgenesis of the internal carotid artery associated with transsphenoidal encephalocele: a neural crest syndrome? *AJNR Am J Neuroradiol*. 1999;20: 1154–7.
6. Quint DJ, Silbergleit R, Young WC. Absence of the carotid canals at skull base CT. *Radiology*. 1992;182:477–81.
7. Kidooka M, Okada T, Handa J. Agenesis of the internal carotid artery – report of a case combined with arachnoid cyst in a child. *No To Shinkei*. 1992;44:371–5.
8. Owada Y, Sakuta Y. Bilateral carotid artery agenesis with corpus callosum hypogenesis – a case report. *No To Shinkei*. 1995;47:589–94.
9. Padget D. The development of the cranial arteries in the human embryo. *Contrib Embryol* 1948; 32:207 – 262.
10. Keen JA. Absence of both internal carotid arteries. *Clin Proc* 1945–1946; 4:588.



GLOBAL JOURNALS INC. (US) GUIDELINES HANDBOOK 2013

WWW.GLOBALJOURNALS.ORG

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards “FARSM” title to individuals. The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The “FARSM” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:



FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.



You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.


We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.





The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.

As FARSM, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.



The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.

The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website <https://associationofresearch.org> which will be helpful to upgrade the dignity.



The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.

The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.





The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.

The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.

The “MARSM” is a dignified ornament which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.



MARSM accrediting is an honor. It authenticates your research activities. After becoming MARSM, you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefits can be availed by you only for next three years from the date of certification.



MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.





We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.



Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.



AUXILIARY MEMBERSHIPS

Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as “Institutional Fellow of Open Association of Research Society” (IFOARS).

The “FARSC” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.



The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as “Institutional Board of Open Association of Research Society”-(IBOARS).

The Institute will be entitled to following benefits:



The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA). The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.



The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of “Open Association of Research Society, U.S.A (OARS)” so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



Journals Research
inducing researches

The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.



We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as “Institutional Fellow” and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.

The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:

Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.



Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.

Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.



We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth \$ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.



- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10% discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in-depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

//

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of “Difference of Opinion [if any]” among the Board members, our decision will be final and binding to everyone.

//



PROCESS OF SUBMISSION OF RESEARCH PAPER

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not convenient, and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



PREFERRED AUTHOR GUIDELINES

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.

Author Guidelines:

1. General,
2. Ethical Guidelines,
3. Submission of Manuscripts,
4. Manuscript's Category,
5. Structure and Format of Manuscript,
6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

- 1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.
- 2) Drafting the paper and revising it critically regarding important academic content.
- 3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5. STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also. Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

- (a) Title should be relevant and commensurate with the theme of the paper.
- (b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.
- (c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.
- (d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.
- (e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.
- (f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;
- (g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.
- (h) Brief Acknowledgements.
- (i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.



The Editorial Board reserves the right to make literary corrections and to make suggestions to improve brevity.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than $1.4 \times 10^{-3} \text{ m}^3$, or 4 mm somewhat than $4 \times 10^{-3} \text{ m}$. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art. A few tips for deciding as strategically as possible about keyword search:



- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.



Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org.



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.



12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.



Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

- Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

- Use standard writing style including articles ("a", "the," etc.)
- Keep on paying attention on the research topic of the paper
- Use paragraphs to split each significant point (excluding for the abstract)
- Align the primary line of each section
- Present your points in sound order
- Use present tense to report well accepted
- Use past tense to describe specific results
- Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives
- Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript-- must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.



- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently. You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.



ADMINISTRATION RULES LISTED BEFORE SUBMITTING YOUR RESEARCH PAPER TO GLOBAL JOURNALS INC. (US)

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.
- Do not give permission to anyone else to "PROOFREAD" your manuscript.
- **Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)**
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILATION)
BY GLOBAL JOURNALS INC. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	C-D	E-F
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring



INDEX

D

Diaphragm · 7, 10, 12, 15, 17, 19, 20
Duodenum · 28

E

Epilepsy · 32, 34, 36, 39, 40, 42, 44

G

Gastrojejunostomy · 29

L

Lymphoma · 2, 8

O

Omdurman · 2

R

Reynolds · 8, 9

S

Shimadzu · 12




save our planet



Global Journal of Medical Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals